

figure-drafts

Linda Herrera, Natalie Chen, June McNally

Set up

Packages

```
# general use and cleaning
library(tidyverse)
library(here)
library(janitor)
library(snakecase)

# wrapping axis labels
library(scales)

# reading in .xlsx files
library(readxl)

# arranging plots
library(patchwork)

# coloring
library(wesanderson)
```

Data

```
# taxonomic information
taxon_list <- read_csv(here("data", "taxon_list.csv"))

# aquatic invertebrates
aq_ins <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
```

```

        sheet = "Aquatic Insects")

# site information
sites <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
                  sheet = "sites")

# water quality information
water_quality <- read_xlsx(here("data", "Aquatic Sampling
↪ Data-2026-03-10.xlsx"),
                          sheet = "Water Quality",
                          na = c("", "n/a", "N/A", "over", "173+"))

```

Cleaning and Wrangling

Cleaning

sites object

```

# creating new object from sites
ncos_sites <- sites |>
  # filter to only include NCOS sites sampled four times a year
  filter(sector == "NCOS" & sampling_frequency == "Quarterly")

```

taxon_list object

```

# creating new clean object from taxon_list
taxon_list_clean <- taxon_list |>
  # converting taxon name to snake case (no spaces or capital letters,
  # only underscores)
  mutate(verbatim_name = to_any_case(verbatim_name, case = "snake")) |>
  select(verbatim_name, scientificName)

```

water_quality object

```

water_quality_clean <- water_quality |>
  clean_names() |>
  semi_join(ncos_sites, by = "site") |>
  unite("date_site", date, site, remove = TRUE) |>

```

```

group_by(date_site) |>
  summarize(
    med_ph      = median(p_h, na.rm = TRUE),
    med_sal     = median(salinity_ppt, na.rm = TRUE),
    med_do      = median(dissolved_oxygen_mg_l, na.rm = TRUE),
    med_temp    = median(temperature_c, na.rm = TRUE)) |> # <-- ADD
    ↪ THIS
ungroup() |>
filter(date_site != "2022-08-12_NVBR")

```

aq_ins object

```

# creating new clean object from aquatic inverts
aq_ins_clean <- aq_ins |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites occurring in ncos_sites
  semi_join(ncos_sites, by = c("site")) |>
  # renaming columns
  rename(date = date_on_vial) |>
  # select columns of interest
  select(date, sample_type, site,
    ostracod,
    copepod,
    cladocera) |>
  # filter to only include "filtered beaker" samples
  filter(sample_type %in% c("FB 250", "FB250")) |>
  # convert the data frame to long format
  pivot_longer(cols = ostracod:cladocera,
    names_to = "taxon",
    values_to = "abundance") |>
  # group by date, site, taxon
  group_by(date, site, taxon) |>
  # calculate average abundance per liter (sum abundance divided by 7.5
  ↪ liters)
  summarize(ave_lit = sum(abundance, na.rm = TRUE)/7.5) |>
  # ungroup data frame
  ungroup() |>
  # create a new column called `date_site` to join with water quality
  unite("date_site", date, site, remove = FALSE) |>
  # join with water quality
  left_join(water_quality_clean, by = c("date_site")) |>

```

```

# join with taxon list
left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>
# add full names of sites
mutate(site_full = case_when(
  site == "NVBR" ~ "North of Venoco Bridge",
  site == "NEC" ~ "South of Venoco Bridge",
  site == "NMC" ~ "Main Channel",
  site == "NPB" ~ "South of Phelps Bridge",
  site == "NPB1" ~ "North of Phelps Bridge",
  site == "NPB2" ~ "Phelps Road",
  site == "NWP" ~ "West Pond",
  site == "NDC" ~ "Devereux Creek"
)) |>
# setting factor levels for site
mutate(site_full = fct_reorder(site_full, med_sal, .fun = "median", .na_rm
  ↪ = TRUE),
  site_full = fct_rev(site_full)) |>

# ADD: log+1 transform of average abundance per liter
mutate(log_ave_lit = log(ave_lit + 1)) |> # <-- Fig
  ↪ 1, 2, 3, 4

# ADD: season derived from date
mutate(season = case_when(
  month(date) %in% c(3, 4, 5) ~ "Spring",
  month(date) %in% c(6, 7, 8) ~ "Summer",
  month(date) %in% c(9, 10, 11) ~ "Fall",
  month(date) %in% c(12, 1, 2) ~ "Winter"
),
season = factor(season, levels = c("Spring", "Summer", "Fall", "Winter")))
  ↪ |># <-- Fig 4
mutate(taxon = factor(taxon, levels = c("copepod", "ostracod",
  ↪ "cladocera")))

```

```

# colors for taxa
taxon_cols <- c(
  "ostracod" = "orange2",
  "copepod" = "tomato3",
  "cladocera" = "turquoise4")

# setting common theme elements
theme_set(

```

```
theme_classic() +
  theme(text = element_text(size = 12),
        plot.title.position = "plot",
        legend.position = "none"))
```

Figure 1

Figure 2 - Mean and Distribution of Invertebrate Abundance (log +1/L) for Three Tax

```
jitter <- ggplot(data = aq_ins_clean,
                 aes(x = taxon,
                     y = log_ave_lit,
                     color = taxon)) +

  geom_jitter(height = 0,
              alpha = 1,
              width = 0.2,
              shape = 21) +
  # second layer: points to represent medians
  stat_summary(geom = "point",
               fun = "mean",
               size = 4,
               color = "black",
               shape = 18) + # 18 = filled diamond

  stat_summary(geom = "errorbar",
               fun.data = mean_se,
               width = 0.1,
               color = "black") +

  scale_color_manual(values = taxon_cols) +
  labs(x = "Taxon",
       y = "log + 1 transformed average abundance/L",
       title = " Mean and Distribution of Invertebrate Abundance (log +1/L)
               for Three Tax")

jitter
```

Mean and Distribution of Invertebrate Abundance (log +1/L) for Three Tax




Figure 3

Figure 4

```
# create new object to store figure 4
figure4 <- ggplot(data = aq_ins_clean,
                  mapping = aes(x = med_temp,
                                y = log_ave_lit,
                                color = taxon)) +

  geom_jitter(height = 0,
              shape = 21) +

  geom_line() +

  facet_wrap(~season,
             scales = "free") +
  labs(title = " Seasonal Patterns Between Water Temperature and Log + 1
    Abundance for Three Dominant Taxa",
       x = "Average Temperature (°C)",
       y = "log + 1 Transformed Average Abundance/L",
```

```

color = "Taxon") +

scale_color_manual(values = taxon_cols) +

theme(legend.position = "right")

```

figure4

Seasonal Patterns Between Water Temperature and Log + 1 Abundance for Three Dominant Taxa

