

# Individual Assignment 5

Natalie Chen

2026-05-04

## Set up

### Packages

```
#general use
library(here)

library(tidyverse)

library(janitor)

library(snakecase)

# packages for visualization customization

library(scales)
library(palettter)
library(ggthemes)

# reading in .xlsx files
library(readxl)

# visualizing missing data
library(naniar)

# arranging plots
library(patchwork)
```

## Data

```
# taxonomic information
taxon_list <- read_csv(here("data", "taxon_list.csv"))

# aquatic invertebrates
aq_ins <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Aquatic Insects")

# site information
sites <- read_xlsx(here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "sites")

# water quality information
water_quality <- read_xlsx(here("data", "Aquatic Sampling
  ↪ Data-2026-03-10.xlsx"),
  sheet = "Water Quality",
  na = c("", "n/a", "N/A", "over", "173+"))
```

## copied from class

```
# creating new object from sites
ncos_sites <- sites |>

  # filter to only include NCOS quarterly sampling sites
  filter(sector == "NCOS" & sampling_frequency == "Quarterly")

water_quality_clean <- water_quality |>

  # cleaning column names
  clean_names() |>

  # filtering join: only include sites in ncos_sites
  # keep all the rows on the left that match up with rows on the right
  semi_join(ncos_sites, by = "site") |>

  # combine date and site together and remove those indiv. columns
  unite("date_site", date, site, remove = TRUE) |>

  # group by date_site
  group_by(date_site) |>
```

```

# create new columns that show median of each variable at each date_site
# because there are multiple samples at the same site at different depths
summarize(med_ph = median(p_h, na.rm = TRUE),
          med_sal = median(salinity_ppt, na.rm = TRUE),
          med_do = median(dissolved_oxygen_mg_l, na.rm = TRUE)) |>

# ungroup
ungroup() |>

# filter out the salinity outlier
filter(date_site != "2022-08-12_NVBR")

```

```

# creating new object from taxon_list
taxon_list_clean <- taxon_list |>

# converted taxon name values to snake case (lowercase)
mutate(verbatim_name = to_any_case(verbatim_name,
                                   case = "snake"))

```

```

# creating new clean aquatic inverts object from aquatic inverts
aq_ins_clean <- aq_ins |>

# cleaning column names
clean_names() |>

# filtering join: only include quarterly NCOS surveys from ncos_sites
semi_join(ncos_sites, by = "site") |>

# renaming column to just date
rename(date = date_on_vial) |>

# select columns of interest
select(date, sample_type, site,

       # 9 most abundant taxon
       ostracod,
       copepod,
       hemiptera_corixidae_boatman,
       cladocera,
       nematode,
       diptera_ceratopogonidae,

```

```

    annelida_oligochaete,
    diptera_chironomid,
    annelida_polychaete) |>

# filter to only include filtered beaker samples
filter(sample_type %in% c("FB250", "FB 250")) |>

pivot_longer(
  # cols = columns to make longer
  cols = ostracod:annelida_polychaete,
  # new column for the old column names
  names_to = "taxon",
  # new column for the values in each of the old columns
  values_to = "abundance"
) |>

# group by the date, site, and taxon
group_by(date, site, taxon) |>

# calculate average abundance per liter (sum abundance divided by 7.5
  ↳ liters)
summarize(avg_per_lit = sum(abundance, na.rm = TRUE)/7.5) |>

# ungroup
ungroup() |>

# create a new column called `date_site` to join with water quality
# don't remove the original columns
unite("date_site", date, site, remove = FALSE) |>

# join with water quality clean
# wqc has `date_site` column as well
left_join(water_quality_clean, by = "date_site") |>

# join with taxonomic information from taxon list
# verbatim_name column in taxon_list_clean matches with taxon column in
  ↳ aq_ins_clean
left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>

# fill in full site names
mutate(site_full = case_when(
  site == "NVBR" ~ "North of Venoco Bridge",

```

```

    site == "NEC" ~ "East Channel",
    site == "NMC" ~ "Main Channel",
    site == "NPB" ~ "South of Phelps Bridge",
    site == "NPB1" ~ "North of Phelps Bridge",
    site == "NPB2" ~ "Phelps Road",
    site == "NWP" ~ "West Pond",
    site == "NDC" ~ "Devereux Creek"
  )) |>

# setting factor levels for full site name by median salinity
mutate(site_full = fct_reorder(site_full, med_sal, .fun = "median", .na_rm
  ↪   = TRUE),
       site_full = fct_rev(site_full))

salinity_plot <- ggplot(data = aq_ins_clean,
                        # x-axis: full site name
                        # y-axis: median salinity
                        mapping = aes(x = site_full,
                                      y = med_sal)) +
# points representing median salinity at each survey
geom_jitter(height = 0,
            shape = 21) +

stat_summary(geom = "point",
            fun = median,
            color = "red",
            size = 2,) +


# wrap x-axis labels
scale_x_discrete(labels = label_wrap(width = 10)) +

# labeling titles and axes
labs(x = "Site",
     y = "Median salinity (ppt)",
     title = "Site Salinity") +

# change theme
theme_bw()

# displaying the object
salinity_plot

```



```
copepods <- aq_ins_clean |>
```

```
  filter(taxon == "copepod") |>
```

```
  mutate(log_avg_per_lit = log(avg_per_lit + 1))
```

```
# creating object to store plot
```

```
copepod_plot <- ggplot(data = copepods,
```

```
  # x-axis: full site name
```

```
  # y-axis: log transformed avg count/L
```

```
  mapping = aes(x = site_full,  
                y = log_avg_per_lit)) +
```

```
  # jitter to represent salinity
```

```
  geom_jitter(height = 0,  
              width = 0.1,  
              shape = 21) +
```

```
# points to represent median copepod abundance
```

```
  stat_summary(geom = "point",  
              fun = median,  
              color = "blue",  
              size = 2) +
```

```
  # wrapping x axis labels
```

```
  scale_x_discrete(labels = label_wrap(width = 10)) +
```

```
  # relabelling axes
```


```

labs(x = "Site",
     y = "log + 1 transformed averag abundance/L",
     title = "Copepod Abundance") +

# different theme
theme_bw()

# display object
copepod_plot

```



## 1. Visualizing differences across sites in major taxon abundance

### a. Plan your figure

- x-axis: site\_full
- y-axis: average abundance of ostracod per liter
- geometry to represent average abundance/L: `geom_jitter()`
- geometry to represent medians: `stat_summary()`

## b. Wrangle your data

```
# creating new object from aquatic insects
ostracod <- aq_ins_clean |>

# filter to only include ostracod
filter(taxon == "ostracod") |>

# calculate the average of ostracod abundance per lit
mutate(ostracod_log_avg_per_lit = log(avg_per_lit + 1))

# display 10 random rows

slice_sample(ostracod, n = 10)
```

# A tibble: 10 x 24

|    | date_site<br><chr> | date<br><dtm>       | site<br><chr> | taxon<br><chr> | avg_per_lit<br><dbl> | med_ph<br><dbl> | med_sal<br><dbl> | med_do<br><dbl> |
|----|--------------------|---------------------|---------------|----------------|----------------------|-----------------|------------------|-----------------|
| 1  | 2023-02-12~        | 2023-02-12 00:00:00 | NEC           | ostr~          | 2.13                 | NA              | NA               | NA              |
| 2  | 2023-05-04~        | 2023-05-04 00:00:00 | NEC           | ostr~          | 0.667                | 8.05            | NA               | 8.55            |
| 3  | 2022-08-15~        | 2022-08-15 00:00:00 | NMC           | ostr~          | 0.133                | 7.61            | NA               | NA              |
| 4  | 2020-01-21~        | 2020-01-21 00:00:00 | NDC           | ostr~          | 2.4                  | 6.53            | 0.3              | 7.6             |
| 5  | 2023-09-12~        | 2023-09-12 00:00:00 | NVBR          | ostr~          | 0                    | 9.14            | NA               | 20.3            |
| 6  | 2023-02-28~        | 2023-02-28 00:00:00 | NPB           | ostr~          | 2.53                 | NA              | NA               | NA              |
| 7  | 2023-05-06~        | 2023-05-06 00:00:00 | NDC           | ostr~          | 56                   | 10.1            | NA               | 13.1            |
| 8  | 2023-03-03~        | 2023-03-03 00:00:00 | NPB2          | ostr~          | 0.133                | 7.71            | 1.98             | 9.59            |
| 9  | 2022-05-12~        | 2022-05-12 00:00:00 | NPB2          | ostr~          | 1198                 | NA              | NA               | NA              |
| 10 | 2022-05-05~        | 2022-05-05 00:00:00 | NEC           | ostr~          | 0.267                | 9.45            | 63.8             | 7.3             |

# i 16 more variables: taxonRank <chr>, scientificName <chr>, kingdom <chr>,  
# Phylum <chr>, Class <chr>, Subclass <chr>, Superorder <chr>, Order <chr>,  
# Infraorder <chr>, Family <chr>, Subfamily <chr>, Tribe <chr>, Genus <chr>,  
# comments <chr>, site\_full <fct>, ostracod\_log\_avg\_per\_lit <dbl>

## c. Code your figure

```
# base layer: create an object that stores the plot
ostracod_plot <- ggplot(data = ostracod,
  # x = site
  # y = log + 1 transformed average abundance/L
```

```

        mapping = aes(x = site_full,
                      y = ostracod_log_avg_per_lit)) +


# first layer: geometry jitter points to represent each sample
geom_jitter(height = 0,
            width = 0.1,
            shape = 21) +
# points to represent median abundance
stat_summary(geom = "point",
            fun = median,
            color = "green",
            size = 2) +
# wrapping x axis labels
scale_x_discrete(labels = label_wrap(width = 10)) +

# labeling axes and title
labs(title = "Ostracod Abundance",
     x = "Site",
     y = "log + 1 transformed average abundance/L") +

# changed theme
theme_bw()


# display plot
ostracod_plot

```



#### d. Create a multipanel figure

```
salinity_plot / (copepod_plot | ostracod_plot)
```



**e. Interpret your figure**

What are the differences between sites in copepod and ostracod abundance?

- Average copepod abundance/L was highest at East Channel and Devereux Creek at respective transformed values of around 4 and 3, and moderate at a value of around 2 at North of Venoco Bridge, South of Phelps Bridge, and Phelps Road. The highest average ostracod abundance/L was moderate around a transformed value of 2 at South of Phelps Bridge and also Devereux Creek, and low at pretty much every other site.

**How does ostracod abundance relate (or not relate) to site-level differences in salinity?**

- Ostracod abundance does not correlate to salinity site-level differences in salinity because the plots do not show the same type of trend for the salinity plot and for the ostracod abundance plot.

## **2. Visualizing total abundance as a function of salinity**

**a. Plan your figure**

**b. Wrangle your data**

**c. Code your figure**

**d. Interpret your figure**