

homework 3

Ngoc Anh Pham

September 2021

Ex1 : $t = y(x, w) + \text{noise}$
 $\Rightarrow w = (X^T X)^{-1} X^T t$

We have: $t = y(x, w) + \text{noise} = N(y(x, w), \beta^{-1})$ Precision parameter $\beta = \frac{1}{\sigma^2}$

$$p(t|x, w, \beta) = N(t|y(x, w), \beta^{-1})$$

The likelihood function:

$$p(t|x, w, \beta) = \prod_{n=1}^N N(t_n|y(x_n, w), \beta^{-1})$$

It is convenient to maximize the logarithm of the likelihood function

$$\begin{aligned} \log p(t|x, w, \beta) &= \sum_{n=1}^N \log (N(t_n|y(x_n, w), \beta^{-1})) \\ &= -\frac{\beta}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2 + \frac{N}{2} \log \beta - \frac{N}{2} \log(2\pi) \\ \max \log p(t|x, w, \beta) &= -\max \frac{-\beta}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2 \\ &= \min \frac{1}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2 \end{aligned}$$

We minimize $P = \frac{1}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2$ to find w . Suppose:

$$X = [x_1 \ x_2 \ \dots \ x_n], w = w_0 w_1$$

$$\Rightarrow P = \|Xw - t\|_2^2$$

$$\nabla P = 2X^T(Xw - t) = 2X^T Xw - 2X^T t$$

Setting this gradient to zero, we have:

$$X^T Xw - X^T t = 0$$

$$\text{so : } w = (X^T X)^{-1} X^T t$$

Ex4 : Prove that $X^T X$ is invertible when X is full rank

Suppose $X^T v = 0$. $\Rightarrow X X^T v = 0$

Conversely, suppose $X X^T v = 0$.

$\Rightarrow v^T X X^T v = 0$, so that $(X^T v)^T (X^T v) = 0$.

This implies $X^T v = 0$.

Hence, we have proved that $X^T v = 0$ if and only if v is in the nullspace of $X^T X$.

But $X^T v = 0$ and $v \neq 0$ if and only if X has linearly dependent rows.

Thus, $X^T X$ is invertible if and only if X has full row rank.