

homework 3

Ngoc Anh Pham

September 2021

Ex1 : $t = y(x, w) + noise$

$$\Rightarrow w = (X^T X)^{-1} X^T t$$

We have:

$$t = y(x, w) + noise = N(y(x, w), \beta^{-1})$$

Precision parameter $\beta = \frac{1}{\sigma^2}$

$$p(t|x, w, \beta) = N(t|y(x, w), \beta^{-1})$$

The likelihood function:

$$p(t|x, w, \beta) = \prod_{n=1}^N N(t_n|y(x_n, w), \beta^{-1})$$

It is convenient to maximize the logarithm of the likelihood function

$$\begin{aligned} \log p(t|x, w, \beta) &= \sum_{n=1}^N \log (N(t_n|y(x_n, w), \beta^{-1})) \\ &= \frac{-\beta}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2 + \frac{N}{2} \log \beta - \frac{N}{2} \log(2\pi) \\ \max \log p(t|x, w, \beta) &= -\max \frac{-\beta}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2 \\ &= \min \frac{1}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2 \end{aligned}$$

We minimize $P = \frac{1}{2} \sum_{n=1}^N (y(x_n, w) - t_n)^2$ to find w . Suppose:

$$X = [x_1, x_2, \dots, x_n], w = [w_0, w_1]$$

$$\Rightarrow P = \|Xw - t\|_2^2$$

$$\nabla P = 2X^T(Xw - t) = 2X^TXw - 2X^Tt$$

Setting this gradient to zero, we have:

$$X^TXw - X^Tt = 0$$

$$\text{so : } w = (X^TX)^{-1}X^Tt$$

Ex4 : Prove that X^TX is invertible when X is full rank

Suppose $X^Tv = 0$. $\Rightarrow XX^Tv = 0$
 Conversely, suppose $XX^Tv = 0$.
 $\Rightarrow v^TXX^Tv = 0$, so that $(X^Tv)^T(X^Tv) = 0$.
 This implies $X^Tv = 0$.
 Hence, we have proved that $X^Tv = 0$ if and only if v is in the nullspace of X^TX .
 But $X^Tv = 0$ and $v \neq 0$ if and only if X has linearly dependent rows.
 Thus, X^TX is invertible if and only if X has full row rank.