

# BRANCHIO-OTO-RENAL SYNDROME: A CASE REPORT & REVIEW OF LITERATURE

Dr. Prithvi Das<sup>1\*</sup>, Dr. Somnath Saha<sup>2</sup>

<sup>1</sup> Senior Resident, Department of ENT, Head & Neck Surgery, Calcutta National Medical College & Hospital, Kolkata.

<sup>2</sup> Professor & Head, Department of ENT, Head & Neck Surgery, Calcutta National Medical College & Hospital, Kolkata.

\*Correspondence:

Dr. Prithvi Das

Email: prithvidas89@gmail.com

## ABSTRACT:

**Background:** Branchio-oto-renal (BOR) syndrome is a rare autosomal dominant disorder characterized by syndromic association of branchial arch anomalies, otologic malformations and renal anomalies. Very few articles on this syndrome have been published in journals related to Otorhinolaryngology.

**Case Report:** A 18 years old male patient, presented with a right sided intermittently discharging neck sinus at the junction of middle 1/3rd and lower 1/3rd of anterior border of sternocleidomastoid muscle along with bilateral pinna deformity and preauricular sinus. Fistulogram revealed a fistulous tract communicating with the right tonsillar bed. NCCT temporal bone showed marked hypoplasia of bilateral external auditory canal, dysplastic middle ear ossicles and microtia. NCCT KUB showed a hypoplastic right kidney and malrotated left kidney although his renal function test were within normal limits. In the presence of 3 major symptoms, branchial fistula, bilateral middle ear malformations and renal structure abnormality this case was clinically diagnosed as branchio-oto-renal syndrome. The patient was treated for his active symptoms hence his fistulous tract was excised after initial medical management.

**Conclusion:** The BOR syndrome is an infrequent but a well described entity that combines branchial arch anomalies, otologic malformations and renal anomalies. The BOR syndrome should be included in the differential diagnosis of deafness and chronic renal failure in childhood and adolescence.

**Key words:** Branchio-Oto-Renal Syndrome, fistula, branchial, renal malrotation, preauricular sinus, renal hypoplasia.

## INTRODUCTION:

Branchio-oto-renal (BOR) syndrome is a rare autosomal dominant disorder with prevalence of approximately 1 in 40,000 new born infants and in about 2% of children with profound deafness<sup>1,2</sup>. Branchio-oto-renal (BOR) syndrome was first described by Melnick et al<sup>3</sup> in 1975 followed by Fraser et al<sup>4</sup>, thus giving the name Melnick-Fraser syndrome.

It is characterized by branchial arch anomalies (branchial clefts, fistula, cysts), hearing impairment (malformation of auricle with preauricular sinuses, conductive, sensorineural, or mixed hearing impairment), and renal malformations (urinary tract malformation, renal hypoplasia or agenesis, renal hypoplasia, renal cysts)<sup>5</sup>. Patients usually present with hearing impairment and discharging branchial fistula.

Such patients should be further investigated for BOR syndrome and thus control its progress.

Phenotypic presentation of BOR syndrome is extremely variable<sup>6</sup>. The combination of deafness with chronic renal failure may be confused with the Alport syndrome. Anterior lenticonus, a conical protrusion of the central portion of lens in to the anterior chamber is pathognomonic for Alport's syndrome. Moreover deafness in the Alport's syndrome manifests at a later age<sup>7,12</sup>.

## CASE REPORT:

An 18 years old male patient presented to the Otorhinolaryngology out-patient department with the chief complaints of right sided intermittently discharging neck sinus since birth.

He was born of non-consanguineous marriage and there was no history of hearing deficits, renal disease and similar discharging sinuses in the parents or in his only sibling. His birth history was insignificant. On examination the patient was noted to have a discharging fistulous tract over right side of the neck at the junction of middle 1/3rd and lower 1/3rd on the anterior border of sternocleidomastoid muscle. The patient also had bilateral pinna deformity along with bilateral preauricular sinus.



Fig 1: showing pinna deformity and discharging sinus over right side of neck



Fig 2: Showing bilateral pinna deformity  
Facial nerve examination was within normal limits on both side and audiological assessment revealed bilateral severe mixed sensorineural hearing loss. On investigating further, following were noted: Fistulogram shows fluid contrast through fistulous tract communicating with the right tonsillar bed.




Fig 3: fistulogram

NCCT temporal bone shows marked hypoplasia of bilateral external auditory canal, dysplastic middle ear ossicles and bilateral microtia. Scans also revealed opacification of middle ear cavity on both side.



Fig 4: NCCT Temporal Bone




Fig 5 Right side

Fig 6 Left side

Although the patient had no complaints related to renal function, but on the basis of suspicion for BOR syndrome a NCCT KUB was requested and it revealed a small right kidney and malrotation of left kidney.



He was then referred to a nephrologist for his renal anomaly and subsequent regular follow up.



Fig 7 CT Abdomen




Fig 8: CT Abdomen

As a part of his management work up the patient was treated only for its active symptoms.

Hence the patient underwent complete excision of the fistulous tract after initial stabilization and medical management. He was advised hearing aids, regular hearing assessment, and periodic renal function monitoring.

In the light of 3 major symptoms, branchial fistula, bilateral middle ear malformations and renal structure abnormality this case was clinically diagnosed as Branchio-oto-renal syndrome.

## DISCUSSION:

The BOR syndrome is a rare but well defined constellation of branchial arch anomalies (branchial fistulas, branchial cysts), external ear malformations with

hearing loss and renal hypoplasia and dysplasia. It has an autosomal dominant transmission pattern with variable clinical expression<sup>6</sup>

Chang et al. developed the diagnostic criteria of BOR syndrome in 2004<sup>8,9</sup>. These include major criteria such as second branchial arch anomalies, hearing loss, preauricular pits, auricular deformity and renal anomalies, and minor criteria such as external auditory canal anomalies, middle ear anomalies, inner ear anomalies, preauricular tags and other symptoms such as facial asymmetry and palate abnormalities.

In patients with a family history, any single major criterion is sufficient for diagnosis of BOR syndrome. Without any family history, three major criteria or two major and two minor criteria are needed to make a confident diagnosis.

EYA1 (8q13.3), the human homologue of the *Drosophila* eyes absent gene, is the most frequent causative gene for BOR syndrome<sup>10</sup>. It expresses very early, between 4th & 6th weeks of human embryogenesis<sup>11</sup>. Deafness relates to abnormalities in the three ossicles of the middle ear derived from the first and second branchial arches, while the branchial fistulae relates to second, third and fourth arches. EYA1 gene is strongly expressed in the human embryonic kidney and in BOR syndrome there is fault between the ureteric bud and metanephric mesenchymal mass as the ureteric bud branches into renal parenchyma, resulting in renal anomalies<sup>12</sup>. Mutations in SIX1 (14q23.1) and SIX5 (19q13.32) have been reported less frequently<sup>13</sup>. SIX1 interacts with EYA1 in the development of various organs<sup>14</sup>. SIX5 has a high degree of homology to SIX1 and directly interacts with EYA1<sup>15</sup>.

Clinical features are highly variable. The most common presenting symptom is deafness (90%) which can be sensorineural or conductive but is mostly mixed (50%)<sup>16</sup>. Pre-auricular pits can be the presenting features in over 70% of the cases and sometimes can be the only external ear finding as seen in our patient, while around 50% of patients have external ear anomalies in the form of microtia to small lop or cupped ears with over folded superior helices<sup>17</sup>. Middle ear anomalies include ossicular malformations and inner ear anomalies include cochlear hypoplasia or dysplasia<sup>18</sup>. Renal anomalies include renal dysplasia and agenesis which can lead to end-stage renal disease<sup>19</sup>.

## CONCLUSION:

BOR syndrome is a rare but well described clinical entity. It combines branchial anomalies with otologic and renal malformations. Most individuals with BOR syndrome do not have a life-threatening condition, and, in many families, it is not uncommon for the disease to go undiagnosed until the birth of a child with severe

manifestations of the BOR phenotype. This is unfortunate, as recognition of the hallmark features of BOR syndrome could ensure that affected persons receive appropriate medical information and care. Integral elements of medical care include audiologic, otologic, head & neck, urologic, and genetic evaluation. Patients with BOR syndrome who receive adequate treatment can lead normal, productive lives. It is therefore essential to provide early diagnosis including the relevant genetic tests, although further studies are needed to unearth the molecular mechanisms and undiscovered causative genes responsible for BOR syndrome.

## REFERENCES:

1. Fraser FC, Sproute JR, Halal F. Frequency of the branchio-otorenal (BOR) syndrome in children with profound hearing loss. *Am J Med Genet* 1980; 7:341-9.
2. Jones KL. Melnick-Fraser syndrome. In: Jones KL, Jones MC, del Campo M, editors. *Smith's recognizable pattern of human malformation*. 4th ed. Philadelphia: W.B. Saunders's Co. 1988.p. 206.
3. Melnick M, Bixler D, Silk K, Yune H, Nance WE. Autosomal dominant branchio-oto-renal dysplasia: two distinct autosomal dominant disorders. *Clin Genet* 1975; 13:425-42.
4. Fraser FC, Ling D, Clogg D, Nogrady B. Genetic aspects of the BOR syndrome: branchial fistulas, ear pits, hearing loss and renal anomalies. *Am J Med Genet* 1978; 2:241-52.
5. Emrah FC, Hatice K, Fatih A, Guleser S, Basak NG, Saliha S. From a branchial fistula to a branchiootorenal syndrome: A case report and review of the literature. *J Pediatr Surg* 2009;44:623-5.
6. Melnick M, Bixler D, Nance W, Dilk K, Yune H. Familial Branchio-otorenal dysplasia: a new addition to the branchial arch syndrome. *Clin Genetics* 1976;9:25-34.
7. Misra M, Nolph KD. Renal failure and deafness: branchio-oto-renal syndrome. *Am J Kidney Dis* 1998; 32: 334-337
8. Smith RJH. Branchiootorenal spectrum disorders. In: Pagon RA, Adam MP, Bird TD et al. (eds). *GeneReviews™* (Internet). University of Washington, Seattle, WA, 1999. Mar 19 [Updated 2013 Jun 20; Cited 23 February 2014.]; Available from URL: <http://www.ncbi.nlm.nih.gov/books/NBK1380/>.
9. Chang EH, Menezes M, Meyer NC et al. Branchio-oto-renal syndrome: The mutation spectrum in EYA1 and its phenotypic consequences. *Hum. Mutat.* 2004; 23: 582-9.
10. Abdelhak S, Kalatzis V, Heilig R et al. A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal (BOR) syndrome and identifies a novel gene family. *Nat. Genet.* 1997; 15: 157-64.
11. Kemperman MH, Stinckens C, Kumar S, Hygen PLM, Joosten FB, Cremers CW. The Branchio-oto-renal syndrome. *Adv Otorhinolaryngol* 2002;61:192-200.
12. Kameswaran M, Anand RS, Murli S, Raghunandhan S, Karthikeyan. Cochlear implantation in Branchio-oto-renal syndrome – A surgical challenge. *Indian J Otolaryngol Head & neck Surg* 2007;59:280-3.
13. Krug P, Morinière V, Marlin S, Koubi V, Gabriel HD, Colin E, et al. Mutation screening of the EYA1, SIX1, and SIX5 genes in a large cohort of patients harboring branchio-oto-renal syndrome calls into question the pathogenic role of SIX5 mutations. *Hum Mutat* 2011; 32:183-90.
14. Mutsuddi M, Chaffee B, Cassidy J, Silver SJ, Tootle TL, Rebay I. Using *Drosophila* to decipher how mutations associated with human branchio-oto-renal syndrome and optical defects compromise the protein tyrosine phosphatase and transcriptional functions of eyes absent. *Genetics* 2005; 170: 687-95.
15. Hoskins BE, Cramer CH, Silvius D et al. Transcription factor SIX5 is mutated in patients with branchio-oto-renal syndrome. *Am. J. Hum. Mutat.* 2007; 80: 800-4.
16. Smith RJ, Schwartz C. Branchio-oto-renal syndrome. *J Common Disord* 1998; 31:411.
17. Kemperman MH, Stinckens C, Kumar S, Hygen PLM, Joosten FB, Cremers CW. Progressive fluctuant hearing loss, enlarged vestibular aqueduct and cochlear hypoplasia in branchio-otorenal syndrome. *Otol Neurotol* 2001; 22:637-43.
18. Ceruti S, Stinckens C, Cremers CW, Casselman JW. Temporal bone anomalies in the branchio-oto-renal syndrome: detailed computed tomographic and magnetic resonance imaging findings. *Otol Neurotol* 2002; 23:200-7.
19. Peterson R, Agarwal I, Sinha R. End-stage renal disease with deafness in an adolescent child. *Indian J Nephrol* 2011; 21: 293-4.
20. Smith RJH. Branchiootorenal Spectrum Disorder. 1999 Mar 19 [Updated 2018 Sep 6]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1380/>