



Trøndelagkonkurranse 2026 - Level II - Solutions

First round - 26.09

1st part


Choose and solve only one of the following problems:

Problem 1

The figure shows an equilateral triangle of side 1, and three internal squares tangent to the base of the triangle and to each other as shown. The respective sides of the squares are a , $2a$ and $3a$, and as shown, the smallest and largest squares each touch one additional side with a vertex. Knowing that a can be expressed as

$$a = \frac{A + B\sqrt{3}}{C},$$

with $A, B, C \in \mathbb{N}$ and the expression is as reduced as possible, determine the value of $A + B + C$.



Problem 2

Bianca has a calculator with two special buttons A and B . Given a certain value x , A takes x into $\frac{1}{x}$, while B takes x into $1 - x$.

If she starts with some number x and presses the special buttons in the order

$$A, B, A, B, A, B, \dots, A, B$$

starting with A and alternating each time such that each button is pressed 1013 times, which value will she obtain at the end?

Problem 1

Let p and q be the small segments of the base of the triangle at the left and right of the squares respectively, hence, if we add them to the sides of the squares we add up to 1, namely

$$p + a + 2a + 3a + q = 1. \quad (1)$$

Notice that p and q are basis of half equilateral triangles. On the left, the one with base p has altitude a and on the right, the one with base q has altitude $3a$. Using the Pythagorean theorem, we find that the ratio between the base and the altitude of half of an equilateral triangle is $1 : \sqrt{3}$, thus we can determine that

$$p = \frac{a}{\sqrt{3}} \quad \text{and} \quad q = a\sqrt{3}.$$

Putting this back into (1) we have

$$\frac{a}{\sqrt{3}} + a + 2a + 3a + a\sqrt{3} = 1, \quad (2)$$

which after grouping and simplifying leads to

$$\begin{aligned} \frac{a}{\sqrt{3}} + 6a + a\sqrt{3} &= 1 \\ a + 6a\sqrt{3} + 3a &= \sqrt{3} \\ a(6\sqrt{3} + 4) &= \sqrt{3} \\ a &= \frac{\sqrt{3}}{6\sqrt{3} + 4} \\ a &= \frac{\sqrt{3}(6\sqrt{3} - 4)}{6^2 \cdot 3 - 4^2} \\ a &= \frac{18 - 4\sqrt{3}}{92} \\ a &= \frac{9 - 2\sqrt{3}}{46}. \end{aligned}$$

Finally, we have that

$$A + B + C = 9 - 2 + 46 = 53.$$

Problem 2

After computing some iterations of A and B for some value x , we find that

$$x \xrightarrow{A} \frac{1}{x} \xrightarrow{B} \frac{x-1}{x} \xrightarrow{A} \frac{x}{x-1} \xrightarrow{B} \frac{1}{1-x} \xrightarrow{A} 1-x \xrightarrow{B} x$$

which means that the sequence A, B, A, B, A, B , takes x into x .

With this in mind, since the original sequence has length 2026, it suffices to find $l < 5$ such that $2026 = 6k + l$ for some $k \in \mathbb{N}$. Notice that $2026 = 2022 + 4$ and 2022 is divisible by 2 and

3, hence it is divisible by 6. This tells us that, since every sequence of A, B, A, B, A, B leaves x unchanged, applying A and B alternating 2026 times is equivalent to just apply

$$A, B, A, B,$$

which, from the above diagram, we can see it will yield $\frac{1}{1-x}$, so this is the number Bianca will obtain.


$$\sqrt{T}$$

Trøndelagkonkurranse 2026 - Level II - Solutions

First round - 26.09

2nd part

Bianca proposes the following problem to Nils, she constructs a big triangular array of numbers as shown below



such that the possible values of a, b, c , and so on are $-1, 0$ and 1 . She first chooses one value for a , and then randomly assigns values making sure that no two adjacent values are equal.

Task 1: If the bottom of the array has 101 numbers, determine the maximum value of the sum of all the values in the array.

Task 2: If the bottom of the array has 2025 numbers, show that the sum of all the values in the array is independent of the choice of a .

Nils, would also like to give Bianca a similar problem, either using arrays of numbers or sums of integers or anything interesting he can think of.

Task 3: Help Nils create a nice problem he can give to Bianca.

See level I for solution.