INITIAL VISIT PACKET

UNIFORM DATA SET (UDS) VERSION 4.0



Form B3: Unified Parkinson's Disease Rating Scale (UPDRS¹) - Motor Exam

ADRC name:	Participant ID:	Form date: / /
Visit #:	Examiner's initials:	
	s form is to be completed by the clinician or other trained s, see UDS Coding Guidebook for Initial Visit Packet, Form	
- I	the clinician completes the UPDRS examination an ox is checked, all items will default to 0 in the database	
1. Speech	Check the most appropriate box: O Normal Slight loss of expression, diction and/or volume Monotone, slurred but understandable; moderately impaired.	☐ 3 Marked impairment, difficult to understand. ☐ 4 Unintelligible ☐ 8 Untestable (SPECIFY):
2. Facial expression	Check the most appropriate box: 0 Normal 1 Minimal hypomimia, could be normal "poker face" 2 Slight but definitely abnormal diminution of facial expression	☐ 3 Moderate hypomimia; lips parted some of the time ☐ 4 Masked or fixed facies with severe or complete loss of facial expression; lips parted ¼ inches or more ☐ 8 Untestable (SPECIFY):
3. Tremor at rest		
3a. Face, lips, chin	Check the most appropriate box: ☐ 0 Absent ☐ 1 Slight and infrequently present ☐ 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present	☐ 3 Moderate in amplitude and present most of the time ☐ 4 Marked in amplitude and present most of the time ☐ 8 Untestable (SPECIFY):
3b. Right hand	Check the most appropriate box: 0 Absent 1 Slight and infrequently present 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present	☐ 3 Moderate in amplitude and present most of the time ☐ 4 Marked in amplitude and present most of the time ☐ 8 Untestable (SPECIFY):

¹ Fahn S, Elton RL, UPDRS Development Committee. The Unified Parkinson's Disease Rating Scale. In Fahn S, Marsden CD, Calne DB, Goldstein M, eds. Recent developments in Parkinson's disease, Vol. 2. Florham Park, NJ: Macmillan Healthcare Information, 1987:153-163, 293-304. Reproduced by permission of the author.

	continued		
Check the most appropriate box: 0 Absent 1 Slight and infrequently present 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present	☐ 3 Moderate in amplitude and present most of the time ☐ 4 Marked in amplitude and present most of the time ☐ 8 Untestable (SPECIFY):		
Check the most appropriate box: 0 Absent 1 Slight and infrequently present 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present	☐ 3 Moderate in amplitude and present most of the time ☐ 4 Marked in amplitude and present most of the time ☐ 8 Untestable (SPECIFY):		
Check the most appropriate box: 0 Absent 1 Slight and infrequently present 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present	☐ 3 Moderate in amplitude and present most of the time ☐ 4 Marked in amplitude and present most of the time ☐ 8 Untestable (SPECIFY):		
l tremor of hands			
Check the most appropriate box: 0 Absent 1 Slight; present with action 2 Moderate in amplitude, present with action	☐ 3 Moderate in amplitude with posture holding as well as action ☐ 4 Marked in amplitude; interferes with feeding ☐ 8 Untestable (SPECIFY):		
Check the most appropriate box: 0 Absent 1 Slight; present with action 2 Moderate in amplitude, present with action	☐ 3 Moderate in amplitude with posture holding as well as action ☐ 4 Marked in amplitude; interferes with feeding ☐ 8 Untestable (SPECIFY):		
5. Rigidity (judged on passive movement of major joints with patient relaxed in sitting position; cogwheeling to be ignored)			
Check the most appropriate box: 0 Absent 1 Slight or detectable only when activated by mirror or other movements 2 Mild to moderate	☐ 3 Marked, but full range of motion easily achieved ☐ 4 Severe; range of motion achieved with difficulty ☐ 8 Untestable (SPECIFY):		
Check the most appropriate box: 0 Absent 1 Slight or detectable only when activated by mirror or other movements 2 Mild to moderate	☐ 3 Marked, but full range of motion easily achieved ☐ 4 Severe; range of motion achieved with difficulty ☐ 8 Untestable (SPECIFY):		
	□ 0 Absent □ 1 Slight and infrequently present □ 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present Check the most appropriate box: □ 0 Absent □ 1 Slight and infrequently present □ 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present Check the most appropriate box: □ 0 Absent □ 1 Slight and infrequently present □ 2 Mild in amplitude and persistent; or moderate in amplitude, but only intermittently present I tremor of hands Check the most appropriate box: □ 0 Absent □ 1 Slight; present with action □ 2 Moderate in amplitude, present with action Check the most appropriate box: □ 0 Absent □ 1 Slight; present with action □ 2 Moderate in amplitude, present with action □ 2 Moderate in amplitude, present with action □ 1 Slight; present with patient relaxed in sitting position of the most appropriate box: □ 0 Absent □ 1 Slight or detectable only when activated by mirror or other movements □ 2 Mild to moderate Check the most appropriate box: □ 0 Absent □ 1 Slight or detectable only when activated by mirror or other movements □ 2 Mild to moderate Check the most appropriate box: □ 0 Absent □ 1 Slight or detectable only when activated by mirror or other movements □ 2 Mild to moderate		

Form date: ____ / ___ / ___ __ Visit #: _

Participant ID: ___

5. Rigidity		continued		
5c. Left upper extremity	Check the most appropriate box: ☐ 0 Absent ☐ 1 Slight or detectable only when activated by mirror or other movements ☐ 2 Mild to moderate	☐ 3 Marked, but full range of motion easily achieved ☐ 4 Severe; range of motion achieved with difficulty ☐ 8 Untestable (SPECIFY):		
5d. Right lower extremity	Check the most appropriate box: □ 0 Absent □ 1 Slight or detectable only when activated by mirror or other movements □ 2 Mild to moderate	☐ 3 Marked, but full range of motion easily achieved ☐ 4 Severe; range of motion achieved with difficulty ☐ 8 Untestable (SPECIFY):		
5e. Left lower extremity	Check the most appropriate box: ☐ 0 Absent ☐ 1 Slight or detectable only when activated by mirror or other movements ☐ 2 Mild to moderate	☐ 3 Marked, but full range of motion easily achieved ☐ 4 Severe; range of motion achieved with difficulty ☐ 8 Untestable (SPECIFY):		
6. Finger taps (patient taps thumb wi	ith index finger in rapid succession)			
6a. Right hand	Check the most appropriate box: O Normal Mild slowing and/or reduction in amplitude Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement 4 Can barely perform the task. 8 Untestable (SPECIFY):		
6b. Left hand	Check the most appropriate box: O Normal 1 Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement 4 Can barely perform the task. 8 Untestable (SPECIFY):		
7. Hand movements				
(patient opens and clos	ses hands in rapid succession)			
7a. Right hand	Check the most appropriate box: O Normal I Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement 4 Can barely perform the task. 8 Untestable (SPECIFY):		
7b. Left hand	Check the most appropriate box: 0 Normal 1 Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	☐ 3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement ☐ 4 Can barely perform the task. ☐ 8 Untestable (SPECIFY):		

Form date: ____ / ____ / ____ Visit #: _

Participant ID: ____

8. Rapid alternating movements of hands (pronation-supination movements of hands, vertically and horizontally, with as large an amplitude as possible, both hands simultaneously)				
8a. Right hand	Check the most appropriate box: 0 Normal 1 Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	☐ 3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement ☐ 4 Can barely perform the task. ☐ 8 Untestable (SPECIFY):		
8b. Left hand	Check the most appropriate box: 0 Normal 1 Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	☐ 3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement ☐ 4 Can barely perform the task. ☐ 8 Untestable (SPECIFY):		
9. Leg agility (patient taps heel on the ground in rapid succession, picking up entire leg; amplitude should be at least 3 inches)				
9a. Right leg	Check the most appropriate box: 0 Normal 1 Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	☐ 3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement ☐ 4 Can barely perform the task. ☐ 8 Untestable (SPECIFY):		
9b. Left leg	Check the most appropriate box: 0 Normal 1 Mild slowing and/or reduction in amplitude 2 Moderately impaired; definite and early fatiguing; may have occasional arrests in movement	☐ 3 Severely impaired; frequent hesitation in initiating movements or arrests in ongoing movement ☐ 4 Can barely perform the task. ☐ 8 Untestable (SPECIFY):		
10. Arising from chair (patient attempts to rise from a straightbacked chair, with arms folded across chest)	Check the most appropriate box: ☐ 0 Normal ☐ 1 Slow; or may need more than one attempt. ☐ 2 Pushes self up from arms of seat.	☐ 3 Tends to fall back and may have to try more than one time, but can get up without help. ☐ 4 Unable to arise without help ☐ 8 Untestable (SPECIFY):		
11. Posture	Check the most appropriate box: ☐ 0 Normal ☐ 1 Not quite erect, slightly stooped posture; could be normal for older person. ☐ 2 Moderately stooped posture, definitely abnormal; can be slightly leaning to one side.	☐ 3 Severely stooped posture with kyphosis; can be moderately leaning to one side. ☐ 4 Marked flexion with extreme abnormality of posture. ☐ 8 Untestable (SPECIFY):		

____ Form date: ____ / ____ / ____ / ____ Visit #: __

Participant ID: ____

12. Gait	Check the most appropriate box: 0 Normal 1 Walks slowly; may shuffle with short steps, but no festination (hastening steps) or propulsion. 2 Walks with difficulty, but requires little or no assistance; may have some festination, short steps, or propulsion.	☐ 3 Severe disturbance of gait requiring assistance. ☐ 4 Cannot walk at all, even with assistance. ☐ 8 Untestable (SPECIFY):
13. Posture stability (response to sudden, strong posterior displacement produced by pull on shoulders while patient erect with eyes open and feet slightly apart; patient is prepared)	Check the most appropriate box: 0 Normal erect 1 Retropulsion, but recovers unaided. 2 Absence of postural response; would fall if not caught by examiner.	☐ 3 Very unstable, tends to lose balance spontaneously. ☐ 4 Unable to stand without assistance. ☐ 8 Untestable (SPECIFY):
14. Body bradykinesia and hypokinesia (combining slowness, hesitancy, decreased arm swing, small amplitude, and poverty of movement in general)	Check the most appropriate box: O None I Minimal slowness, giving movement a deliberate character; could be normal for some persons; possibly reduced amplitude. D Mild degree of slowness and poverty of movement which is definitely abnormal. Alternatively, some reduced amplitude.	☐ 3 Moderate slowness, poverty or small amplitude of movement. ☐ 4 Marked slowness, poverty or small amplitude of movement. ☐ 8 Untestable (SPECIFY):

Participant ID: _____ Form date: ___ / ___ / ___ Visit #: ____ Visit #: