



Boxes and Locks






:

\square *Bool*



$\vdash t : A$




$$\frac{\vdash A}{\Gamma \vdash \Box A} \text{ NEC.}$$




$$K: \Gamma \vdash \Box(A \rightarrow B) \rightarrow \Box A \rightarrow \Box B$$




$$\mathsf{T}: \Gamma \vdash \Box A \rightarrow A$$



$$4: \Gamma \vdash \Box A \rightarrow \Box \Box A$$





Fitch-style IK

$\Gamma ::= \cdot \mid \Gamma, x : A \mid \Gamma, \blacksquare$

$$\frac{}{\Gamma, x : A, \Gamma' \vdash x : A} \blacksquare \notin \Gamma'$$

$$\frac{\Gamma \vdash t : \Box A}{\Gamma, \blacksquare, \Gamma' \vdash \mathbf{unbox} \ t : A} \blacksquare \notin \Gamma'$$

$$\frac{\Gamma, \blacksquare \vdash t : A}{\Gamma \vdash \mathbf{box} \ t : \Box A}$$

Axiom K is derivable

$$\frac{\frac{\dots, \blacksquare \vdash \mathbf{unbox} f : A \rightarrow B \quad \dots, \blacksquare \vdash \mathbf{unbox} x : A}{\Gamma, f : \Box(A \rightarrow B), x : \Box A, \blacksquare \vdash \mathbf{app} (\mathbf{unbox} f) (\mathbf{unbox} x) : B}}{\Gamma, f : \Box(A \rightarrow B), x : \Box A \vdash \mathbf{box} (\mathbf{app} (\mathbf{unbox} f) (\mathbf{unbox} x)) : \Box B}$$

$$\frac{\Gamma \vdash t : \Box A}{\Gamma, \blacksquare, \Gamma' \vdash \mathbf{unbox} t : A} \blacksquare \notin \Gamma' \quad \frac{\Gamma, \blacksquare \vdash t : A}{\Gamma \vdash \mathbf{box} t : \Box A}$$

Necessitation is admissible

$$\vdash A \implies \vdash \Box A$$

$$\frac{\vdash t : A}{\mathsf{lock} \vdash t : A} \text{WK} \\ \hline \vdash \mathbf{box} \, t : \Box A$$

What about "Denecessitation"?

$$\vdash \Box A \implies \vdash A$$

How would you prove this?

$$\frac{\Gamma \vdash t : \Box A}{\Gamma, \blacksquare, \Gamma' \vdash \mathbf{unbox} \ t : A} \blacksquare \notin \Gamma' \qquad \frac{\Gamma, \blacksquare \vdash t : A}{\Gamma \vdash \mathbf{box} \ t : \Box A}$$

Normalization Rules

$$(\beta) \quad \mathbf{unbox} (\mathbf{box} \, t) \longrightarrow t$$

$$(\eta) \quad \Gamma \vdash t \longrightarrow \mathbf{box} (\mathbf{unbox} \, t) : \Box A$$

Normal Forms

$$\frac{\Gamma, \mathbf{\boxdot} \vdash_{\text{nf}} t : A}{\Gamma \vdash_{\text{nf}} \mathbf{box} \ t : \Box A}$$

$$\frac{\Gamma \vdash_{\text{ne}} t : \Box A}{\Gamma, \mathbf{\boxdot}, \Gamma' \vdash_{\text{ne}} \mathbf{unbox} \ t : A} \quad \mathbf{\boxdot} \notin \Gamma'$$

Normalization by Evaluation

$$(\llbracket _ \rrbracket) : (\Gamma \vdash A) \rightarrow \llbracket \Gamma \rrbracket_{\Delta} \rightarrow \llbracket A \rrbracket_{\Delta}$$

$$\downarrow_A : \llbracket A \rrbracket_{\Gamma} \rightarrow (\Gamma \vdash_{\text{nf}} A)$$

$$\text{norm } t = \downarrow (\llbracket t \rrbracket \gamma_{id})$$

Interpretation

$$\llbracket \tau \rrbracket_{\Gamma} = \Gamma \vdash_{\text{nf}} \tau$$

$$\llbracket A \rightarrow B \rrbracket_{\Gamma} = \Gamma' \leq \Gamma \rightarrow \llbracket A \rrbracket_{\Gamma'} \rightarrow \llbracket B \rrbracket_{\Gamma'}$$

$$\llbracket \Box A \rrbracket_{\Gamma} = \text{Box}_{\Gamma} \llbracket A \rrbracket$$

$$\llbracket \cdot \rrbracket_{\Gamma} = \top$$

$$\llbracket \Delta, A \rrbracket_{\Gamma} = \llbracket \Delta \rrbracket_{\Gamma} \times \llbracket A \rrbracket_{\Gamma}$$

$$\llbracket \Delta, \mathbf{\Delta} \rrbracket_{\Gamma} = \text{Lock}_{\Gamma} \llbracket \Delta \rrbracket$$

Boxes and Locks

$\mathcal{A} : \text{Ctx} \rightarrow \text{Set}$

$$\frac{x : \mathcal{A}_{\Gamma, \blacksquare}}{\text{box } x : \text{Box}_{\Gamma} \mathcal{A}}$$

$$\frac{x : \mathcal{A}_{\Gamma}}{\text{lock } x : \text{Lock}_{\Gamma, \blacksquare, \Gamma'} \mathcal{A}} \blacksquare \notin \Gamma'$$

$$\text{unbox} : \text{Box}_{\Gamma} \llbracket A \rrbracket \rightarrow \llbracket A \rrbracket_{\Gamma, \blacksquare, \Gamma'} \quad (\blacksquare \notin \Gamma')$$

$$\text{unbox} (\text{box } x) = wk \ x$$

Evaluation

$$(\llbracket _ \rrbracket) : (\Gamma \vdash A) \rightarrow \llbracket \Gamma \rrbracket_{\Delta} \rightarrow \llbracket A \rrbracket_{\Delta}$$

$$(\llbracket \mathbf{box} \ t \rrbracket) \ \gamma = \mathbf{box} \ ((\llbracket t \rrbracket) \ \gamma)$$

$$(\llbracket \mathbf{unbox} \ t \rrbracket) \ \langle \gamma, _ \rangle = (\llbracket \mathbf{unbox} \ t \rrbracket) \ \gamma$$

$$(\llbracket \mathbf{unbox} \ t \rrbracket) \ (\mathbf{lock} \ \gamma) = \mathbf{unbox} \ ((\llbracket t \rrbracket) \ \gamma)$$

$$\frac{\Gamma \vdash t : \Box A}{\Gamma, \blacksquare, \Gamma' \vdash \mathbf{unbox} \ t : A} \blacksquare \notin \Gamma' \qquad \frac{\Gamma, \blacksquare \vdash t : A}{\Gamma \vdash \mathbf{box} \ t : \Box A}$$

Reification and Reflection

$$\downarrow_A : \llbracket A \rrbracket_\Gamma \rightarrow (\Gamma \vdash_{\text{nf}} A)$$

$$\downarrow_{\Box A} (\mathbf{box} \ x) = \mathbf{box} \ (\downarrow_A \ x)$$

$$\uparrow_A : (\Gamma \vdash_{\text{ne}} A) \rightarrow \llbracket A \rrbracket_\Gamma$$

$$\uparrow_{\Box A} m = \mathbf{box} \ (\uparrow_A \ (\mathbf{unbox} \ m))$$

Denecessitation can be *proved using normal forms*

$$\vdash \Box A \implies \vdash A$$

$$\frac{\frac{\times}{\vdash_{\text{ne}} t : \Box B}}{\text{🔒} \vdash_{\text{ne}} \mathbf{unbox} \ t : B} \cdot \cdot \frac{\text{🔒} \vdash_{\text{nf}} t : A}{\vdash_{\text{nf}} \mathbf{box} \ t : \Box A}$$

No neutrals in empty context, dismissed!

Only culprit that introduces 🔒

Remove 🔒 to get $\vdash_{\text{nf}} t : \Box A$



Why?

1. Useful

- Towards confluence, decidability, consistency, etc.
- Simplify / prove new application-specific theorems?
(e.g., partial evaluation/staging theorems, noninterference, etc.)
- Type-directed partial evaluation for modal type systems

2. Extensible: Boxes, Locks and (fake) Diamonds!


IK♦

$$\frac{\Gamma \vdash t : A}{\Gamma, \blacksquare, \Gamma' \vdash \mathbf{dia} \ t : \blacklozenge A} \blacksquare \notin \Gamma' \quad \frac{\Gamma \vdash t : \blacklozenge A \quad A, \blacksquare \vdash u : B}{\Gamma \vdash \mathbf{bind} \ t \ u : B}$$

Interpreting ♦

$$\frac{x : \text{Lock}_\Gamma \mathcal{A}}{\text{val } x : \text{Dia}_\Gamma \mathcal{A}} \quad \frac{m : \Gamma \vdash_{\text{ne}} \blacklozenge A \quad y : \text{Dia}_{A, \blacksquare} \mathcal{A}}{\text{bindst } m \ y : \text{Dia}_\Gamma \mathcal{A}} \blacksquare \notin \Gamma'$$

2. Extensible: IS₄, IK₄, IR



3. Comprehensible: Reduction traces

DEMO


4. Elegant: Boxes and Locks, in an Adjoint relationship

$$unit : \mathcal{A} \multimap \text{Box} (\text{Lock } \mathcal{A})$$

$$unit\ x = \text{box} (\text{lock } x)$$

$$counit : \text{Lock} (\text{Box } \mathcal{A}) \multimap \mathcal{A}$$

$$counit (\text{lock} (\text{box } x)) = wk\ x$$



What would you like to see more of?

Modalities (S_4 , K_4 , R, etc.)

Semantics

Applications

Extensions (0, +, Nat, etc.)

EOM

