







Rank



Try the Next Challenge | Try a Random Challenge







Data Structures Challenges

Your Minimum Absolute Difference in an Array submission got 15.00 points.

Comparti

Tweet

×

# Minimum Absolute Difference in an





Problem

Submissions

Leaderboard

Discussions

Editorial

Consider an array of integers,  $A = a_0, a_1, \dots, a_{n-1}$ . We define the absolute difference between two elements,  $a_i$  and  $a_j$  (where  $i \neq j$ ), to be the absolute value of  $a_i - a_j$ .

Given an array of  $m{n}$  integers, find and print the minimum absolute difference between any two elements in the array.

### **Input Format**

The first line contains a single integer denoting n (the number of integers).

The second line contains n space-separated integers describing the respective values of  $a_0, a_1, \ldots, a_{n-1}$ .

# Constraints

- $2 \le n \le 10^5$
- $-10^9 \le a_i \le 10^9$

#### **Output Format**

Print the minimum absolute difference between any two elements in the array.

## Sample Input 0

3 3 -7 0

#### **Sample Output 0**


3

# **Explanation 0**

With n = 3 integers in our array, we have three possible pairs: (3, -7), (3, 0), and (-7, 0). The absolute values of the differences between these pairs are as follows:


- $|3--7| \Rightarrow 10$
- $|3-0| \Rightarrow 3$
- $|-7-0| \Rightarrow 7$

Notice that if we were to switch the order of the numbers in these pairs, the resulting absolute values would still be the same. The smallest of these possible absolute differences is **3**, so we print **3** as our answer.



Run Code

Submit Code





Copyright © 2017 HackerRank. All Rights Reserved

Join us on IRC at #hackerrank on freenode for hugs or bugs.

Contest Calendar | Interview Prep | Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Terms Of Service | Privacy Policy | Request a Feature

Test against custom input

**1** Upload Code as File