

Building Single Molecules from Single Atoms

Yichao Yu

Lee Liu, Kenneth Wang, Lewis Picard, Jonathan Hood
Jessie T. Zhang, Eliot Fenton, Yen-Wei Lin

Ni Group/Harvard

Sep 5, 2019


Atom

- Laser cooling/trapping
 - Internal state control
 - High fidelity imagining
- ⋮

From Atom to Molecule

Atom

- Laser cooling/trapping
- Internal state control
- High fidelity imagining
- ⋮



Nature 545, 462-466 (2017)


From Atom to Molecule

Atom

- Laser cooling/trapping
- Internal state control
- High fidelity imagining
- ⋮

Molecule

- Strong interaction
- Rich internal structure




Nature 545, 462-466 (2017)

From Atom to Molecule

Atom


- Laser cooling/trapping
- Internal state control
- High fidelity imaging

Quantum Chemistry



Nature 464, 1324 (2010)

Quantum Simulation




Nat. Phys. 2, 341 (2006)

Molecule


- Strong interaction
- Rich internal structure

Quantum Computing



Phys. Rev. Lett. 97, 33003 (2006)

Precision Measurement




Science 343, p. 269-272 (2014)

Path to Ultracold Molecules

Direct molecule cooling

“Diagonal molecules”:
CaF, SrF, YO, . . .




Phys. Rev. Lett. 119, 103201 (2017)

Path to Ultracold Molecules

Direct molecule cooling

“Diagonal molecules”:
CaF, SrF, YO, . . .



Phys. Rev. Lett. 119, 103201 (2017)

Making molecule from atoms


Bi-Alkali/Alkaline Earth:
KRb, RbCs, NaK, LiNa, . . .

- 1 Cold gas
- 2 Feshbach association
- 3 STIRAP

Path to Ultracold Molecules

Assemble molecule in tweezers

- Single site detection
- Single site addressing
- Control loading and interaction by rearranging.



Making molecule from atoms

Bi-Alkali/Alkaline Earth:
KRb, RbCs, NaK, LiNa, ...

- ① Cold gas
- ② Feshbach association
- ③ STIRAP

Outline

1 System Overview

2 Trapping and Cooling of Atoms

3 Atom-Atom Interaction and Molecule Formation

Steps



Loading




Steps




Loading

Cooling


Steps




Steps




Steps



Science 360 6391, 2018






Single Atom in Tweezer

- Previously done with Rb
- Works for Cs
- Doesn't work for Na

Single Atom in Tweezer

- Previously done with Rb
- Works for Cs
- Doesn't work for Na



Single Atom in Tweezer

- Previously done with Rb
- Works for Cs
- Doesn't work for Na




Single Atom in Tweezer

- Previously done with Rb
- Works for Cs
- Doesn't work for Na


Issues with Na

- Low vapor pressure
- Broad linewidth
- Low mass
- Small hyperfine structure


Real Issue with Na: Light Shift



Real Issue with Na: Light Shift



Cs single atom imaging




Real Issue with Na: Light Shift

Trap modulation


Alternate between trap and resonant (cooling and imaging) light at 2.5 MHz

$$f_{trap} = 100 \sim 500 \text{ kHz}$$

$$\Gamma = 2\pi \times 10 \text{ MHz}$$



Cs single atom imaging




Real Issue with Na: Light Shift

Trap modulation


Alternate between trap and resonant (cooling and imaging) light at 2.5 MHz

$$f_{trap} = 100 \sim 500 \text{ kHz}$$

$$\Gamma = 2\pi \times 10 \text{ MHz}$$



Cs single atom imaging




Real Issue with Na: Light Shift

Trap modulation



Alternate between trap and resonant (cooling and imaging) light at 2.5 MHz

$$f_{trap} = 100 \sim 500 \text{ kHz}$$


$$\Gamma = 2\pi \times 10 \text{ MHz}$$




Cs single atom imaging




Raman Sideband Cooling



Raman Sideband Cooling



Axial sideband spectrum



Outline


1 System Overview

2 Trapping and Cooling of Atoms


3 Atom-Atom Interaction and Molecule Formation

Optical Transfer to Molecular State

Binding energy
 $\approx 150\text{THz}$




Molecule




Atoms

Optical Transfer to Molecular State

Binding energy
 $\approx 150\text{THz}$




Binding energy
 $\approx 300\text{MHz}$




Molecule

Weakly-Bound
Molecule




Atoms


Optical Transfer to Molecular State




Photoassociation (PA) Spectroscopy




Photoassociation (PA) Spectroscopy



Probability of empty trap




Photoassociation (PA) Spectroscopy



Optical Transfer to Weakly-Bound Molecular


Scattering length a

- Binding energy
- Molecular potential
- Feshbach resonance
- Molecule formation
- ⋮




Scattering length a

- Binding energy
- Molecular potential
- Feshbach resonance
- Molecule formation
- ⋮




Scattering length a

- Binding energy
- Molecular potential
- Feshbach resonance
- Molecule formation
- ⋮




Scattering length a

- Binding energy
- Molecular potential
- Feshbach resonance
- Molecule formation
- ⋮




Scattering length a


- Binding energy
- Molecular potential
- Feshbach resonance
- Molecule formation
- ⋮





Interaction shift



Interaction shift




Interaction shift



Interaction shift

$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{m_1 \omega_{1,i}^2 x_{1,i}^2}{2} + \frac{p_{1,i}^2}{2m_1} \right)}_{\text{Na}} + \underbrace{\sum_{i=x,y,z} \left(\frac{m_2 \omega_{2,i}^2 x_{2,i}^2}{2} + \frac{p_{2,i}^2}{2m_2} \right)}_{\text{Cs}} + V_{int}(\vec{r}_1 - \vec{r}_2)$$

Interaction



Interaction shift

$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{m_1 \omega_{1,i}^2 X_{1,i}^2}{2} + \frac{p_{1,i}^2}{2m_1} \right)}_{\text{Na}} + \underbrace{\sum_{i=x,y,z} \left(\frac{m_2 \omega_{2,i}^2 X_{2,i}^2}{2} + \frac{p_{2,i}^2}{2m_2} \right)}_{\text{Cs}} + V_{int}(\vec{r}_1 - \vec{r}_2) \underbrace{\quad}_{\text{Interaction}}$$

To center of mass
and relative coordinates

$$M = m_1 + m_2$$

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$

$$\Omega_i^2 = \frac{m_1 \omega_{1,i}^2 + m_2 \omega_{2,i}^2}{m_1 + m_2}$$


$$\omega_{R,i}^2 = \frac{m_2 \omega_{1,i}^2 + m_1 \omega_{2,i}^2}{m_1 + m_2}$$

$$X_i = \frac{m_1 x_{1,i} + m_2 x_{2,i}}{m_1 + m_2}$$

$$x_{R,i} = x_{1,i} - x_{2,i}$$

$$P_i = p_{1,i} + p_{2,i}$$



$$p_{R,i} = \frac{m_2 p_{1,i} - m_1 p_{2,i}}{m_1 + m_2}$$



Center of mass



$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{M \Omega_i^2 X_i^2}{2} + \frac{P_i^2}{2M} \right)}_{\text{Center of mass}} + \underbrace{\sum_{i=x,y,z} \left(\frac{\mu \omega_{R,i}^2 X_{R,i}^2}{2} + \frac{p_{R,i}^2}{2\mu} \right) + V_{int}(\vec{r}_R)}_{\text{Relative}} + \underbrace{\sum_{i=x,y,z} \mu (\omega_{1,i}^2 - \omega_{2,i}^2) X_i X_{R,i}}_{\text{Mixing}}$$

Interaction shift





$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{M\Omega_i^2 X_i^2}{2} + \frac{P_i^2}{2M} \right)}_{\text{Center of mass}} + \underbrace{\sum_{i=x,y,z} \left(\frac{\mu\omega_{R,i}^2 X_{R,i}^2}{2} + \frac{p_{R,i}^2}{2\mu} \right) + V_{int}(\vec{r}_R)}_{\text{Relative}} + \underbrace{\sum_{i=x,y,z} \mu(\omega_{1,i}^2 - \omega_{2,i}^2) X_i X_{R,i}}_{\text{Mixing}}$$

Interaction shift




$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{M\Omega_i^2 X_i^2}{2} + \frac{P_i^2}{2M} \right)}_{\text{Center of mass}} + \underbrace{\sum_{i=x,y,z} \left(\frac{\mu\omega_{R,i}^2 X_{R,i}^2}{2} + \frac{p_{R,i}^2}{2\mu} \right) + V_{int}(\vec{r}_R)}_{\text{Relative}} + \underbrace{\sum_{i=x,y,z} \mu(\omega_{1,i}^2 - \omega_{2,i}^2) X_i X_{R,i}}_{\text{Mixing}}$$

Interaction shift




$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{M\Omega_i^2 X_i^2}{2} + \frac{P_i^2}{2M} \right)}_{\text{Center of mass}} + \underbrace{\sum_{i=x,y,z} \left(\frac{\mu\omega_{R,i}^2 X_{R,i}^2}{2} + \frac{p_{R,i}^2}{2\mu} \right) + V_{int}(\vec{r}_R)}_{\text{Relative}} + \underbrace{\sum_{i=x,y,z} \mu(\omega_{1,i}^2 - \omega_{2,i}^2) X_i X_{R,i}}_{\text{Mixing}}$$

Interaction shift



$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{M\Omega_i^2 X_i^2}{2} + \frac{P_i^2}{2M} \right)}_{\text{Center of mass}} + \underbrace{\sum_{i=x,y,z} \left(\frac{\mu\omega_{R,i}^2 X_{R,i}^2}{2} + \frac{P_{R,i}^2}{2\mu} \right) + V_{int}(\vec{r}_R)}_{\text{Relative}} + \underbrace{\sum_{i=x,y,z} \mu(\omega_{1,i}^2 - \omega_{2,i}^2) X_i X_{R,i}}_{\text{Mixing}}$$

Interaction shift




$$H = \underbrace{\sum_{i=x,y,z} \left(\frac{M\Omega_i^2 X_i^2}{2} + \frac{P_i^2}{2M} \right)}_{\text{Center of mass}} + \underbrace{\sum_{i=x,y,z} \left(\frac{\mu\omega_{R,i}^2 X_{R,i}^2}{2} + \frac{P_{R,i}^2}{2\mu} \right) + V_{int}(\vec{r}_R)}_{\text{Relative}} + \underbrace{\sum_{i=x,y,z} \mu(\omega_{1,i}^2 - \omega_{2,i}^2) X_i X_{R,i}}_{\text{Mixing}}$$


Na (1, -1) Cs (3, -3) Feshbach resonance



Na (1, -1) Cs (3, -3) Feshbach resonance



Na (1, -1) Cs (3, -3) Feshbach resonance



	<i>s</i> -wave	<i>p</i> -wave
Predicted (based on interaction shift) ¹	663 G	799 G
Measured	652(3) G	791.2(2) G

¹In collaboration with Bo Gao

