

# Game theory and adaptive dynamics

LSM 4255 Methods in Mathematical Biology

Week 12

# Outline

- Brief history of game theory
- 2-strategy games, evolutionary analysis
- Adaptive dynamics
- Tragedy of the Commons

# Natural selection

- In previous lectures, you looked at population genetic models like this

$$p_{n+1} = u (p_n + q_n)^2 \frac{1}{d_n} \text{ freq. AA} \quad (1)$$

$$q_{n+1} = v (p_n + q_n) (q_n + r_n) \frac{1}{d_n} \text{ half freq. Aa} \quad (2)$$

$$r_{n+1} = w (q_n + r_n)^2 \frac{1}{d_n} \text{ freq aa} \quad (3)$$

$$d_n = u(p_n + q_n)^2 2v(p_n + q_n)(q_n + r_n) w(q_n + r_n)^2 \quad (4)$$

where  $u, v, w$  are fitness.

- Where does fitness come from?

- Previously assumed a constant, reflecting something about a static or averaged environment,
- But might be more complicated than that

## Fitness depends on genotypes and frequencies of others

Fitness is determined not just by an individual's own genotype but by the types and frequencies of other genes in the population

# Fitness depends on genotypes and frequencies of others

Fitness is determined not just by an individual's own genotype but by the types and frequencies of other genes in the population



# Fitness depends on genotypes and frequencies of others

Fitness is determined not just by an individual's own genotype but by the types and frequencies of other genes in the population



photo BBC

# Fitness depends on genotypes and frequencies of others

Fitness is determined not just by an individual's own genotype but by the types and frequencies of other genes in the population



photo BBC

Need some mathematical framework to deal with this

- Game theory
- Came to biology from economics / military

# A brief history of game theory - 1. cooperative game theory

- John von Neumann and Oskar Morgenstern in 1944
  - Book – *Theory of Games and Economic Behavior*
    - "we wish to find the mathematically complete principles which define 'rational behaviour' for the participants in a social economy, and to derive from them the general characteristics of that behaviour"
  - cooperative game theory
    - About humans forming coalitions, making agreements, splitting costs/profits
    - e.g. Several nearby towns want a water supply
  - Not yet so interesting to biologists ...



Morgenstern (left) and von Neumann (right)

# A brief history of game theory - 2. non-cooperative games

- John Nash - non-cooperative games
  - A more general theory
  - No enforcement mechanisms (e.g. contracts to split costs) outside the game itself
  - Not about coalitions, agreements and side-payments possible between players, but rather individual strategies and payoffs
- Individualistic
  - ‘Darwinian’ view of the world: each works for themselves and maximises own payoff



# A brief history of game theory - 2. non-cooperative games

- John Nash - non-cooperative games
  - A more general theory
  - No enforcement mechanisms (e.g. contracts to split costs) outside the game itself
  - Not about coalitions, agreements and side-payments possible between players, but rather individual strategies and payoffs
- Individualistic
  - 'Darwinian' view of the world: each works for themselves and maximises own payoff
- 'Games'
  - All players are rational and have full information about game
  - Players cannot coordinate with each other
  - Single-shot
  - Only objective is to maximise own payoff



# Example: The stag hunt

Two people go on a hunt. Each can either hunt stag or hunt hare. A stag is worth more but takes two people. A hare is easily caught alone.

- What should an individualistic rational player do?

<b>S</b> <b>h</b>	COOPERATE	DEFECT
COOPERATE		
DEFECT		

## Example: The stag hunt

Two people go on a hunt. Each can either hunt stag or hunt hare. A stag is worth more but takes two people. A hare is easily caught alone.

- What should an individualistic rational player do?
- Payoff matrix

	<b>s</b>	<b>h</b>
	 COOPERATE	 DEFECT
 COOPERATE	3 3	0 2
 DEFECT	2 0	1 1

## Example: The stag hunt

Two people go on a hunt. Each can either hunt stag or hunt hare. A stag is worth more but takes two people. A hare is easily caught alone.

- What should an individualistic rational player do?
- Payoff matrix
  - If other player hunting stag?

	<b>s</b>	<b>h</b>
	 COOPERATE	 DEFECT
 COOPERATE	3 3	0 2
 DEFECT	2 0	1 1

# Example: The stag hunt

Two people go on a hunt. Each can either hunt stag or hunt hare. A stag is worth more but takes two people. A hare is easily caught alone.

- What should an individualistic rational player do?
- Payoff matrix
  - If other player hunting stag?
  - If other player hunting hare?

	<b>s</b>	<b>h</b>	
			<b>2</b>
	<b>COOPERATE</b>	<b>DEFECT</b>	
<b>1</b>		<b>3</b>	<b>2</b>
<b>DEFECT</b>		<b>3</b>	<b>0</b>
<b>2</b>		<b>0</b>	<b>1</b>
<b>1</b>			<b>1</b>

## Example: The stag hunt

Two people go on a hunt. Each can either hunt stag or hunt hare. A stag is worth more but takes two people. A hare is easily caught alone.

- What should an individualistic rational player do?
- Payoff matrix
  - If other player hunting stag?
  - If other player hunting hare?
- My best strategy depends on others' strategy

	<b>s</b>	<b>h</b>	
			<b>COOPERATE</b>
<b>COOPERATE</b>		<b>3</b>	<b>2</b>
<b>DEFECT</b>		<b>3</b>	<b>0</b>

	<b>s</b>	<b>h</b>	
			<b>COOPERATE</b>
<b>COOPERATE</b>		<b>3</b>	<b>2</b>
<b>DEFECT</b>		<b>3</b>	<b>0</b>

	<b>s</b>	<b>h</b>	
			<b>COOPERATE</b>
<b>COOPERATE</b>		<b>3</b>	<b>2</b>
<b>DEFECT</b>		<b>2</b>	<b>1</b>

# Example: The stag hunt

- Nash equilibrium:

- A strategy is a Nash equilibrium ( $x^*$ ) if no player can do better by unilaterally changing their strategy.

$$f(x^*, x^*) \geq f(x, x^*) \quad (5)$$

for all other strategies  $x$

- The stag hunt has two Nash equilibria:


$$f(\text{stag, stag}) > f(\text{hare, stag}) \quad (6)$$

$$f(\text{hare, hare}) > f(\text{stag, hare}) \quad (7)$$

Notation: payoff to a focal individual is  $f$ (focal individ's strategy, others' strategy)

		S <sub>h</sub>	
		COOPERATE	DEFECT
COOPERATE	COOPERATE	2	1
	DEFECT	1	0
DEFECT	COOPERATE	3	2
	DEFECT	0	1

# Example of stag hunt in nature - carousel hunting



# Evolutionary game theory

Maynard Smith & Price (1973):

- Males often compete for territory, etc. → transmission of genes
- Might expect natural selection to favour maximally effective weapons and fighting styles for a “total war” strategy, battles to the death
- Instead, a “limited war” type is common
- ‘Group selection’ type explanation was accepted explanation



© Arkaprava Ghosh / Barcroft India

# Evolutionary game theory

- Maynard Smith & Price recast the game theory into biological context:
  - 'Game' → interaction that determines fitness (e.g. snakes fighting for territory)
  - 'Strategy' → genetically encoded behaviour or trait
  - 'Player' → individual animal, though better to think of as gene
  - 'Payoff' → fitness

# Evolutionary game theory

- Maynard Smith & Price recast the game theory into biological context:
  - 'Game' → interaction that determines fitness (e.g. snakes fighting for territory)
  - 'Strategy' → genetically encoded behaviour or trait
  - 'Player' → individual animal, though better to think of as gene
  - 'Payoff' → fitness
- Recall the four processes of population genetics:
  - ① Selection
  - ② Mutation
  - ③ Genetic drift
  - ④ Gene flow

Basic evolutionary game theory only includes the first (replicator dynamics), and second (ESS, adaptive dynamics), though can be extended to include others

# Evolutionary game theory - Hawk-Dove

- 'Hawk-Dove' game (see Hastings text)
  - two animals contesting favoured territory with value  $V$
  - fighting over it has injury cost  $C > V$
- So ... total war?

		2	
	COOPERATE	 2	 DEFECT
1	COOPERATE	$V/2$	$V$
1	DEFECT	$V/2$	0
		0	$(V-C)/2$
		$V$	$(V-C)/2$

# Evolutionary game theory - Hawk-Dove

- 'Hawk-Dove' game (see Hastings text)
  - two animals contesting favoured territory with value  $V$
  - fighting over it has injury cost  $C > V$
- So ... total war?

		2	
	COOPERATE	 2	
 1	COOPERATE	$V/2$	$V$
 1	DEFECT	$V/2$	0
		0	$(V-C)/2$
		$V$	$(V-C)/2$

$$f(\text{hawk, hawk}) < f(\text{dove, hawk})$$

# Evolutionary game theory - Hawk-Dove

- 'Hawk-Dove' game (see Hastings text)
  - two animals contesting favoured territory with value  $V$
  - fighting over it has injury cost  $C > V$
- So ... total war? Peace?

	!d	2	2
	COOPERATE	2	DEFECT
!d	COOPERATE	$V/2$	$V$
1	DEFECT	$V/2$	0

Diagram of the Hawk-Dove game payoff matrix:

- Player 1 (row) has strategies: !d (Cooperate) and 1 (Defect).
- Player 2 (column) has strategies: COOPERATE and DEFECT.
- Payoffs:
  - (!d, COOPERATE): Player 1 gets  $V/2$ , Player 2 gets  $V$ .
  - (!d, DEFECT): Player 1 gets  $V$ , Player 2 gets 0.
  - (1, COOPERATE): Player 1 gets  $V/2$ , Player 2 gets  $V/2$ .
  - (1, DEFECT): Player 1 gets 0, Player 2 gets  $(V-C)/2$ .

$$f(\text{hawk}, \text{hawk}) < f(\text{dove}, \text{hawk})$$

$$f(\text{dove}, \text{dove}) < f(\text{hawk}, \text{dove})$$

# Evolutionary game theory - Hawk-Dove

- 'Hawk-Dove' game (see Hastings text)
  - two animals contesting favoured territory with value  $V$
  - fighting over it has injury cost  $C > V$
- So ... total war? Peace?
- Recall Nash equilibrium

$$f(x^*, x^*) \geq f(x, x^*) \forall x$$

		2	2
	COOPERATE	COOPERATE	DEFECT
!d	COOPERATE	$V/2$	$V$
1	$V/2$	0	$(V-C)/2$
	DEFECT	0	$(V-C)/2$
		$V$	$(V-C)/2$

$$f(\text{hawk}, \text{hawk}) < f(\text{dove}, \text{hawk})$$

$$f(\text{dove}, \text{dove}) < f(\text{hawk}, \text{dove})$$

# Evolutionary game theory - Hawk-Dove

- 'Hawk-Dove' game (see Hastings text)
  - two animals contesting favoured territory with value  $V$
  - fighting over it has injury cost  $C > V$
- So ... total war? Peace?
- Recall Nash equilibrium

$$f(x^*, x^*) \geq f(x, x^*) \forall x$$

- Neither pure strategy is a Nash equilibrium

		2	2
	COOPERATE	COOPERATE	DEFECT
h	COOPERATE	$V/2$	$V$
d	DEFECT	$V/2$	0
1	COOPERATE	0	$(V-C)/2$
	DEFECT	$V$	$(V-C)/2$

$$f(\text{hawk}, \text{hawk}) < f(\text{dove}, \text{hawk})$$

$$f(\text{dove}, \text{dove}) < f(\text{hawk}, \text{dove})$$

# Evolutionary game theory - A technical point

- Nash equilibrium  $f(x^*, x^*) \geq f(x, x^*) \forall x$

# Evolutionary game theory - A technical point

- Nash equilibrium  $f(x^*, x^*) \geq f(x, x^*) \forall x$
- Maynard Smith concept of an *evolutionarily stable strategy* (ESS), two criteria:
  - ①  $f(x^*, x^*) > f(x, x^*) \forall x$  OR
  - ②
    - ①  $f(x^*, x^*) = f(x, x^*)$  AND
    - ②  $f(x^*, x) > f(x, x) \forall x$

# Evolutionary game theory - A technical point

- Nash equilibrium  $f(x^*, x^*) \geq f(x, x^*) \forall x$
- Maynard Smith concept of an *evolutionarily stable strategy* (ESS), two criteria:
  - ①  $f(x^*, x^*) > f(x, x^*) \forall x$  OR
  - ②
    - ①  $f(x^*, x^*) = f(x, x^*)$  AND
    - ②  $f(x^*, x) > f(x, x) \forall x$
- Meaning of two ESS criteria:
  - ① Strong ESS: no alternative strategy can invade
  - ② Weak ESS: if an alternative strategy is neutral,  $x^*$  cannot be eliminated from population

# Evolutionary game theory - A technical point

- Nash equilibrium  $f(x^*, x^*) \geq f(x, x^*) \forall x$
- Maynard Smith concept of an *evolutionarily stable strategy* (ESS), two criteria:
  - ①  $f(x^*, x^*) > f(x, x^*) \forall x$  OR
  - ②
    - ①  $f(x^*, x^*) = f(x, x^*)$  AND
    - ②  $f(x^*, x) > f(x, x) \forall x$
- Meaning of two ESS criteria:
  - ① Strong ESS: no alternative strategy can invade
  - ② Weak ESS: if an alternative strategy is neutral,  $x^*$  cannot be eliminated from population

ESS → Nash equilibrium

- For these 2-strategy games they are equivalent

# Evolutionary game theory - Hawk-Dove

- Neither hawk nor dove are an ESS, intuition suggests that best strategy is something ‘in between’
- Your text (Hastings) investigates a *mixed strategy*, where players pursue ‘hawk’ or ‘dove’ with some probability
- An alternative is to look at the evolution of the *proportions* of hawk- and dove-strategests in a population


	1	2
1	 COOPERATE	 DEFECT
2	 COOPERATE	 DEFECT
1	$V/2$	$V$
2	$V/2$	$0$
1	$0$	$(V-C)/2$
2	$V$	$(V-C)/2$

# Replicator dynamics

- Agents do not have to be conscious or rational: all they need is a strategy that they pass on
- No change in strategy, no mutation to new strategy
- Interested in change in distribution of strategies in population
- ‘Goal’ of the game is to produce as many replicates of oneself as possible


# Replicator dynamics

- Agents do not have to be conscious or rational: all they need is a strategy that they pass on
- No change in strategy, no mutation to new strategy
- Interested in change in distribution of strategies in population
- ‘Goal’ of the game is to produce as many replicates of oneself as possible




# Replicator dynamics

- Agents do not have to be conscious or rational: all they need is a strategy that they pass on
- No change in strategy, no mutation to new strategy
- Interested in change in distribution of strategies in population
- ‘Goal’ of the game is to produce as many replicates of oneself as possible




# Replicator dynamics

- Agents do not have to be conscious or rational: all they need is a strategy that they pass on
- No change in strategy, no mutation to new strategy
- Interested in change in distribution of strategies in population
- ‘Goal’ of the game is to produce as many replicates of oneself as possible




# Replicator dynamics

- Agents do not have to be conscious or rational: all they need is a strategy that they pass on
- No change in strategy, no mutation to new strategy
- Interested in change in distribution of strategies in population
- ‘Goal’ of the game is to produce as many replicates of oneself as possible



# Replicator dynamics


- Agents do not have to be conscious or rational: all they need is a strategy that they pass on
- No change in strategy, no mutation to new strategy
- Interested in change in distribution of strategies in population
- ‘Goal’ of the game is to produce as many replicates of oneself as possible



# Replicator dynamics equation

Following Chapter 9 of Webb *Game Theory: Decisions, Interaction and Evolution*


- $n_i$ : number of individuals pursuing strategy  $i$
- $N$ : total number of individuals
- $\beta$ : background reproduction rate



# Replicator dynamics equation

Following Chapter 9 of Webb *Game Theory: Decisions, Interaction and Evolution*


- $n_i$ : number of individuals pursuing strategy  $i$
- $N$ : total number of individuals
- $\beta$ : background reproduction rate
- Assume that fitness is a function of *proportions* of strategies in population
  - $f_i(\mathbf{p})$ : fitness effect of strategy  $i$



# Replicator dynamics equation

Following Chapter 9 of Webb *Game Theory: Decisions, Interaction and Evolution*


- $n_i$ : number of individuals pursuing strategy  $i$
- $N$ : total number of individuals
- $\beta$ : background reproduction rate
- Assume that fitness is a function of *proportions* of strategies in population
  - $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $p_i = n_i/N$ : proportion of individuals pursuing strategy  $i$
- Want dynamics of  $p_i$



# Replicator dynamics equation

- Dynamics of  $i$  strategists

$$\frac{dn_i}{dt} = \dot{n}_i = n_i(\beta + f_i) \quad (8)$$



- $n_i$ : no. individts of strategy  $i$
- $N$ : total no. of individts
- $p_i$ : proportions of strategy  $i$
- $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $\beta$ : background reprodn rate


# Replicator dynamics equation

- Dynamics of  $i$  strategists

$$\frac{dn_i}{dt} = \dot{n}_i = n_i(\beta + f_i) \quad (8)$$

- Dynamics of whole population

$$\begin{aligned}\dot{N} &= \sum_i \dot{n}_i \\ &= \beta \sum_i n_i + \sum_i f_i n_i \\ &= \beta N + N \sum_i f_i \frac{n_i}{N} \\ &= N(\beta + \bar{f})\end{aligned} \quad (9)$$



- $n_i$ : no. individvs of strategy  $i$
- $N$ : total no. of individvs
- $p_i$ : proportions of strategy  $i$
- $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $\beta$ : background reprodn rate

where  $\bar{f}$  is mean fitness


# Replicator dynamics equation

- Dynamics of  $i$  strategists

$$\frac{dn_i}{dt} = \dot{n}_i = n_i(\beta + f_i) \quad (8)$$

- Dynamics of whole population

$$\dot{N} = N(\beta + \bar{f}) \quad (9)$$



- $n_i$ : no. individvs of strategy  $i$
- $N$ : total no. of individvs
- $p_i$ : proportions of strategy  $i$
- $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $\beta$ : background reprodn rate

# Replicator dynamics equation


- Dynamics of  $i$  strategists

$$\frac{dn_i}{dt} = \dot{n}_i = n_i(\beta + f_i) \quad (8)$$

- Dynamics of whole population

$$\dot{N} = N(\beta + \bar{f}) \quad (9)$$

- Dynamics of proportions  $p_i = \frac{n_i}{N} \rightarrow n_i = p_i N$



- $n_i$ : no. individvs of strategy  $i$
- $N$ : total no. of individvs
- $p_i$ : proportions of strategy  $i$
- $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $\beta$ : background reprodn rate

# Replicator dynamics equation

- Dynamics of  $i$  strategists


$$\frac{dn_i}{dt} = \dot{n}_i = n_i(\beta + f_i) \quad (8)$$

- Dynamics of whole population

$$\dot{N} = N(\beta + \bar{f}) \quad (9)$$

- Dynamics of proportions  $p_i = \frac{n_i}{N} \rightarrow n_i = p_i N$

$$\dot{n}_i = \dot{p}_i N + p_i \dot{N}$$



- $n_i$ : no. individvs of strategy  $i$
- $N$ : total no. of individvs
- $p_i$ : proportions of strategy  $i$
- $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $\beta$ : background reprodn rate

# Replicator dynamics equation

- Dynamics of  $i$  strategists

$$\frac{dn_i}{dt} = \dot{n}_i = n_i(\beta + f_i) \quad (8)$$


- Dynamics of whole population

$$\dot{N} = N(\beta + \bar{f}) \quad (9)$$

- Dynamics of proportions  $p_i = \frac{n_i}{N} \rightarrow \dot{n}_i = p_i \dot{N}$

$$\dot{n}_i = \dot{p}_i N + p_i \dot{N}$$

$$\dot{p}_i N = \dot{n}_i - p_i \dot{N}$$




- $n_i$ : no. individvs of strategy  $i$
- $N$ : total no. of individvs
- $p_i$ : proportions of strategy  $i$
- $f_i(\mathbf{p})$ : fitness effect of strategy  $i$
- $\beta$ : background reprodn rate

# Replicator dynamics equation

- Dynamics of proportions

$$\dot{p}_i N = \dot{n}_i - p_i \dot{N}$$

$$\dot{p}_i N = (\beta + f_i) n_i - p_i N (\beta + \bar{f})$$



$$\dot{n}_i = n_i (\beta + f_i)$$

$$\dot{N} = N (\beta + \bar{f})$$

# Replicator dynamics equation

- Dynamics of proportions


$$\dot{p}_i N = \dot{n}_i - p_i \dot{N}$$

$$\dot{p}_i N = (\beta + f_i) n_i - p_i N (\beta + \bar{f})$$

$$\dot{p}_i = (\beta + f_i) \frac{n_i}{N} - p_i (\beta + \bar{f})$$

$$= (\beta + f_i) p_i - p_i (\beta + \bar{f})$$

$$\dot{p}_i = p_i (f_i - \bar{f})$$




$$\dot{n}_i = n_i (\beta + f_i)$$

$$\dot{N} = N (\beta + \bar{f})$$

# Replicator dynamics equation

- Replicator dynamics

$$\dot{p}_i = p_i(f_i - \bar{f}) \quad (10)$$




# Replicator dynamics equation

- Replicator dynamics

$$\dot{p}_i = p_i(f_i - \bar{f}) \quad (10)$$

- For two strategies

$$\dot{p}_1 = p_1(1 - p_1)(f_1 - f_2) \quad (11)$$



# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

		 1	 2
 1	COOPERATE	$\frac{V}{2}$	$V$
	DEFECT	$V$	$\frac{(V-C)}{2}$

# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

- Fitness effect of hawk strategy?

		2	
1		COOPERATE	DEFECT
1		 COOPERATE	 DEFECT
 HAWK		$V/2$	$V$
 DOVE		$V/2$	0
 HAWK		0	$(V-C)/2$
 DOVE		$V$	$(V-C)/2$

# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

- Fitness effect of hawk strategy? Hint:  $f_i(\mathbf{p})$

		 1	 2
 1	COOPERATE	 $V/2$	 $V$
	DEFECT	 $V/2$	 $0$

# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

- Fitness effect of hawk strategy? Hint:  $f_i(\mathbf{p})$

$$f_H(\mathbf{p}) = p_H \frac{V - C}{2} + p_D V$$

	 1	 2
 1	 COOPERATE	 DEFECT
 2	$V/2$	$V$
	$V/2$	0

	 1	 2
 1	 COOPERATE	 DEFECT
 2	$V/2$	$V$
	$V/2$	0

	 1	 2
 1	 COOPERATE	 DEFECT
 2	$V/2$	$V$
	$V/2$	0

# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

- Fitness effect of hawk strategy? Hint:  $f_i(\mathbf{p})$

$$f_H(\mathbf{p}) = p_H \frac{V - C}{2} + p_D V$$

		 <b>1</b>	
		 <b>2</b>	 <b>2</b>
 <b>2</b>		<b>COOPERATE</b>	<b>DEFECT</b>
 <b>2</b>	 <b>1</b>	$V/2$	$V$
 <b>1</b>	 <b>2</b>	$V/2$	$(V-C)/2$
		$0$	$0$
		$(V-C)/2$	$(V-C)/2$

# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

- Fitness effect of hawk strategy? Hint:  $f_i(\mathbf{p})$

$$f_H(\mathbf{p}) = p_H \frac{V - C}{2} + p_D V$$

$$f_D(\mathbf{p}) = p_H 0 + p_D \frac{V}{2}$$

 1 	 COOPERATE  DEFECT	 2 
		 $V/2$  0
	 0  $(V-C)/2$	

# Hawk-Dove replicator dynamics

- Replicator dynamics

$$\dot{p}_H = p_H(1 - p_H)(f_H - f_D)$$

- Fitness effect of hawk strategy? Hint:  $f_i(\mathbf{p})$

$$f_H(\mathbf{p}) = p_H \frac{V - C}{2} + p_D V$$

$$f_D(\mathbf{p}) = p_H 0 + p_D \frac{V}{2}$$

- Sub  $(1 - p_H) = p_D$ , rearrange

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

	$\frac{V}{2}$	$V$
	$0$	$\frac{(V-C)}{2}$

# Hawk-Dove replicator analysis

$$\dot{p}_H = \frac{dp_H}{dt} = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

- Equilibrium?

	 1	 2
 COOPERATE	$V/2$	$V$
 DEFECT	$V/2$	$0$

	 1	 2
 COOPERATE	$V/2$	$V$
 DEFECT	$V/2$	$0$

# Hawk-Dove replicator analysis

$$\dot{p}_H = \frac{dp_H}{dt} = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

- Equilibrium? Solve  $\frac{dp_H}{dt} = 0$

	<b>COOPERATE</b>	<b>DEFECT</b>
	<b>DEFECT</b>	<b>COOPERATE</b>

A game matrix for the Hawk-Dove game. The rows represent Player 1's strategies: COOPERATE (top) and DEFECT (bottom). The columns represent Player 2's strategies: COOPERATE (left) and DEFECT (right). Payoffs are listed in the matrix cells. The payoffs are: (Player 1, Player 2) = (COOP, COOP) → (V/2, V/2); (Player 1, Player 2) = (COOP, DEFECT) → (V/2, 0); (Player 1, Player 2) = (DEFECT, COOP) → (0, V/2); (Player 1, Player 2) = (DEFECT, DEFECT) → (V-C/2, V-C/2).

# Hawk-Dove replicator analysis

$$\dot{p}_H = \frac{dp_H}{dt} = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

- Equilibrium? Solve  $\frac{dp_H}{dt} = 0$

$$p_H^* = 0, 1, \frac{V}{C}$$

	 1	 2
 1	 COOPERATE $V/2$	 DEFECT $V$
 2	 COOPERATE $V/2$	 DEFECT $0$
	 1	 COOPERATE $(V-C)/2$

# Hawk-Dove replicator analysis

$$\dot{p}_H = \frac{dp_H}{dt} = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

- Equilibrium? Solve  $\frac{dp_H}{dt} = 0$

$$p_H^* = 0, 1, \frac{V}{C}$$

- Asymptotic stability?

	 1	 2
 1	 COOPERATE $V/2$	 DEFECT $V$
 2	 COOPERATE $V/2$	 DEFECT $0$
	 1	 DEFECT $(V-C)/2$

# Hawk-Dove replicator analysis

$$\dot{p}_H = \frac{dp_H}{dt} = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

- Equilibrium? Solve  $\frac{dp_H}{dt} = 0$

$$p_H^* = 0, 1, \frac{V}{C}$$

- Asymptotic stability?

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

		2 COOPERATE	DEFECT
1 COOPERATE	V/2	V	(V-C)/2
0	V/2	0	(V-C)/2
DEFECT	V	(V-C)/2	

# Hawk-Dove replicator analysis

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

		2	
		COOPERATE	DEFECT
		1	2
1	COOPERATE	 $\frac{V}{2}$	 $V$
	DEFECT	 $\frac{V}{2}$	 $0$
2	COOPERATE	 $0$	 $\frac{(V-C)}{2}$
	DEFECT	 $V$	 $\frac{(V-C)}{2}$

$$p_H^* = 0, 1, \frac{V}{C}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

# Hawk-Dove replicator analysis

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

$$(uvw)' = u'vw + uv'w + uwv'$$

		Hawk	
		Cooperate	Defect
Dove	Cooperate	$\frac{V}{2}$	$V$
	Defect	$\frac{V}{2}$	0
1	Defect	0	$\frac{(V-C)}{2}$
		$V$	$\frac{(V-C)}{2}$

$$p_H^* = 0, 1, \frac{V}{C}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

# Hawk-Dove replicator analysis

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

$$(uvw)' = u'vw + uv'w + uwv'$$

$$\begin{aligned}\frac{d\dot{p}_H}{dp_H} &= (1 - p_H) \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad - p_H \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad + p_H (1 - p_H) \left( \frac{V - C}{2} - \frac{V}{2} \right)\end{aligned}$$

		2	
	COOPERATE	$\frac{V}{2}$	$V$
	COOPERATE	$\frac{V}{2}$	0
1	DEFECT	0	$\frac{(V-C)}{2}$
		$V$	$\frac{(V-C)}{2}$

$$p_H^* = 0, 1, \frac{V}{C}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

# Hawk-Dove replicator analysis

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

$$(uvw)' = u'vw + uv'w + uwv'$$

$$\begin{aligned}\frac{d\dot{p}_H}{dp_H} &= (1 - p_H) \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad - p_H \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad + p_H (1 - p_H) \left( \frac{V - C}{2} - \frac{V}{2} \right)\end{aligned}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=0} = \frac{V}{2} > 0$$

		2	
	COOPERATE	V/2	V
1	COOPERATE	V/2	0
	DEFECT	0	(V-C)/2
		V	(V-C)/2

$$p_H^* = 0, 1, \frac{V}{C}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

# Hawk-Dove replicator analysis

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

$$(uvw)' = u'vw + uv'w + uwv'$$

$$\begin{aligned}\frac{d\dot{p}_H}{dp_H} &= (1 - p_H) \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad - p_H \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad + p_H (1 - p_H) \left( \frac{V - C}{2} - \frac{V}{2} \right)\end{aligned}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=0} = \frac{V}{2} > 0$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=1} = -\frac{V - C}{2} > 0$$

		2	
	COOPERATE	V/2	V
1	DEFECT	V/2	0
		0	(V-C)/2
		V	(V-C)/2

$$p_H^* = 0, 1, \frac{V}{C}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

# Hawk-Dove replicator analysis

$$\dot{p}_H = p_H(1 - p_H) \left( p_H \frac{V - C}{2} + (1 - p_H) \frac{V}{2} \right)$$

$$(uvw)' = u'vw + uv'w + uwv'$$

$$\begin{aligned}\frac{d\dot{p}_H}{dp_H} &= (1 - p_H) \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad - p_H \left( \frac{V - C}{2} p_H + \frac{V}{2} (1 - p_H) \right) \\ &\quad + p_H (1 - p_H) \left( \frac{V - C}{2} - \frac{V}{2} \right)\end{aligned}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=0} = \frac{V}{2} > 0$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=1} = -\frac{V - C}{2} > 0$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=\frac{V}{C}} = -\frac{V}{2} \left( 1 - \frac{V}{C} \right) < 0$$

	h D	2 COOPERATE	2 DEFECT
1 COPRO	V/2 0	V 0	(V-C)/2 (V-C)/2
D EFFECT	V 0	(V-C)/2 (V-C)/2	

$$p_H^* = 0, 1, \frac{V}{C}$$

$$\frac{d\dot{p}_H}{dp_H} \Big|_{p_H=p_H^*} < 0$$

# Replicator dynamics - A technical point

ESS → asymptotically stable → Nash equilibrium

- e.g. variants on rock-scissors-paper have stable replicator dynamics but are not ESS

# Replicator dynamics - A technical point

ESS → asymptotically stable → Nash equilibrium

- e.g. variants on rock-scissors-paper have stable replicator dynamics but are not ESS
- For 2-player 2-strategy games they are all equivalent

# Adaptive dynamics

- Adaptive dynamics (1990s onwards)
  - About evolution of some continuous ‘trait’ in a ‘resident population’ by successive invasions of a mutant traits
  - ‘Ecoevolutionary’: feedbacks between ecological dynamics and evolutionary dynamics
    - In replicator dynamics, fitness a function of *proportions*
    - In AD, fitness a function of absolute population size
  - Good introduction ‘Hitchhiker’s guide to Adaptive Dynamics’ (Bränström et al. 2013)

# Adaptive dynamics

- Adaptive dynamics (1990s onwards)
  - About evolution of some continuous ‘trait’ in a ‘resident population’ by successive invasions of a mutant traits
  - ‘Ecoevolutionary’: feedbacks between ecological dynamics and evolutionary dynamics
    - In replicator dynamics, fitness a function of *proportions*
    - In AD, fitness a function of absolute population size
  - Good introduction ‘Hitchhiker’s guide to Adaptive Dynamics’ (Bränström et al. 2013)
- Key assumptions of Adaptive Dynamics
  - ① Clonal reproduction
  - ② Separation of timescales
    - Resident population in ecological equilibrium when new mutant appears
  - ③ Small mutational steps and few mutants
    - Mutant does not affect fitness of residents but residents’ strategy affects fitness of mutant

# Adaptive dynamics

- Adaptive dynamics (1990s onwards)
  - About evolution of some continuous ‘trait’ in a ‘resident population’ by successive invasions of a mutant traits
  - ‘Ecoevolutionary’: feedbacks between ecological dynamics and evolutionary dynamics
    - In replicator dynamics, fitness a function of *proportions*
    - In AD, fitness a function of absolute population size
  - Good introduction ‘Hitchhiker’s guide to Adaptive Dynamics’ (Bränström et al. 2013)
- Key assumptions of Adaptive Dynamics
  - ① Clonal reproduction
  - ② Separation of timescales
    - Resident population in ecological equilibrium when new mutant appears
  - ③ Small mutational steps and few mutants
    - Mutant does not affect fitness of residents but residents’ strategy affects fitness of mutant
- Clarifies meaning of evolutionary trajectory and end-points

# Adaptive dynamics example - migratory birds

Reference - Johansson & Jonzén (2012)


- Timing of arrival back from migration



# Adaptive dynamics example - migratory birds

Reference - Johansson & Jonzén (2012)


- Timing of arrival back from migration
  - Arrive too early - cold, low food



# Adaptive dynamics example - migratory birds

Reference - Johansson & Jonzén (2012)


- Timing of arrival back from migration
  - Arrive too early - cold, low food
  - Arrive too late - not enough time to nest, lay eggs, raise chicks before winter



# Adaptive dynamics example - migratory birds

Reference - Johansson & Jonzén (2012)


- Timing of arrival back from migration
  - Arrive too early - cold, low food
  - Arrive too late - not enough time to nest, lay eggs, raise chicks before winter
  - So obviously best arrival time is at peak?



# Adaptive dynamics example - migratory birds

Reference - Johansson & Jonzén (2012)


- Timing of arrival back from migration
  - Arrive too early - cold, low food
  - Arrive too late - not enough time to nest, lay eggs, raise chicks before winter
  - So obviously best arrival time is at peak?
- But there is competition for limited nesting territories
  - Those who arrive earlier have a better chance at obtaining and defending a territory
  - If everyone else arrives at peak, may be better overall for an individual to arrive a bit earlier?



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)

- Combine
  - Overall reproduction is hump-shaped with arrival time





# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)

- Combine


- Overall reproduction is hump-shaped with arrival time
- Greater chance to obtain breeding territory with earlier arrival



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)


- Combine
  - Overall reproduction is hump-shaped with arrival time
  - Greater chance to obtain breeding territory with earlier arrival
  - Fitness function of mutants when all others at reproduction peak



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)


- Combine
  - Overall reproduction is hump-shaped with arrival time
  - Greater chance to obtain breeding territory with earlier arrival
  - Fitness function of mutants when all others at reproduction peak
- Successive invasions
  - Fitness of mutant who arrives earlier is higher – can invade



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)


- Combine
  - Overall reproduction is hump-shaped with arrival time
  - Greater chance to obtain breeding territory with earlier arrival
  - Fitness function of mutants when all others at reproduction peak
- Successive invasions
  - Fitness of mutant who arrives earlier is higher – can invade
  - Invader grows, establishes, new mutant fitness landscape



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)


- Combine
  - Overall reproduction is hump-shaped with arrival time
  - Greater chance to obtain breeding territory with earlier arrival
  - Fitness function of mutants when all others at reproduction peak
- Successive invasions
  - Fitness of mutant who arrives earlier is higher – can invade
  - Invader grows, establishes, new mutant fitness landscape



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)


- Combine
  - Overall reproduction is hump-shaped with arrival time
  - Greater chance to obtain breeding territory with earlier arrival
  - Fitness function of mutants when all others at reproduction peak
- Successive invasions
  - Fitness of mutant who arrives earlier is higher – can invade
  - Invader grows, establishes, new mutant fitness landscape



# Adaptive dynamics example

Reference - Johansson & Jonzén (2012)

- Combine
  - Overall reproduction is hump-shaped with arrival time
  - Greater chance to obtain breeding territory with earlier arrival
  - Fitness function of mutants when all others at reproduction peak
- Successive invasions
  - Fitness of mutant who arrives earlier is higher – can invade
  - Invader grows, establishes, new mutant fitness landscape
- End-point where no further invasions possible




# Adaptive dynamics clarifies concepts

- AD is interested in the two concepts from example above:
  - ① Which strategy will evolutionary trajectory head towards?  
*Convergence stability*
  - ② Can population pursuing strategy be invaded by alternative strategy?  
*Evolutionary stability*
- Not every ESS is an evolutionary attractor
  - e.g. gregarious behaviour (Eshel 1983)
- *Continuously stable strategy*: An evolutionarily singular strategy that is both evolutionarily stable and convergence stable

# Adaptive dynamics approach

- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
- Adaptive dynamics approach works ‘backwards’ from end-point




# Adaptive dynamics approach

- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
  - Adaptive dynamics approach works ‘backwards’ from end-point
- ① Find evolutionarily singular strategy  $x^*$

$$\bullet g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = 0$$

$x^*$  is where mutant fitness gradient is 0



# Adaptive dynamics approach


- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
- Adaptive dynamics approach works ‘backwards’ from end-point

- ① Find evolutionarily singular strategy  $x^*$

- $$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = 0$$

$x^*$  is where mutant fitness gradient is 0

- ② Is  $x^*$  an *evolutionarily stable strategy*?



# Adaptive dynamics approach

- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
- Adaptive dynamics approach works ‘backwards’ from end-point

- ① Find evolutionarily singular strategy  $x^*$


- $$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = 0$$

$x^*$  is where mutant fitness gradient is 0

- ② Is  $x^*$  an *evolutionarily stable strategy*?

- $$\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$$

fitness gradient at  $x^*$  a maximum wrt change in mutant trait



# Adaptive dynamics approach

- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
- Adaptive dynamics approach works ‘backwards’ from end-point

- ① Find evolutionarily singular strategy  $x^*$

- $$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = 0$$


$x^*$  is where mutant fitness gradient is 0

- ② Is  $x^*$  an *evolutionarily stable strategy*?

- $$\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$$

fitness gradient at  $x^*$  a maximum wrt change in mutant trait

- ③ Is  $x^*$  a *convergence stable strategy*?



# Adaptive dynamics approach

- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
- Adaptive dynamics approach works ‘backwards’ from end-point

- ① Find evolutionarily singular strategy  $x^*$

- $$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = 0$$

$x^*$  is where mutant fitness gradient is 0

- ② Is  $x^*$  an *evolutionarily stable strategy*?


- $$\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$$

fitness gradient at  $x^*$  a maximum wrt change in mutant trait

- ③ Is  $x^*$  a *convergence stable strategy*?

- $$\frac{dg_x}{dx} \Big|_{x=x^*} < 0$$

fitness gradient at  $x^*$  a maximum wrt change in resident trait



# Adaptive dynamics approach

- Call fitness  $w(y, x)$ , where  $y$  is mutant trait and  $x$  is resident trait
- Adaptive dynamics approach works ‘backwards’ from end-point

- ① Find evolutionarily singular strategy  $x^*$

- $$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = 0$$

$x^*$  is where mutant fitness gradient is 0

- ② Is  $x^*$  an *evolutionarily stable strategy*?


- $$\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$$

fitness gradient at  $x^*$  a maximum wrt change in mutant trait

- ③ Is  $x^*$  a *convergence stable strategy*?

- $$\frac{dg_x}{dx} \Big|_{x=x^*} < 0$$


fitness gradient at  $x^*$  a maximum wrt change in resident trait



All combinations possible

## Aside: Adaptive dynamics and speciation

- How could an evolutionarily singular strategy be convergence stable but not evolutionarily stable?




- Because AD assumes clonal reproduction, leads to polymorphism or 'speciation'
- In reality however, gene flow prevents trait divergence
- See Dieckmann & Doebeli (1999) versus Gavrilets (2005)

# Adaptive dynamics worked example - preliminaries

First consider population all following same strategy – *resident strategy*

- No. offspring fnc arrival time

$$R(x) = R_0 e^{-\frac{x^2}{2}} \quad (12)$$




# Adaptive dynamics worked example - preliminaries

First consider population all following same strategy – *resident strategy*

- No. offspring fnc arrival time

$$R(x) = R_0 e^{-\frac{x^2}{2}} \quad (12)$$



- Probability of obtaining a territory assuming  $n_t > K$


$$p = \frac{K}{n_t} \quad (13)$$

# Adaptive dynamics worked example - preliminaries

First consider population all following same strategy – *resident strategy*

- No. offspring fnc arrival time

$$R(x) = R_0 e^{-\frac{x^2}{2}} \quad (12)$$



- Probability of obtaining a territory assuming  $n_t > K$

$$p = \frac{K}{n_t} \quad (13)$$

- Population dynamics

$$n_{t+1} = pR(x) n_t \quad (14)$$

# Adaptive dynamics worked example - separation of timescales

- Population dynamics

$$n_{t+1} = pR(x) n_t \quad (15)$$

# Adaptive dynamics worked example - separation of timescales

- Population dynamics

$$n_{t+1} = pR(x) n_t \quad (15)$$

- Assume that evolutionary dynamics happening on a slower timescale than population dynamics

# Adaptive dynamics worked example - separation of timescales

- Population dynamics

$$n_{t+1} = pR(x) n_t \quad (15)$$


- Assume that evolutionary dynamics happening on a slower timescale than population dynamics
- Therefore assume population at steady state

$$n_{t+1} = n_t = n^* \quad (16)$$

$$p_r = \frac{1}{R(x)} = \frac{K}{n^*(x)} \quad (17)$$

# Adaptive dynamics worked example - very few mutants


- Residents' probability of obtaining a territory  $p_r = \frac{1}{R(x)}$
- Assume only one mutation at a time
- What happens if a mutant with a different arrival time  $y$  arises?



# Adaptive dynamics worked example - very few mutants

- Residents' probability of obtaining a territory  $p_r = \frac{1}{R(x)}$
- Assume* only one mutation at a time
- What happens if a mutant with a different arrival time  $y$  arises?
- Assume* large enough population, doesn't affect residents' fitness

$$\text{keep } p_r = \frac{1}{R(x)} \quad (18)$$




# Adaptive dynamics worked example - very few mutants

- Residents' probability of obtaining a territory  $p_r = \frac{1}{R(x)}$
- Assume only one mutation at a time
- What happens if a mutant with a different arrival time  $y$  arises?
- Assume large enough population, doesn't affect residents' fitness

$$\text{keep } p_r = \frac{1}{R(x)} \quad (18)$$

- But mutant definitely affected by residents
  - For purpose of example, assume a linear relationship truncated between 0 and 1

$$p(y, x) = p_r (1 + a (x - y)) \quad (19)$$



## Adaptive dynamics worked example - invasion fitness

- Now have two components:  $R(y)$  and  $p(y, x)$
- Recall the population dynamics

$$n_{t+1} = pR(x) n_t \quad (20)$$

## Adaptive dynamics worked example - invasion fitness

- Now have two components:  $R(y)$  and  $p(y, x)$
- Recall the population dynamics

$$n_{t+1} = pR(x) n_t \quad (20)$$

- Growth rate is that  $pR(x)$  term
  - Growth rate of resident  $pR(x) = 1$
  - Growth rate of mutant  $p(y, x)R(y)$

## Adaptive dynamics worked example - invasion fitness

- Now have two components:  $R(y)$  and  $p(y, x)$
- Recall the population dynamics

$$n_{t+1} = pR(x) n_t \quad (20)$$

- Growth rate is that  $pR(x)$  term
  - Growth rate of resident  $pR(x) = 1$
  - Growth rate of mutant  $p(y, x)R(y)$
- Fitness of mutant with  $y$  arrival time in a resident population with  $x$  therefore

$$w(y, x) = p(y, x)R(y) \quad (21)$$

## Adaptive dynamics worked example - invasion fitness

- Now have two components:  $R(y)$  and  $p(y, x)$
- Recall the population dynamics

$$n_{t+1} = pR(x) n_t \quad (20)$$

- Growth rate is that  $pR(x)$  term
  - Growth rate of resident  $pR(x) = 1$
  - Growth rate of mutant  $p(y, x)R(y)$
- Fitness of mutant with  $y$  arrival time in a resident population with  $x$  therefore


$$w(y, x) = p(y, x)R(y) \quad (21)$$

- Notice the assumptions
  - Separation of timescales - resident population at ecological equilibrium, one mutant at a time
  - Resident fitness not affected by mutant, but mutant fitness affected by resident strategy

# Adaptive dynamics worked example - find singular strategy

Find evolutionarily singular strategy  $x^*$

- $g_x = \frac{\partial w(y,x)}{\partial y} \Big|_{y=x} = 0$
- Using:  
 $w(y, x) = p(y, x)R(y)$



# Adaptive dynamics worked example - find singular strategy

Find evolutionarily singular strategy  $x^*$


- $g_x = \frac{\partial w(y,x)}{\partial y} \Big|_{y=x} = 0$

- Using:

$$w(y, x) = p(y, x)R(y)$$

- Find the partial derivative wrt  $y$

$$\frac{\partial w(y, x)}{\partial y} = -R(y) (ap_r + yp(y, x)) \quad (22)$$



# Adaptive dynamics worked example - find singular strategy

Find evolutionarily singular strategy  $x^*$

- $g_x = \frac{\partial w(y,x)}{\partial y} \Big|_{y=x} = 0$

- Using:


$$w(y, x) = p(y, x)R(y)$$

- Find the partial derivative wrt  $y$

$$\frac{\partial w(y, x)}{\partial y} = -R(y) (ap_r + yp(y, x)) \quad (22)$$

- Evaluate partial derivative at the resident strategy

$$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = -R(x) (ap_r + xp_r) \quad (23)$$



# Adaptive dynamics worked example - find singular strategy

Find evolutionarily singular strategy  $x^*$

- $g_x = \frac{\partial w(y,x)}{\partial y} \Big|_{y=x} = 0$

- Using:

$$w(y, x) = p(y, x)R(y)$$

- Find the partial derivative wrt  $y$


$$\frac{\partial w(y, x)}{\partial y} = -R(y) (ap_r + yp(y, x)) \quad (22)$$

- Evaluate partial derivative at the resident strategy

$$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = -R(x) (ap_r + xp_r) \quad (23)$$

- $x^*$  is when  $g_x = 0$


$$\rightarrow x^* = -a \quad (24)$$



# Adaptive dynamics worked example - is singular strategy evolutionarily stable?

Is  $x^*$  an *evolutionarily stable strategy*?

- $\frac{\partial^2 w(y,x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$
- Using:  $x^* = -a$




# Adaptive dynamics worked example - is singular strategy evolutionarily stable?

Is  $x^*$  an *evolutionarily stable strategy*?

- $\frac{\partial^2 w(y,x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$

- Using:  $x^* = -a$
- Find the second partial derivative wrt  $y$

$$\frac{\partial^2 w(y,x)}{\partial y^2} = R(y) (p(y,x)(y^2 - 1) + 2ayp_r) \quad (25)$$



# Adaptive dynamics worked example - is singular strategy evolutionarily stable?

Is  $x^*$  an *evolutionarily stable strategy*?


- $\frac{\partial^2 w(y,x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$

- Using:  $x^* = -a$
- Find the second partial derivative wrt  $y$

$$\frac{\partial^2 w(y,x)}{\partial y^2} = R(y) (p(y,x)(y^2 - 1) + 2ayp_r) \quad (25)$$

- Evaluate second partial derivative at the resident strategy

$$\frac{\partial^2 w(y,x)}{\partial y^2} \Big|_{y=x} = R(x)p_r (x^2 - 1 + 2ax) \quad (26)$$



# Adaptive dynamics worked example - is singular strategy evolutionarily stable?

Is  $x^*$  an *evolutionarily stable strategy*?

- $\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} < 0$

- Using:  $x^* = -a$
- Find the second partial derivative wrt  $y$


$$\frac{\partial^2 w(y, x)}{\partial y^2} = R(y) (p(y, x)(y^2 - 1) + 2ayp_r) \quad (25)$$

- Evaluate second partial derivative at the resident strategy

$$\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x}} = R(x)p_r (x^2 - 1 + 2ax) \quad (26)$$

- Evaluate at  $x^*$


$$\frac{\partial^2 w(y, x)}{\partial y^2} \Big|_{\substack{y=x \\ x=x^*}} = R(-a)p_r (-1 - a^2) < 0 \quad (27)$$



# Adaptive dynamics worked example - is singular strategy convergence stable?

Is  $x^*$  a *convergence stable strategy*?

- $\frac{dg_x}{dx} \Big|_{x=x^*} < 0$
- Using:  $x^* = -a$




# Adaptive dynamics worked example - is singular strategy convergence stable?

Is  $x^*$  a *convergence stable strategy*?

- $\frac{dg_x}{dx} \Big|_{x=x^*} < 0$
- Using:  $x^* = -a$
- Recall

$$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = -p_r R(x) (a + x) \quad (28)$$



# Adaptive dynamics worked example - is singular strategy convergence stable?


Is  $x^*$  a *convergence stable strategy*?

- $\frac{dg_x}{dx} \Big|_{x=x^*} < 0$
- Using:  $x^* = -a$
- Recall

$$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = -p_r R(x) (a + x) \quad (28)$$

- Find the derivative of  $g(x)$  wrt  $x$

$$\frac{dg_x}{dx} = -p_r \left( \frac{dR(x)}{dx} (a + x) + R(x) \right) \quad (29)$$



# Adaptive dynamics worked example - is singular strategy convergence stable?

Is  $x^*$  a *convergence stable strategy*?

- $\frac{dg_x}{dx} \Big|_{x=x^*} < 0$
- Using:  $x^* = -a$
- Recall


$$g_x = \frac{\partial w(y, x)}{\partial y} \Big|_{y=x} = -p_r R(x) (a + x) \quad (28)$$

- Find the derivative of  $g(x)$  wrt  $x$

$$\frac{dg_x}{dx} = -p_r \left( \frac{dR(x)}{dx} (a + x) + R(x) \right) \quad (29)$$



- Evaluate at  $x^*$

$$\frac{dg_x}{dx} \Big|_{x=x^*} = -p_r R(-a) < 0 \quad (30)$$



# Adaptive dynamics worked example - meaning

- One singular strategy  $x^* = -a$ 
  - In example right, I've set  $a = 2$
- Continuously stable strategy
- What does it mean biologically?





# Adaptive dynamics worked example - meaning

- One singular strategy  $x^* = -a$ 
  - In example right, I've set  $a = 2$
- Continuously stable strategy
- What does it mean biologically?

Species	Date	Location	Conditions	Numbers	Source
(x) Mortality during spring migration					
Various species (> 23 species)	April 1881	Off Louisiana coast <sup>*</sup>	Gale	'Many thousands'	Frazar (1881)
Lapland Longspurs <i>Calcarius lapponicus</i>	March 1904	Minnesota-Iowa	Snowstorm	1.5 million	Roberts (1907a, 1907b)
Mainly Lapland Longspurs <i>Calcarius lapponicus</i>	February 1922	Nebraska	Snowstorm	'Thousands'	Reed (1922), Sibley (1922)
Magnolia Warblers <i>Dendroica magnolia</i> and others (39 species)	May 1951	Off Texas coast <sup>*</sup>	Rainstorm	> 10 000	James (1956)
Ducks, geese and swans	April 1954	Wisconsin	Hailstorm	'Many'	Hochbaum (1955)
Various (> 14 species)	May 1954	Minnesota	Snowstorm	> 175	Frenell and Marshall (1954)



"In each documented example, the migrants could have avoided the cold spell if they had arrived in breeding areas some days later than they did" Newton (2007, *Ibis*)



# Prisoner's dilemma

Two criminals interrogated separately by police. If both stay silent, get a lesser charge and 1 year jail. But police make an offer - testify against the other, and if the other doesn't testify, you go free and they 3 years jail. But if both testify against each other, both get 2 years jail.

Should the criminals cooperate with each other and stay silent?

		1	2
		COOPERATE	DEFECT
1	COOPERATE	-1 -1	0 -3
2	DEFECT	0 -3	-2

# Prisoner's dilemma

Two criminals interrogated separately by police. If both stay silent, get a lesser charge and 1 year jail. But police make an offer - testify against the other, and if the other doesn't testify, you go free and they 3 years jail. But if both testify against each other, both get 2 years jail.

Should the criminals cooperate with each other and stay silent?

- If other is defecting, should defect  
 $f(D, D) > f(C, D)$

		Player 2	
		COOPERATE	DEFECT
Player 1	COOPERATE	-1	0
	DEFECT	-1	-3

		Player 2	
		COOPERATE	DEFECT
Player 1	COOPERATE	-1	0
	DEFECT	-1	-3

		Player 2	
		COOPERATE	DEFECT
Player 1	COOPERATE	-1	0
	DEFECT	-1	-3

		Player 2	
		COOPERATE	DEFECT
Player 1	COOPERATE	-1	0
	DEFECT	-1	-3

# Prisoner's dilemma

Two criminals interrogated separately by police. If both stay silent, get a lesser charge and 1 year jail. But police make an offer - testify against the other, and if the other doesn't testify, you go free and they 3 years jail. But if both testify against each other, both get 2 years jail.

Should the criminals cooperate with each other and stay silent?

- If other is defecting, should defect  
 $f(D, D) > f(C, D)$
- If other is cooperating, should still defect  
 $f(C, C) < f(D, C)$

	2	
	COOPERATE	DEFECT
COOPERATE	-1	0
DEFECT	-1	-3
1	-3	-2
0	0	-2

# Prisoner's dilemma

Two criminals interrogated separately by police. If both stay silent, get a lesser charge and 1 year jail. But police make an offer - testify against the other, and if the other doesn't testify, you go free and they 3 years jail. But if both testify against each other, both get 2 years jail.

Should the criminals cooperate with each other and stay silent?

- If other is defecting, should defect  
 $f(D, D) > f(C, D)$
- If other is cooperating, should still defect  
 $f(C, C) < f(D, C)$
- Defect is only ESS / Nash equilibrium

		p	d
		COOPERATE	2
1	COOPERATE	-1	0
1	DEFECT	-1	-3
		DEFECT	-3
			-2

# Generalised Prisoner's Dilemma and replicator dynamics

		2	
		COOPERATE	DEFECT
1		b	d
		b	c
		c	a
		d	a

$$d > b > a > c$$

# Generalised Prisoner's Dilemma and replicator dynamics

- Fitness effects:

$$f_C = p_C b + p_D c$$

$$f_D = p_C d + p_D a$$

		2	
		COOPERATE	DEFECT
1		b	d
COOPERATE		b	c
DEFECT		c	a
		d	

$$d > b > a > c$$

# Generalised Prisoner's Dilemma and replicator dynamics

- Fitness effects:

$$f_C = p_C b + p_D c$$

$$f_D = p_C d + p_D a$$

- Replicator dynamics

$$\dot{p}_C = p_C(1 - p_C)(f_C - f_D)$$

$$= p_C(1 - p_C)\{b p_C + c(1 - p_C) - d p_C - (1 - p_C)a\}$$

	2	
	COOPERATE	DEFECT
1	b	d
COOPERATE	b	c
DEFECT	c	a
	d	a

$$d > b > a > c$$

# Generalised Prisoner's Dilemma and replicator dynamics

- Fitness effects:

$$f_C = p_C b + p_D c$$

$$f_D = p_C d + p_D a$$

- Replicator dynamics

$$\dot{p}_C = p_C(1 - p_C)(f_C - f_D)$$

$$= p_C(1 - p_C)\{b p_C + c(1 - p_C) - d p_C - (1 - p_C)a\}$$

- Steady states:

$$p_C^* = 0, 1, \frac{a - c}{a - c + b - d}$$

		2	
		COOPERATE	DEFECT
1	COOPERATE	b	d
	DEFECT	c	a
		d	a

$$d > b > a > c$$

# Generalised Prisoner's Dilemma and replicator dynamics

- Fitness effects:

$$f_C = p_C b + p_D c$$

$$f_D = p_C d + p_D a$$

- Replicator dynamics

$$\dot{p}_C = p_C(1 - p_C)(f_C - f_D)$$

$$= p_C(1 - p_C)\{b p_C + c(1 - p_C) - d p_C - (1 - p_C)a\}$$

- Steady states:  $p_C^* = 0, 1$

		2	
		COOPERATE	DEFECT
1	COOPERATE	b	d
	DEFECT	c	a
		d	a

$$d > b > a > c$$

# Generalised Prisoner's Dilemma and replicator dynamics

- Fitness effects:

$$f_C = p_C b + p_D c$$

$$f_D = p_C d + p_D a$$

- Replicator dynamics

$$\dot{p}_C = p_C(1 - p_C)(f_C - f_D)$$

$$= p_C(1 - p_C)\{b p_C + c(1 - p_C) - d p_C - (1 - p_C)a\}$$

- Steady states:  $p_C^* = 0, 1$

- Check asymptotic stability

$$\frac{d\dot{p}_C}{dp_C} = (1 - p_C)\{ \} - p_C\{ \} + p_C(1 - p_C)\frac{d\{ \}}{dp_C}$$

$$d > b > a > c$$

	2	
	COOPERATE	DEFECT
1	COOPERATE	b d
	DEFECT	b c c a d a

# Generalised Prisoner's Dilemma and replicator dynamics

- Fitness effects:

$$f_C = p_C b + p_D c$$
$$f_D = p_C d + p_D a$$

- Replicator dynamics

$$\dot{p}_C = p_C(1 - p_C)(f_C - f_D)$$
$$= p_C(1 - p_C)\{b p_C + c(1 - p_C) - d p_C - (1 - p_C)a\}$$

- Steady states:  $p_C^* = 0, 1$
- Check asymptotic stability

$$\frac{d\dot{p}_C}{dp_C} = (1 - p_C)\{ \} - p_C\{ \} + p_C(1 - p_C)\frac{d\{ \}}{dp_C}$$

$$d > b > a > c$$

$$\left. \frac{d\dot{p}_C}{dp_C} \right|_{p_C=1} = -(b - d) > 0; \text{ and } \left. \frac{d\dot{p}_C}{dp_C} \right|_{p_C=0} = (c - a) < 0$$

	1	2
	COOPERATE	DEFECT
COOPERATE	b	d
DEFECT	c	a

# Prisoner's dilemma and the environment

- Two countries.
- A fossil-fuel economy is worth 20 units.
- A switch to renewables reduces economic benefit to 10 units.
- But cost of CO<sub>2</sub> emissions paid by both countries via climate change.
- Cost is 6 units per polluting country.

# Prisoner's dilemma and the environment

- Two countries.
- A fossil-fuel economy is worth 20 units.
- A switch to renewables reduces economic benefit to 10 units.
- But cost of CO<sub>2</sub> emissions paid by both countries via climate change.
- Cost is 6 units per polluting country.

	2	
	COOPERATE	DEFECT
1	10	14
	COOPERATE	DEFECT
10	4	8
	14	8

# Prisoner's dilemma and the environment

- Two countries.
- A fossil-fuel economy is worth 20 units.
- A switch to renewables reduces economic benefit to 10 units.
- But cost of CO<sub>2</sub> emissions paid by both countries via climate change.
- Cost is 6 units per polluting country.

	2	
	COOPERATE	DEFECT
1	10	14
	COOPERATE	
	10	4
	DEFECT	
	4	8
		14

It is in the interests of each country, *regardless of what the other is doing*, to keep on polluting.

# The Tragedy of the Commons

Hardin (1968, *Science*)

- William Forster Lloyd's *Tragedy of the Commons*
  - Herders share a common, limited-size, pasture
  - Utility to an individual herder of adding one more sheep is greater than the cost to themselves, a cost shared by all
  - Rational herders will put as many sheep on the commons as possible




Figure 10.2  
*Environmental Science*  
© 2012 W. H. Freeman and Company

# The Tragedy of the Commons

Hardin (1968)

- Overpopulation
  - 'They think that farming the seas or developing new strains of wheat will solve the problem – technologically'
  - Adam Smith's 'invisible hand'
    - Market settles on production and prices beneficial to all
    - ...but not invariably true



# The Tragedy of the Commons

Hardin (1968)

- Overpopulation
  - 'They think that farming the seas or developing new strains of wheat will solve the problem – technologically'
  - Adam Smith's 'invisible hand'
    - Market settles on production and prices beneficial to all
    - ...but not invariably true
- How to solve? Appeal to people's conscience?
  - Appealing to conscience will lead to the elimination of conscience
  - Also creates a 'double-bind'
    - you're immoral if you don't cooperate,
    - you're stupid if you do



# The Tragedy of the Commons

Hardin (1968)

- Overpopulation
  - 'They think that farming the seas or developing new strains of wheat will solve the problem – technologically'
  - Adam Smith's 'invisible hand'
    - Market settles on production and prices beneficial to all
    - ...but not invariably true
- How to solve? Appeal to people's conscience?
  - Appealing to conscience will lead to the elimination of conscience
  - Also creates a 'double-bind'
    - you're immoral if you don't cooperate,
    - you're stupid if you do
- Key point is that ToC occurs even if all individuals understand the consequences of their actions



# The Tragedy of the Commons

Hardin (1968)

- Overpopulation
  - 'They think that farming the seas or developing new strains of wheat will solve the problem – technologically'
  - Adam Smith's 'invisible hand'
    - Market settles on production and prices beneficial to all
    - ...but not invariably true
- How to solve? Appeal to people's conscience?
  - Appealing to conscience will lead to the elimination of conscience
  - Also creates a 'double-bind'
    - you're immoral if you don't cooperate,
    - you're stupid if you do
- Key point is that ToC occurs even if all individuals understand the consequences of their actions
- Next lab - fishing game



... but wait

- Do you behave like this in daily life?

... but wait

- Do you behave like this in daily life?
- Consider the cleaner fish - why doesn't the big fish just eat it when it's done?



... but wait

- Do you behave like this in daily life?
- Consider the cleaner fish - why doesn't the big fish just eat it when it's done?
- Consider the vampire bat
  - Feeds by biting a small hole in some mammal and lapping up blood
  - Two nights without food will die
  - Roost together in colonies in caves
  - Bats who had a feed that night will regurgitate food for those who had none



... but wait

## A BAT'S DILEMMA

Game theory can model the choice to share a meal with a hungry neighbor.

		Bat A shares	Bat A doesn't share
			
Bat B shares	Bat A shares	Both survive, if a little hungrier. Bat A fitness: 0.9   Bat B: 0.9	Bat A stays full; Bat B dies. Bat A fitness: 1   Bat B: 0
	Bat A doesn't share		
		Bat B stays full; Bat A dies. Bat A fitness: 0   Bat B: 1	Each survives alone; much hungrier. Bat A fitness: 0.4   Bat B: 0.4



Olena Shmahalo/Quanta Magazine

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

- ① First play is 'cooperate'

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

- ① First play is 'cooperate'
- ② Then repeat:
  - ① If opponent played 'defect', next play is 'defect'

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

- ① First play is 'cooperate'
- ② Then repeat:
  - ① If opponent played 'defect', next play is 'defect'
  - ② If opponent played 'cooperate', next play is 'cooperate'

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

- ① First play is 'cooperate' - be nice
- ② Then repeat:
  - ① If opponent played 'defect', next play is 'defect'
  - ② If opponent played 'cooperate', next play is 'cooperate'

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

- ① First play is 'cooperate' - be nice
- ② Then repeat:
  - ① If opponent played 'defect', next play is 'defect' - be provokable
  - ② If opponent played 'cooperate', next play is 'cooperate'

# The Axelrod Tournaments

- 1980s Robert Axelrod held a series of tournaments
  - Scientists could submit their code to play PD
  - Each algorithm would be played against each other for multiple rounds of PD
- Notice: *repeated* plays against same opponent
  - So far looked at *one-shot* PD
  - This is the *iterated* PD
- Many clever algorithms submitted



R. Axelrod (left)

Winner - Anatol Rapoport - with a very simple program tit-for-tat:

- ① First play is 'cooperate' - be nice
- ② Then repeat:
  - ① If opponent played 'defect', next play is 'defect' - be provokable
  - ② If opponent played 'cooperate', next play is 'cooperate' - be forgiving

# The usefulness of evolutionary game theory

- Small communities actually don't just overgraze and ruin their commons
- People and other animals do find ways to cooperate in PD-like situations



# The usefulness of evolutionary game theory

- Small communities actually don't just overgraze and ruin their commons
- People and other animals do find ways to cooperate in PD-like situations
- Conditions for cooperation in PD incl.:
  - ① May play the same player again
  - ② Ability to recognise other player
    - Repeated game like Axelrod's Tournament
    - Fixed meeting place, or individual recognition
  - ③ Unknown number of future games



# The usefulness of evolutionary game theory

- Small communities actually don't just overgraze and ruin their commons
- People and other animals do find ways to cooperate in PD-like situations
- Conditions for cooperation in PD incl.:
  - ① May play the same player again
  - ② Ability to recognise other player
    - Repeated game like Axelrod's Tournament
    - Fixed meeting place, or individual recognition
  - ③ Unknown number of future games
- But for large-scale problems, like climate change, conditions above are not met



# The usefulness of evolutionary game theory

- Small communities actually don't just overgraze and ruin their commons
- People and other animals do find ways to cooperate in PD-like situations
- Conditions for cooperation in PD incl.:
  - ① May play the same player again
  - ② Ability to recognise other player
    - Repeated game like Axelrod's Tournament
    - Fixed meeting place, or individual recognition
  - ③ Unknown number of future games
- But for large-scale problems, like climate change, conditions above are not met
- Hardin - a necessity that we recognise the problem
  - Privatisation? ('Injustice is preferable to ruin')
  - 'Mutual coercion'



# Summary

- Evolutionary game theory
  - Game theory introduced to biology from economics / political science
  - A way to study Darwinian evolution in a mathematical framework
  - Allows fitness to be a function of others' traits or strategies

# Summary

- Evolutionary game theory
  - Game theory introduced to biology from economics / political science
  - A way to study Darwinian evolution in a mathematical framework
  - Allows fitness to be a function of others' traits or strategies
- Key concepts
  - Nash equilibrium
  - Evolutionarily stable strategy *sensu* Maynard Smith
  - Evolutionary and convergence stability *sensu* adaptive dynamics

# Summary

- Evolutionary game theory
  - Game theory introduced to biology from economics / political science
  - A way to study Darwinian evolution in a mathematical framework
  - Allows fitness to be a function of others' traits or strategies
- Key concepts
  - Nash equilibrium
  - Evolutionarily stable strategy *sensu* Maynard Smith
  - Evolutionary and convergence stability *sensu* adaptive dynamics
- Explanatory power of evolutionary game theory
  - Hawk-Dove – explains limited warfare
  - Prisoner's Dilemma and Tragedy of the Commons
    - Explains why cooperation can be difficult to achieve

# Summary

- Evolutionary game theory
  - Game theory introduced to biology from economics / political science
  - A way to study Darwinian evolution in a mathematical framework
  - Allows fitness to be a function of others' traits or strategies
- Key concepts
  - Nash equilibrium
  - Evolutionarily stable strategy *sensu* Maynard Smith
  - Evolutionary and convergence stability *sensu* adaptive dynamics
- Explanatory power of evolutionary game theory
  - Hawk-Dove – explains limited warfare
  - Prisoner's Dilemma and Tragedy of the Commons
    - Explains why cooperation can be difficult to achieve
- Iterated Prisoner's Dilemma
  - One example of how cooperation can evolve