

# Differential equations as mathematical models

Nadia Lafrenière

ENGR 213 Lecture 6  
September 21, 2023

# Mathematical models

The mathematical description of a social, physical or economic phenomenon is called a **mathematical model**, and is generally made of equations.

It is a simplification of reality, even though it can be very accurate if we take into account the right variables.

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.

Since the model is a **simplification**, we may choose to use only some variables at first.

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.  
Since the model is a **simplification**, we may choose to use only some variables at first.
3. Specify the **level of resolution** of the model

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.  
Since the model is a **simplification**, we may choose to use only some variables at first.
3. Specify the **level of resolution** of the model
4. Make assumptions about the **interactions** of the variables: those can be, for example, physical laws

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.  
Since the model is a **simplification**, we may choose to use only some variables at first.
3. Specify the **level of resolution** of the model
4. Make assumptions about the **interactions** of the variables: those can be, for example, physical laws
5. Express your model through **equations**. For differential equations, your equations should have derivatives.

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.  
Since the model is a **simplification**, we may choose to use only some variables at first.
3. Specify the **level of resolution** of the model
4. Make assumptions about the **interactions** of the variables: those can be, for example, physical laws
5. Express your model through **equations**. For differential equations, your equations should have derivatives.
6. Identify any **initial conditions** (or values at specific times) that you know.



# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.  
Since the model is a **simplification**, we may choose to use only some variables at first.
3. Specify the **level of resolution** of the model
4. Make assumptions about the **interactions** of the variables: those can be, for example, physical laws
5. Express your model through **equations**. For differential equations, your equations should have derivatives.
6. Identify any **initial conditions** (or values at specific times) that you know.
7. **Solve** the differential equation.

# Construction of models

Steps to constructing a mathematical model:

1. Identify the **variables**, and the known information
2. Choose the variables to use in the model.  
Since the model is a **simplification**, we may choose to use only some variables at first.
3. Specify the **level of resolution** of the model
4. Make assumptions about the **interactions** of the variables: those can be, for example, physical laws
5. Express your model through **equations**. For differential equations, your equations should have derivatives.
6. Identify any **initial conditions** (or values at specific times) that you know.
7. **Solve** the differential equation.
8. If needed, **adjust your model** by taking more variables into account. Example: Friction, or air resistance in physical models.

# A few models

- ▶ Growth and decay: population dynamics, radioactive decay

# A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling

# A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling
- ▶ Spread of a disease

# A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling
- ▶ Spread of a disease
- ▶ Mixtures and concentration of a solution

# A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling
- ▶ Spread of a disease
- ▶ Mixtures and concentration of a solution
- ▶ Current in electrical circuits - Kirchoff's laws

# A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling
- ▶ Spread of a disease
- ▶ Mixtures and concentration of a solution
- ▶ Current in electrical circuits - Kirchoff's laws
- ▶ Drainage of a bath tub - Torricelli's law of hydrodynamics



# A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling
- ▶ Spread of a disease
- ▶ Mixtures and concentration of a solution
- ▶ Current in electrical circuits - Kirchoff's laws
- ▶ Drainage of a bath tub - Torricelli's law of hydrodynamics
- ▶ Falling bodies - Newton's second law of motion

## A few models

- ▶ Growth and decay: population dynamics, radioactive decay
- ▶ Cooling and warming - Newton's empirical law of cooling
- ▶ Spread of a disease
- ▶ Mixtures and concentration of a solution
- ▶ Current in electrical circuits - Kirchoff's laws
- ▶ Drainage of a bath tub - Torricelli's law of hydrodynamics
- ▶ Falling bodies - Newton's second law of motion
- ▶ Springs - Newton's second law of motion and Hooke's law

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

## Hypothesis

*The rate at which a population grows is proportional to the size of a population.*

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

## Hypothesis

*The rate at which a population grows is proportional to the size of a population.*

- ▶ Equation:  $\frac{dP}{dt} = cP$ , for some constant  $c$

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

## Hypothesis

*The rate at which a population grows is proportional to the size of a population.*

- ▶ Equation:  $\frac{dP}{dt} = cP$ , for some constant  $c$
- ▶ Initial conditions:  $P(0) = P_0$ .

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

## Hypothesis

*The rate at which a population grows is proportional to the size of a population.*

- ▶ Equation:  $\frac{dP}{dt} = cP$ , for some constant  $c$
- ▶ Initial conditions:  $P(0) = P_0$ .
- ▶ Solving yields  $P = P_0 e^{ct}$ .



# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

## Hypothesis

*The rate at which a population grows is proportional to the size of a population.*

- ▶ Equation:  $\frac{dP}{dt} = cP$ , for some constant  $c$
- ▶ Initial conditions:  $P(0) = P_0$ .
- ▶ Solving yields  $P = P_0 e^{ct}$ .
- ▶ Adjustments: Could we take into account deaths? Emigration and immigration?

# Population dynamics

## Problem

*How large will a population be in some fixed amount of time?*

- ▶ Variable(s): population size  $P$ .

## Hypothesis

*The rate at which a population grows is proportional to the size of a population.*

- ▶ Equation:  $\frac{dP}{dt} = cP$ , for some constant  $c$
- ▶ Initial conditions:  $P(0) = P_0$ .
- ▶ Solving yields  $P = P_0 e^{ct}$ .
- ▶ Adjustments: Could we take into account deaths? Emigration and immigration?
- ▶ Remark: the simplified model with only births is useful to model populations that reproduce fast over a short period of time, such as bacteria or invasive species

# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

- ▶ Variable(s): population size  $P$

# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

- ▶ Variable(s): population size  $P$

## Hypothesis

*The rate of birth is proportional to the size of a population, and so is the rate of death.*

# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

- ▶ Variable(s): population size  $P$

## Hypothesis

*The rate of birth is proportional to the size of a population, and so is the rate of death.*

- ▶ Equation:  $\frac{dP}{dt} = bP - d \cdot P$ , where  $b$  represents the birth rate and  $d$  represents the death rate.

# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

- ▶ Variable(s): population size  $P$

## Hypothesis

*The rate of birth is proportional to the size of a population, and so is the rate of death.*

- ▶ Equation:  $\frac{dP}{dt} = bP - d \cdot P$ , where  $b$  represents the birth rate and  $d$  represents the death rate.
- ▶ Initial conditions: initial population  $P_0 = P(0)$

# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

- ▶ Variable(s): population size  $P$

## Hypothesis

*The rate of birth is proportional to the size of a population, and so is the rate of death.*

- ▶ Equation:  $\frac{dP}{dt} = bP - d \cdot P$ , where  $b$  represents the birth rate and  $d$  represents the death rate.
- ▶ Initial conditions: initial population  $P_0 = P(0)$
- ▶ Solving yields  $P = P_0 e^{(b-d)t}$ .



# Birth and death

## Problem

*How large will a population be in some fixed amount of time, assuming there is no migration (such as the population of planet Earth)?*

- ▶ Variable(s): population size  $P$

## Hypothesis

*The rate of birth is proportional to the size of a population, and so is the rate of death.*

- ▶ Equation:  $\frac{dP}{dt} = bP - d \cdot P$ , where  $b$  represents the birth rate and  $d$  represents the death rate.
- ▶ Initial conditions: initial population  $P_0 = P(0)$
- ▶ Solving yields  $P = P_0 e^{(b-d)t}$ .
- ▶ Adjustments: We could still take migration into account.

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

- ▶ Variable(s): Temperature of the object ( $T$ ), temperature of the room  $T_m$ , initial temperature of the object ( $T(0) = T_0$ ).

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

- ▶ Variable(s): Temperature of the object ( $T$ ), temperature of the room  $T_m$ , initial temperature of the object ( $T(0) = T_0$ ).

## Hypothesis (Newton's empirical law of cooling)

*The rate at which the temperature of a body changes is proportional to the difference between the temperature of the body and the temperature of the surrounding medium.*

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

- ▶ Variable(s): Temperature of the object ( $T$ ), temperature of the room  $T_m$ , initial temperature of the object ( $T(0) = T_0$ ).

## Hypothesis (Newton's empirical law of cooling)

*The rate at which the temperature of a body changes is proportional to the difference between the temperature of the body and the temperature of the surrounding medium.*

- ▶ Equation:  $\frac{dT}{dt} = k(T - T_m)$ , for some constant  $k$  (depending on what the object is made of). We can reason that  $k < 0$ .

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

- ▶ Variable(s): Temperature of the object ( $T$ ), temperature of the room  $T_m$ , initial temperature of the object ( $T(0) = T_0$ ).

## Hypothesis (Newton's empirical law of cooling)

*The rate at which the temperature of a body changes is proportional to the difference between the temperature of the body and the temperature of the surrounding medium.*

- ▶ Equation:  $\frac{dT}{dt} = k(T - T_m)$ , for some constant  $k$  (depending on what the object is made of). We can reason that  $k < 0$ .
- ▶ Initial conditions:  $T(0) = T_0$ .

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

- ▶ Variable(s): Temperature of the object ( $T$ ), temperature of the room  $T_m$ , initial temperature of the object ( $T(0) = T_0$ ).

## Hypothesis (Newton's empirical law of cooling)

*The rate at which the temperature of a body changes is proportional to the difference between the temperature of the body and the temperature of the surrounding medium.*

- ▶ Equation:  $\frac{dT}{dt} = k(T - T_m)$ , for some constant  $k$  (depending on what the object is made of). We can reason that  $k < 0$ .
- ▶ Initial conditions:  $T(0) = T_0$ .
- ▶ This is a separable equation, so it is solvable.

# Cooling and warming

## Problem

*A hot object is taken to a cold room at time 0. At what temperature is it at time  $t$ ?*

- ▶ Variable(s): Temperature of the object ( $T$ ), temperature of the room  $T_m$ , initial temperature of the object ( $T(0) = T_0$ ).

## Hypothesis (Newton's empirical law of cooling)

*The rate at which the temperature of a body changes is proportional to the difference between the temperature of the body and the temperature of the surrounding medium.*

- ▶ Equation:  $\frac{dT}{dt} = k(T - T_m)$ , for some constant  $k$  (depending on what the object is made of). We can reason that  $k < 0$ .
- ▶ Initial conditions:  $T(0) = T_0$ .
- ▶ This is a separable equation, so it is solvable.
- ▶ Adjustments: This assumes that the object is small enough compared to the medium as not to alter the temperature of the medium.



## Example problem (on the board)

When a cake is removed from an oven, its temperature is measured at  $300^{\circ}\text{F}$ . Three minutes later its temperature is  $200^{\circ}\text{F}$ . How long will it take for the cake to cool off to  $100^{\circ}\text{F}$  in a room at a temperature of  $70^{\circ}\text{F}$ ?

# Spread of a disease

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

# Spread of a disease

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Variable(s):  $x$  number of infected people,  $y$  number of non-infected people interacting with them,  $R_0$ , the reproduction rate of the virus.

# Spread of a disease

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Variable(s):  $x$  number of infected people,  $y$  number of non-infected people interacting with them,  $R_0$ , the reproduction rate of the virus.

## Hypothesis

*Once people are infected, they have immunity. The reproduction rate of the virus is constant.*

# Spread of a disease

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Variable(s):  $x$  number of infected people,  $y$  number of non-infected people interacting with them,  $R_0$ , the reproduction rate of the virus.

## Hypothesis

*Once people are infected, they have immunity. The reproduction rate of the virus is constant.*

- ▶ Equation:  $\frac{dx}{dt} = R_0xy$ .

# Spread of a disease

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Variable(s):  $x$  number of infected people,  $y$  number of non-infected people interacting with them,  $R_0$ , the reproduction rate of the virus.

## Hypothesis

*Once people are infected, they have immunity. The reproduction rate of the virus is constant.*

- ▶ Equation:  $\frac{dx}{dt} = R_0xy$ .  
Issue:  $x$  and  $y$  both vary over time. A hypothesis one can make is that, in a small town where none of the  $n$  people who live there is infected comes an infected person. Then  $x + y = n + 1$ , so  $y = n + 1 - x$ , and we reduce to a single variable.

# Spread of a disease (continued)

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

# Spread of a disease (continued)

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Initial conditions: Often, we assume that  $x(0) = 1$ , where time 0 means the apparition of the virus.



# Spread of a disease (continued)

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Initial conditions: Often, we assume that  $x(0) = 1$ , where time 0 means the apparition of the virus.
- ▶ Equation:  $\frac{dx}{dt} = R_0xy$ .  
We suggest the substitution  $y = n + 1 - x$  for a small community.

# Spread of a disease (continued)

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Initial conditions: Often, we assume that  $x(0) = 1$ , where time 0 means the apparition of the virus.
- ▶ Equation:  $\frac{dx}{dt} = R_0xy$ .  
We suggest the substitution  $y = n + 1 - x$  for a small community.
- ▶ The new equation  $\frac{dx}{dt} = R_0x(n + 1 - x)$  is a Bernoulli equation:  $(\frac{dx}{dt} - R_0(n + 1)x = -R_0x^2)$ , so we can solve it through substitution.

# Spread of a disease (continued)

## Problem

*A new virus is spreading through infected people coming into contact with non-infected ones. How many people are infected at time  $t$ ?*

- ▶ Initial conditions: Often, we assume that  $x(0) = 1$ , where time 0 means the apparition of the virus.
- ▶ Equation:  $\frac{dx}{dt} = R_0xy$ .  
We suggest the substitution  $y = n + 1 - x$  for a small community.
- ▶ The new equation  $\frac{dx}{dt} = R_0x(n + 1 - x)$  is a Bernoulli equation:  $(\frac{dx}{dt} - R_0(n + 1)x = -R_0x^2)$ , so we can solve it through substitution.
- ▶ Adjustments can be made for taking into account that the virus is evolving, or that people don't interact with each other uniformly in a community.

# Mixtures




FIGURE 1.3.3 Mixing tank

Picture credit: Zill's textbook.

## Problem

*A solution containing some concentration of salt is slowly poured into another one with a different concentration into a leaky tank. What is leaking is the mixed solution. What is the new concentration of the solution?*

# Mixtures




FIGURE 1.3.3 Mixing tank

Picture credit: Zill's textbook.

## Problem

*A solution containing some concentration of salt is slowly poured into another one with a different concentration into a leaky tank. What is leaking is the mixed solution. What is the new concentration of the solution?*

- Variable(s): Concentration of the solution,  $C$ .

# Mixtures




FIGURE 1.3.3 Mixing tank

Picture credit: Zill's textbook.

## Problem

*A solution containing some concentration of salt is slowly poured into another one with a different concentration into a leaky tank. What is leaking is the mixed solution. What is the new concentration of the solution?*

- ▶ Variable(s): Concentration of the solution,  $C$ .
- ▶ Initial conditions: Concentration of the initial solution ( $C_0$ ), concentration of the solution that is poured in ( $C_1$ ).

# Mixtures




FIGURE 1.3.3 Mixing tank

Picture credit: Zill's textbook.

## Problem

*A solution containing some concentration of salt is slowly poured into another one with a different concentration into a leaky tank. What is leaking is the mixed solution. What is the new concentration of the solution?*

- ▶ Variable(s): Concentration of the solution,  $C$ .
- ▶ Initial conditions: Concentration of the initial solution ( $C_0$ ), concentration of the solution that is poured in ( $C_1$ ).
- ▶ The amount of salt that is in the tank is modeled by  $\frac{dA}{dt} = A_1 - A_2$ , where  $A_1$  is the amount of salt poured in in one unit of time, and  $A_2$  is the amount of salt poured out in one unit of time. The concentration is the amount over the volume.

# Mixtures




FIGURE 1.3.3 Mixing tank

Picture credit: Zill's textbook.

## Problem

*A solution containing some concentration of salt is slowly poured into another one with a different concentration into a leaky tank. What is leaking is the mixed solution. What is the new concentration of the solution?*


- ▶ Variable(s): Concentration of the solution,  $C$ .
- ▶ Initial conditions: Concentration of the initial solution ( $C_0$ ), concentration of the solution that is poured in ( $C_1$ ).
- ▶ The amount of salt that is in the tank is modeled by  $\frac{dA}{dt} = A_1 - A_2$ , where  $A_1$  is the amount of salt poured in in one unit of time, and  $A_2$  is the amount of salt poured out in one unit of time. The concentration is the amount over the volume.



## Example problem (solved on the board)

A tank contains 200 liters of fluid in which 30 grams of salt is dissolved. Brine containing 1 gram of salt per liter is then pumped into the tank at a rate of 4 L /min; the well-mixed solution is pumped out at the same rate. Find the number  $A(t)$  of grams of salt in the tank at time  $t$ .

# Electrical circuits




(a)  $LRC$ -series circuit

Picture credit: Zill's textbook.

## Problem

*A series circuit contains an inductor, a resistor and a capacitor. What is the charge on the capacitor,  $q(t)$ , at a given time?*

# Electrical circuits



(a) LRC-series circuit

## Problem

*A series circuit contains an inductor, a resistor and a capacitor. What is the charge on the capacitor,  $q(t)$ , at a given time?*

Picture credit: Zill's textbook.

## Hypothesis

*Kirchoff's laws:*


$$i = i_L = i_R = i_C, \quad E(t) = u_L + u_R + u_C.$$

*Also,*

$$u_L = L \frac{di}{dt}, \quad u_R = Ri, \quad u_C = \frac{1}{C}q,$$

*where  $L$  is the inductance,  $R$  the resistance and  $C$  the capacitance. The charge is related to the current through  $\frac{dq}{dt} = i(t)$ .*

# Electrical circuits (continued)




(a) *LRC*-series circuit

## Hypothesis

$$i = i_L = i_R = i_C, \quad E(t) = u_L + u_R + u_C$$
$$u_L = L \frac{di}{dt}, \quad u_R = Ri, \quad u_C = \frac{1}{C}q.$$

Picture credit: Zill's textbook.

## Electrical circuits (continued)



(a) *LRC*-series circuit

### Hypothesis

$$i = i_L = i_R = i_C, \quad E(t) = u_L + u_R + u_C$$
$$u_L = L \frac{di}{dt}, \quad u_R = Ri, \quad u_C = \frac{1}{C}q.$$


Picture credit: Zill's textbook.

► Plugging  $i = \frac{dq}{dt}$  into  $E(t)$ :

$$E(t) = L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{1}{C}q,$$

which is a second-order differential equation.

## Electrical circuits (continued)



(a) *LRC*-series circuit

### Hypothesis

$$i = i_L = i_R = i_C, \quad E(t) = u_L + u_R + u_C$$
$$u_L = L \frac{di}{dt}, \quad u_R = Ri, \quad u_C = \frac{1}{C}q.$$

Picture credit: Zill's textbook.

- ▶ Plugging  $i = \frac{dq}{dt}$  into  $E(t)$ :

$$E(t) = L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{1}{C}q,$$

which is a second-order differential equation.

- ▶ Solving second-order differential equations is hard, but we can easily solve LR-circuits or RC-circuits, since they give rise to the first-order equations

$$E(t) = L \frac{di}{dt} + Ri \quad \text{and} \quad E(t) = R \frac{dq}{dt} + \frac{1}{C}q.$$

# Drainage of a tank




FIGURE 1.3.4 Water draining from a tank

Picture credit: Zill's textbook.

## Problem

*A tank (such as a bath tub) is emptying itself through a sharp hole. If the water is filled up to height  $h$ , how is  $h$  changing over time?*

# Drainage of a tank




FIGURE 1.3.4 Water draining from a tank

Picture credit: Zill's textbook.

## Problem

*A tank (such as a bath tub) is emptying itself through a sharp hole. If the water is filled up to height  $h$ , how is  $h$  changing over time?*

## Hypothesis (Torricelli's law)

*The speed  $v$  of efflux of water through a sharp-edged hole at the bottom of a tank filled to a depth  $h$  is the same as the speed that a body (in this case a drop of water) would acquire in falling freely from a height  $h$ ; that is,  $v = \sqrt{2gh}$ , where  $g$  is the acceleration due to gravity.*



# Drainage of a tank (continued)




FIGURE 1.3.4 Water draining from a tank

Picture credit: Zill's textbook.

## Hypothesis (Torricelli's law)

*The speed of efflux of water is  $v = \sqrt{2gh}$ , where  $g$  is the acceleration due to gravity.*

# Drainage of a tank (continued)




FIGURE 1.3.4 Water draining from a tank

## Hypothesis (Torricelli's law)

*The speed of efflux of water is  $v = \sqrt{2gh}$ , where  $g$  is the acceleration due to gravity.*

Picture credit: Zill's textbook.

- ▶ Variable(s):  $h$  is changing. The area of the hole,  $A_h$ , impacts the volume that goes out. The volume of the tank,  $V$ , helps to measure the height of the water that is left through  $V = hA_w$ , where  $A_w$  is the area of a cross-section of the tank (on the picture here, a cylinder).

# Drainage of a tank (continued)




FIGURE 1.3.4 Water draining from a tank

Picture credit: Zill's textbook.

## Hypothesis (Torricelli's law)

*The speed of efflux of water is  $v = \sqrt{2gh}$ , where  $g$  is the acceleration due to gravity.*

- ▶ Variable(s):  $h$  is changing. The area of the hole,  $A_h$ , impacts the volume that goes out. The volume of the tank,  $V$ , helps to measure the height of the water that is left through  $V = hA_w$ , where  $A_w$  is the area of a cross-section of the tank (on the picture here, a cylinder).
- ▶ Equation: On one hand,  $\frac{dV}{dt} = \frac{dh}{dt}A_w$ , if  $A_w$  is constant (if the tank is a cube or a cylinder). Also,  $\frac{dV}{dt} = -A_h v = -A_h \sqrt{2gh}$ . Solving for  $\frac{dh}{dt}$ , we get  $\frac{dh}{dt} = -\frac{A_h \sqrt{2gh}}{A_w}$ , which is a separable equation (when the volume is proportional to the height.)