

Ecological trait variation across beetle families

	Family	Guild	Feeding type	Forest type
<i>Anthaxia fulgurans</i>	Buprestidae	fresh wood	xylophagous	deciduous
<i>Anthaxia quadripunctata</i>	Buprestidae	fresh wood	xylophagous	coniferous
<i>Chrysobothris affinis</i>	Buprestidae	fresh wood	xylophagous	deciduous
<i>Buprestis haemorrhoidalis</i>	Buprestidae	fresh wood	xylophagous	coniferous
<i>Rhagium bifasciatum</i>	Cerambycidae	old wood	xylophagous	both
<i>Rhagium inquisitor</i>	Cerambycidae	fresh wood	xylophagous	coniferous
<i>Rhagium mordax</i>	Cerambycidae	fresh wood	xylophagous	deciduous
<i>Grammoptera ruficornis</i>	Cerambycidae	old wood	xylophagous	deciduous
<i>Pachytodes cerambyciformis</i>	Cerambycidae	old wood	xylophagous	both
<i>Rutpela maculata</i>	Cerambycidae	old wood	xylophagous	both
<i>Stenurella melanura</i>	Cerambycidae	old wood	xylophagous	both
<i>Corymbia rubra</i>	Cerambycidae	old wood	xylophagous	coniferous
<i>Bolitophagus reticulatus</i>	Tenebrionidae	wood fungi	mycetophagous	deciduous
<i>Ips typographus</i>	Curculionidae	fresh wood	xylophagous	coniferous
<i>Pterostichus aethiops</i>	Carabidae	old wood	zoophagous	both
<i>Uleiota planatus</i>	Silvanidae	old wood	zoophagous	both