


AGENTIC SYSTEMS

**Most agent
orchestration
is a queue in
a costume.**


which works until two of them care about the same thing



THE STAKES

**Nothing errors.
Nothing alerts. It
just never settles.**

The hardest failures in a fleet are the ones where every component reports healthy, because every component is healthy.




WHAT EVERYONE ASSUMES

Coordination is a scheduling problem.

Two things want one resource, so you order them. That is a real problem, it is solved, and it is not the one you have.

THE BREAK

Nothing is waiting for anything.





WHAT ACTUALLY HAPPENS

One nudges every thirty seconds.

The other looks every ten minutes
and makes a large correction —
based on a picture the fast one
has already partly fixed. So it
overshoots, and the fast one
spends its life undoing that.

**There is no
contention
to resolve.**

Both loops can run whenever they like. The conflict is not over access. It is over the same variable, at different rates.



WHAT BIOLOGY DOES

**It separates
the timescales.**

Loops that touch the same thing
are placed at rates far enough
apart — commonly an order of
magnitude — that the fast one
settles before the slow one looks.



WHY THAT WORKS

**It stops being
coordination and
becomes arithmetic.**

You are no longer negotiating
between two controllers. You are
guaranteeing one of them sees a
settled world.



WHERE THIS STOPS WORKING

**Some loops
genuinely need
the same rate.**

Then separation is not available
and you need one controller, not
two. Splitting them was the
mistake, and no amount of timing
fixes it.

the rule has an edge, and this is it



THE TEST

How far apart are their periods?

If two of your loops touch the same value and the answer is “similar”, you have this already. You just have not attributed it yet.

check the periods before you check the logic