

Number Theoretic Functions

Md Nafis Ul Haque Shifat

October 23, 2023

Contents

1	Multiplicative Function	2
2	Dirichlet Convolution	2
3	More on common multiplicative functions	5
3.1	Number-of-Divisors Function	5
3.2	Sum-of-Divisors Function	5
3.3	Möbius Function	5
3.4	Euler's Totient Function	6
4	The Möbius Inversion	7
5	Problems	7

1 Multiplicative Function

Definition 1.1 (Multiplicative Function). A number-theoretic function f is **multiplicative** if $f(1) = 1$ and $f(mn) = f(m)f(n)$, $\forall m, n \in \mathbb{N}$ such that $\gcd(m, n) = 1$. Additionally, f is called **completely multiplicative** if $f(mn) = f(m)f(n)$, $\forall m, n \in \mathbb{N}$.

Examples.

- $1(n) = 1$: The constant function.
- $Id(n) = n$: The identity function.
- $\varepsilon(n) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } n > 1 \end{cases}$: The unit function
- $\tau(n) = \sum_{d|n} 1$: The number of divisors function.
- $\sigma(n) = \sum_{d|n} d$: The sum of divisors function.
- $\phi(n) = \sum_{i=1}^n [\gcd(i, n) = 1]$: Euler's Totient Function (here the third brackets serve as a boolean function, which returns 1 if the condition is true, 0 otherwise.)

Note that $1, Id, \varepsilon$ are completely multiplicative as well, while ϕ, τ, σ aren't.

Theorem 1. If f is a multiplicative function and if $n = \prod_{i=1}^r p_i^{e_i}$, then $f(n) = \prod_{i=1}^r f(p_i^{e_i})$.

Proof. Since $\gcd(p_i^{e_i}, p_j^{e_j}) = 1, \forall i \neq j$, induction on r proves the theorem. □

2 Dirichlet Convolution

Definition 2.1 (Dirichlet Convolution). If f and g are two arithmetic functions, then their **dirichlet convolution**, denoted by $f * g$, is defined as

$$(f * g)(n) = \sum_{d|n} f(d)g(n/d)$$

Alternatively, we can write $(f * g)(n) = \sum_{ab=n} f(a)g(b)$.

Properties of Dirichlet Convolution

- Convolution is Commutative: $f * g = g * f$
- It is Associative: $(f * g) * h = f * (g * h)$

Proof. $((f * g) * h)(n) = \sum_{dc=n} (f * g)(d)h(c) = \sum_{dc=n} \sum_{ab=d} f(a)g(b)h(c) = \sum_{abc=n} f(a)g(b)h(c)$

Similarly, $(f * (g * h))(n) = \sum_{ad=n} f(a)(g * h)(d) = \sum_{ad=n} f(a) \sum_{bc=d} g(b)h(c) = \sum_{abc=n} f(a)g(b)h(c)$ □

- $f * \varepsilon = f$

Proof.

$$(f * \varepsilon)(n) = \sum_{d|n} f(d)\varepsilon(n/d)$$

Now if $d < n$, $n/d > 1$, so $\varepsilon(n/d) = 0$. Therefore

$$(f * \varepsilon)(n) = f(n)\varepsilon(1) = f(n)$$

□

Theorem 2. *if f and g are both multiplicative, so is $f * g$.*

Proof. Let $h = f * g$. Now, if $\gcd(m, n) = 1$,

$$h(mn) = \sum_{d|mn} f(d)g(mn/d)$$

Now since $d|mn$ and $\gcd(m, n) = 1$, $d = ab$ where $a|m$, $b|n$ and $\gcd(a, b) = 1$. Hence,

$$\begin{aligned} h(mn) &= \sum_{a|m, b|n} f(ab)g(mn/ab) \\ &= \sum_{a|m, b|n} f(a)f(b)g(m/a)g(n/b) \\ &= \sum_{a|m} f(a)g(m/a) \sum_{b|n} f(b)g(n/b) \\ &= h(m)h(n) \end{aligned}$$

□

Corollary 1. *if f is a multiplicative function, and $F(n) = \sum_{d|n} f(n)$, F is multiplicative as well.*

Proof. $F(n) = \sum_{d|n} f(n) = \sum_{d|n} f(n)1(\frac{n}{d}) = (f * 1)(n)$.

Since $f, 1$ both are multiplicative, by **theorem 2**, F is also multiplicative. □

Theorem 3. *if $h = f * g$ and h, g are both multiplicative, so is f .*

Proof. Suppose f is not multiplicative. So, there exists a pair of positive integers (m, n) with $\gcd(m, n) = 1$ such that $f(mn) \neq f(m)f(n)$. We take such a pair with smallest mn .

If $mn = 1$ then $f(1) \neq f(1)f(1)$ which implies $f(1) \neq 1$. Now since g is multiplicative, $g(1) = 1$. But $h(1) = (f * g)(1) = f(1)g(1) \neq 1$ since $f(1) \neq 1$. This is a contradiction since h is multiplicative and $h(1) = 1$.

If $mn > 1$ then

$$\begin{aligned}
h(mn) &= \sum_{d|mn} f(d)g(mn/d) \\
&= \sum_{a|m, b|n} f(ab)g(mn/ab) \\
&= \sum_{a|m, b|n, ab < mn} f(a)f(b)g(m/a)g(n/b) + f(mn)g(1) \\
&= \sum_{a|m, b|n} f(a)f(b)g(m/a)g(n/b) - f(m)f(n) + f(mn) \\
&= \sum_{a|m} f(a)g(m/a) \sum_{b|n} f(b)g(n/b) - f(m)f(n) + f(mn) \\
&= h(m)h(n) - f(m)f(n) + f(mn)
\end{aligned}$$

Now if h is multiplicative, $f(m)f(n) = f(mn)$, which implies f is multiplicative as well. \square

Definition 2.2 (Dirichlet Inverse). *If f is an arithmetic function, we call g **dirichlet inverse** of f , denoted by f^{-1} , if $f * g = \varepsilon$.*

Recall that ε is the unit function: $\varepsilon(n) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } n > 1 \end{cases}$

Theorem 4. *If f is an arithmetic function where $f(1) \neq 0$, then f^{-1} exists.*

Proof. Let $g = \begin{cases} \frac{1}{f(1)} & \text{if } n = 1 \\ -\frac{\sum_{d|n, d < n} f(n/d)g(d)}{f(1)} & \text{if } n > 1 \end{cases}$ Clearly, $(f * g)(1) = 1$. Now for $n > 1$,

$$\begin{aligned}
(f * g)(n) &= (g * f)(n) \\
&= \sum_{d|n} g(d)f(n/d) \\
&= g(n)f(1) + \sum_{d|n, d < n} g(d)f(n/d)
\end{aligned}$$

Now,

$$g(n) = -\frac{\sum_{d|n, d < n} g(d)f(n/d)}{f(1)} \implies -g(n)f(1) = \sum_{d|n, d < n} g(d)f(n/d)$$

So, $(f * g)(n) = g(n)f(1) - g(n)f(1) = 0$. Hence, $(f * g) = \varepsilon$, thus $g = f^{-1}$. \square

Corollary 2. *If f is multiplicative, so is f^{-1}*

Proof. Since $\varepsilon = f * f^{-1}$, and both ε and f are multiplicative, by **theorem 3**, f^{-1} is also multiplicative. \square

3 More on common multiplicative functions

3.1 Number-of-Divisors Function

Let $\tau(n)$ denote the number of divisors of n . So, $\tau(n) = \sum_{d|n} 1$. It is easy to see that τ is multiplicative, since $\tau(n) = \sum_{d|n} 1 = \sum_{d|n} 1(d)1(n/d) = (1 * 1)(n)$. Recall $1(n)$ is the constant function, which is completely multiplicative. Hence, by **theorem 2**, τ is also multiplicative.

Now it is easy to derive the formula of this function. What is $\tau(p^x)$ where p is prime? Of course the divisors of p^x are $1, p, p^2, \dots, p^x$, so total $(x + 1)$ divisors. Now for any $n = \prod_{i=1}^r p_i^{e_i}$, we can simply apply **theorem 1** and get $\tau(n) = \prod_{i=1}^r (e_i + 1)$. $\tau(n)$ can be calculated with a simple loop in $O(\sqrt{n})$.

3.2 Sum-of-Divisors Function

The sum of divisors function is denoted by σ . We can write $\sigma(n) = \sum_{d|n} d$. Similar to the previous function, we can write $\sigma(n) = \sum_{d|n} d = \sum_{d|n} Id(d)1(n/d) = (Id * 1)(n)$. Since Id and 1 are both multiplicative, by **theorem 2**, σ is also multiplicative.

Now, note that for a prime p , $\sigma(p^x) = 1 + p + p^2 + \dots + p^x = \frac{p^{x+1}-1}{p-1}$. So, for $n = \prod_{i=1}^r p_i^{e_i}$, $\sigma(n) = \prod_{i=1}^r \frac{p_i^{e_i+1}-1}{p_i-1}$.

3.3 Möbius Function

Definition 3.1. For a positive integer n , the **möbius function**, denoted by μ , is defined as,

$$\mu(n) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } p^2 | n \text{ for some prime } p \\ (-1)^r & \text{if } n = p_1 p_2 \dots p_r, \text{ where } p_i \text{ are distinct primes} \end{cases}$$

The next theorem shows a very important property of möbius function.

Theorem 5. $\sum_{d|n} \mu(d) = \begin{cases} 1 & \text{if } n = 1 \\ 0 & \text{if } n > 1 \end{cases} = \varepsilon(n)$

Proof. For $n = 1$, $\sum_{d|n} \mu(d) = \mu(1) = 1$

When $n > 1$, let $n = \prod_{i=1}^r p_i^{e_i}$. If $d|n$, then $d = \prod_{i=1}^r p_i^{f_i}$, where $f_i \leq e_i$, for $1 \leq i \leq r$. Now if $\exists i$ such that $f_i > 1$, $\mu(d)$ will be 0. So, the only divisors we care about are the ones with $f_i \leq 1, \forall i$. This means we have r distinct primes, and we will go through each possible combinations and add their contribution to the sum accordingly. For example, if d contains k primes, then we have $\binom{r}{k}$

possible values for d , and each of them will contribute $(-1)^k$ to the sum. So,

$$\begin{aligned}
\sum_{d|n} \mu(d) &= \sum_{\substack{(f_1, f_2, \dots, f_r) \\ f_i \in \{0, 1\}}} \mu\left(\prod_{i=1}^r p_i^{f_i}\right) \\
&= (-1)^0 \binom{r}{0} + (-1)^1 \binom{r}{1} + \dots + (-1)^r \binom{r}{r} \\
&= \sum_{i=0}^r \binom{r}{i} (-1)^i \\
&= (1 - 1)^r \\
&= 0
\end{aligned}$$

□

We also conclude that μ is **multiplicative** from this theorem, since,

$$\sum_{d|n} \mu(d) = \sum_{d|n} \mu(d) 1(n/d) = (\mu * 1)(n) = \varepsilon(n)$$

This implies μ is the dirichlet inverse of the constant function. So, by **Corollary 2**, μ is multiplicative. Since μ is the dirichlet inverse of the constant function, using the definition of g in **theorem 4**, we can write that,

$$\mu(n) = \sum_{d|n, d < n} -\mu(d)$$

Now we can use this in sieve to calculate the value of möbius function for integers from 1 to n . Here's an implementation in C++:

```

mu[1] = 1;
for(int i = 1; i < N; i++) {
    for(int j = 2 * i; j < N; j+=i) {
        mu[j] -= mu[i];
    }
}

```

This is basically the standard sieve: instead of iterating over d for each n , for each d , we loop over its multiples and add its contribution to the answer of the multiples. So, it runs in $O(n \log n)$ time.

3.4 Euler's Totient Function

Definition 3.2. For a positive integer n , **Euler's Totient Function**, denotes as $\phi(n)$, is defined as the number of positive integers i up to n such that $\gcd(i, n) = 1$.

For example, $\phi(4) = 2$, since from 1 to 4, only 1 and 3 are coprimes with 4.

Theorem 6. $\sum_{d|n} \phi(d) = n$

Proof. Let S_d be the set of all positive integers up to n whose gcd with n is d . Now, $\gcd(n, m) = d$ implies $\gcd(n/d, m/d) = 1$. By definition, the number of such m is $\phi(n/d)$. So,

$$\sum_{d|n} |S_d| = \sum_{d|n} \phi(n/d)$$

Clearly, the sets S_d for all d are pairwise disjoint. Moreover, for every integer $1 \leq k \leq n$, there exists a d such that $d|n$ and $k \in S_d$. Hence, $\sum_{d|n} |S_d| = \sum_{d|n} \phi(n/d) = \sum_{d|n} \phi(d) = n$. \square

Since, $\sum_{d|n} \phi(d) = \sum_{d|n} \phi(d)1(n/d) = (\phi * 1)(n) = Id(n)$, we also see that ϕ is a multiplicative function.

Note that if p is prime, $\phi(p) = p - 1$, and $\phi(p^k) = p^k - p^{k-1} = p^{k-1}(p - 1)$. So, if $n = \prod_{i=1}^r p_i^{e_i}$, by **theorem 1**, we get

$$\phi(n) = \prod_{i=1}^r p_i^{e_i-1}(p_i - 1) = \prod_{i=1}^r p_i^{e_i} \frac{(p_i - 1)}{p_i} = n \prod_{i=1}^r \left(1 - \frac{1}{p_i}\right)$$

which is the well-known formula for calculating $\phi(n)$.

About implementation, we can simply initialize $\phi(n) = n$, and for each prime divisor p , we update its value by multiplying it with $(1 - \frac{1}{p}) = \frac{p-1}{p}$. Here's the code:

```
for (int i = 1; i <= n; i++) phi[i] = i;
for(int p = 2; p <= n; p++) {
    if (phi[p] == p) {
        phi[p] = p - 1;
        for (int i = 2 * p; i <= n; i += p) {
            phi[i] = (phi[i] / p) * (p - 1);
        }
    }
}
```

4 The Möbius Inversion

Theorem 7. Let f and g be two arithmetic functions. Then, $g(n) = \sum_{d|n} f(d)$ if and only if $f(n) = \sum_{d|n} g(d)\mu(n/d)$.

Proof. Note that $g(n) = \sum_{d|n} f(d) = \sum_{d|n} f(d)1(n/d) = (f * 1)(n)$, which implies $g = f * 1$. Similarly, $f(n) = \sum_{d|n} g(d)\mu(n/d) = (g * \mu)(n)$, which implies $f = g * \mu$. Hence it suffices to show that $g = f * 1$ if and only if $f = g * \mu$.

$$g * \mu = (f * 1) * \mu = f * (1 * \mu) = f * \varepsilon = f$$

. Now for converse,

$$f * 1 = (g * \mu) * 1 = g * (1 * \mu) = g * \varepsilon = g$$

.

\square

I kept this single theorem in a separate section only because it is probably the most useful part of the entire note!

5 Problems

Problem 1. Given n , calculate the number of pairs (i, j) such that $i, j \in [1, n]$ and $\gcd(i, j) = 1$.

Solution. We basically need to find the value of $\sum_{i=1}^n \sum_{j=1}^n [\gcd(i, j) = 1]$. Now note that $[\gcd(i, j) = 1] = \varepsilon(\gcd(i, j))$. Hence,

$$\sum_{i=1}^n \sum_{j=1}^n [\gcd(i, j) = 1] = \sum_{i=1}^n \sum_{j=1}^n \varepsilon(\gcd(i, j)) = \sum_{i=1}^n \sum_{j=1}^n \sum_{d|\gcd(i, j)} \mu(d)$$

Now, $d|\gcd(i, j)$ means $d|i$ and $d|j$. So,

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n \sum_{d|\gcd(i, j)} \mu(d) &= \sum_{i=1}^n \sum_{j=1}^n \sum_{d=1}^n [d|i][d|j] \mu(d) \\ &= \sum_{d=1}^n \mu(d) \sum_{i=1}^n [d|i] \sum_{j=1}^n [d|j] \\ &= \sum_{d=1}^n \mu(d) \left(\left\lfloor \frac{n}{d} \right\rfloor\right)^2 \end{aligned}$$

Note that the number of multiples of d from 1 to n is $\lfloor \frac{n}{d} \rfloor$. We can calculate the value of $\mu(i)$ for integers up to n with sieve, and calculate the final answer with a simple loop.

The sequence $\lfloor \frac{n}{1} \rfloor, \lfloor \frac{n}{2} \rfloor, \dots, \lfloor \frac{n}{n} \rfloor$ contains at most $2\sqrt{n}$ unique values, so if we precalculate the $\mu(i)$, we can calculate the answer for each n in $O(\sqrt{n})$ time.

```
for (int l = 1, r; l <= n; l = r + 1) {
    r = n / (n / l);
    //now for all l <= i <= r, n / i is equal.
}
```

Problem 2. Given n , calculate the number of pairs (i, j) such that $1 \leq i < j \leq n$ and $\gcd(i, j) = 1$.

Solution. We need to calculate $\sum_{i=1}^n \sum_{j=i+1}^n [\gcd(i, j) = 1]$.

$$\begin{aligned} \sum_{i=1}^n \sum_{j=i+1}^n [\gcd(i, j) = 1] &= \sum_{j=2}^n \sum_{i=1}^{j-1} [\gcd(i, j) = 1] \\ &= \sum_{j=2}^n \phi(j) \end{aligned}$$

Now it's a simple loop!

Problem 3. Given a, b, k , calculate the number of pairs (x, y) such that $x \in [1, a], y \in [1, b]$ and $\gcd(x, y) = k$. Also, (x, y) and (y, x) are considered the same.

Solution. First of all, note that $\gcd(x, y) = k$ implies that $\gcd(x/k, y/k) = 1$. Hence, we need to find $\sum_{i=1}^{\lfloor \frac{a}{k} \rfloor} \sum_{j=1}^{\lfloor \frac{b}{k} \rfloor} [\gcd(i, j) = 1]$.

Let $n = \lfloor \frac{a}{k} \rfloor$ and $m = \lfloor \frac{b}{k} \rfloor$. For simplicity, let's assume $n \geq m$ (otherwise we can simply swap n, m). We need to calculate $\sum_{i=1}^n \sum_{j=1}^{\min(m, i)} [\gcd(i, j) = 1]$.

$$\begin{aligned}
\sum_{i=1}^n \sum_{j=1}^{\min(m,i)} [\gcd(i,j) = 1] &= \sum_{i=1}^m \sum_{j=1}^i [\gcd(i,j) = 1] + \sum_{i=m+1}^n \sum_{j=1}^m [\gcd(i,j) = 1] \\
&= \sum_{i=1}^m \phi(i) + \sum_{i=m+1}^n \sum_{j=1}^m \sum_{d|\gcd(i,j)} \mu(d) \\
&= \sum_{i=1}^m \phi(i) + \sum_{i=m+1}^n \sum_{j=1}^m \sum_{d|i, d|j} \mu(d) \\
&= \sum_{i=1}^m \phi(i) + \sum_{i=m+1}^n \sum_{j=1}^m \sum_{d=1}^m [d|i][d|j] \mu(d) \\
&= \sum_{i=1}^m \phi(i) + \sum_{d=1}^m \mu(d) \sum_{i=m+1}^n [d|i] \sum_{j=1}^m [d|j] \\
&= \sum_{i=1}^m \phi(i) + \sum_{d=1}^m \mu(d) (\lfloor \frac{n}{d} \rfloor - \lfloor \frac{m}{d} \rfloor) (\lfloor \frac{m}{d} \rfloor)
\end{aligned}$$

Now you can calculate it with a simple loop in $O(m)$.

Problem 4. Given n , calculate $\sum_{i=1}^n \sum_{j=1}^n \gcd(i,j)$.

Solution. From **theorem 6**, we know that $\sum_{d|n} \phi(d) = n$. So,

$$\begin{aligned}
\sum_{i=1}^n \sum_{j=1}^n \gcd(i,j) &= \sum_{i=1}^n \sum_{j=1}^n \sum_{d|\gcd(i,j)} \phi(d) \\
&= \sum_{i=1}^n \sum_{j=1}^n \sum_{d=1}^n [d|i][d|j] \phi(d) \\
&= \sum_{d=1}^n \phi(d) (\sum_{i=1}^n [d|i]) (\sum_{j=1}^n [d|j]) \\
&= \sum_{d=1}^n \phi(d) \lfloor \frac{n}{d} \rfloor^2
\end{aligned}$$

which can be calculated in $O(\sqrt{n})$ for each n after the sieve (which would take $O(n \log n)$ or $O(n)$ if you use linear sieve).

Problem 5. (Hyperrectangle GCD) Let there be a k -dimensional hyperrectangle with dimensions n_1, n_2, \dots, n_k . Each unit cell of the hyperrectangle is addressed by (p_1, p_2, \dots, p_k) where $1 \leq p_i \leq n_i$ for all $i \leq k$. The value of a cell is the **gcd** of its coordinates. Calculate the sum of value over all cells of the hyperrectangle.

Solution. We are basically asked to calculate the value of:

$$\sum_{x_1=1}^{n_1} \sum_{x_2=1}^{n_2} \cdots \sum_{x_k=1}^{n_k} \gcd(x_1, x_2, \dots, x_k)$$

Let $\text{cnt}[x]$ be the number of such tuples whose gcd is x . Also, let $n = \min_{i=1}^k n_i$. Then our answer will be $\sum_{g=1}^n g \times \text{cnt}[g]$. We can write $g = \sum_{d|g} \phi(d)$. So,

$$\sum_{g=1}^n g \times \text{cnt}[g] = \sum_{g=1}^n \sum_{d|g} \phi(d) \times \text{cnt}[g]$$

Instead of iterating over divisors d of each g , we can iterate over multiples g of each d . So,

$$\begin{aligned} \sum_{g=1}^n g \times \text{cnt}[g] &= \sum_{g=1}^n \sum_{d|g} \phi(d) \times \text{cnt}[g] \\ &= \sum_{d=1}^n \phi(d) \sum_{d|g} \text{cnt}[g] \end{aligned}$$

Now, $\sum_{d|g} \text{cnt}[g]$ is basically the number of tuples whose gcd is divisible by d . It is easy to see that the number of such tuples are $\lfloor \frac{n_1}{d} \rfloor \times \lfloor \frac{n_2}{d} \rfloor \times \dots \times \lfloor \frac{n_k}{d} \rfloor$. Hence the answer is,

$$\sum_{d=1}^n \phi(d) \left(\lfloor \frac{n_1}{d} \rfloor \times \lfloor \frac{n_2}{d} \rfloor \times \dots \times \lfloor \frac{n_k}{d} \rfloor \right)$$

which is calculatable in $O(nk)$.

Problem 6. (Devu and Birthday Celebration) Answer Q ($Q \leq 10^5$) queries: Given n and f ($n, f \leq 10^5$), calculate the number of ways to distribute n sweets among f friends so that if i -th friend got a_i sweets, $a_i \geq 1$ for all i and $\text{gcd}(a_1, a_2, \dots, a_f) = 1$. Note that the ordering matters: $a = [1, 2]$ and $a = [2, 1]$ are considered different.

Solution. First of all, how many ways are there to distribute n sweets among f people so that everyone gets at least one sweet (ignoring the gcd condition)? For that, we can represent a distribution with a string of 'o' and '|', where 'o' means we give a sweet to the current person and '|' means we move to the next person. For example, if $n = 6$ and $f = 3$, then 'oo|ooo|o' denotes $a = \{2, 3, 1\}$. Since the string has n o's, we have $(n - 1)$ places to insert $(f - 1)$ |'s, which can be done in $\binom{n-1}{f-1}$ ways.

Now for a fixed f , let $h(n)$ be the number of ways to distribute n sweets among f people so that everyone gets at least one sweet. From the explanation above, we get $h(n) = \binom{n-1}{f-1}$. Now let $f(n)$ be the number of ways to distribute n sweets among f people so that each gets at least 1 and gcd of all a_i 's is 1 (which is basically the answer of the problem).

For a particular distribution, if $\text{gcd}(a_1, a_2, \dots, a_f) = d$, then $d|n$.

$$\begin{aligned} a_1 + a_2 + \dots + a_f &= n \\ \implies d(a'_1 + a'_2 + \dots + a'_f) &= n \\ \implies a'_1 + a'_2 + \dots + a'_f &= \frac{n}{d} \end{aligned}$$

where $\text{gcd}(a'_1, a'_2, \dots, a'_f) = 1$. Since $a'_1 + a'_2 + \dots + a'_f = \frac{n}{d}$ and $\text{gcd}(a'_1, a'_2, \dots, a'_f) = 1$, by definition, the number of such a' is $f(n/d)$. Hence, we get

$$h(n) = \sum_{d|n} f(n/d) = \sum_{d|n} f(d)$$

Finally, we can apply möbius inversion!

$$f(n) = \sum_{d|n} h(d) \mu(n/d) = \sum_{d|n} \binom{d-1}{f-1} \mu(n/d)$$

We can simply precalculate the $\mu(i)$ and the divisors for integer up to 10^5 to answer the queries efficiently.

Problem 7. (COPRIME3) Given an array a of length n ($1 \leq a[i] \leq 10^6, 1 \leq n \leq 10^5$), calculate the number of triplets $i < j < k$ such that $\gcd(a[i], a[j], a[k]) = 1$.

Solution. We basically need to calculate $\sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=j+1}^n [\gcd(a[i], a[j], a[k]) = 1]$. So,

$$\begin{aligned} \sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=j+1}^n [\gcd(a[i], a[j], a[k]) = 1] &= \sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=j+1}^n \varepsilon(\gcd(a[i], a[j], a[k])) \\ &= \sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=j+1}^n \sum_{d|\gcd(a[i], a[j], a[k])} \mu(d) \\ &= \sum_{d=1}^{MAXV} \mu(d) \sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=j+1}^n [d|\gcd(a[i], a[j], a[k])] \end{aligned}$$

Now, $d|\gcd(a[i], a[j], a[k])$ implies d separately divides each of them.

So, $\sum_{i=1}^n \sum_{j=i+1}^n \sum_{k=j+1}^n [d|\gcd(a[i], a[j], a[k])]$ is basically the number of triplets whose elements are divisible by d . If $cnt[d]$ is the number of elements in the array a divisible by d , then number of such triplets will be $\binom{cnt[d]}{3}$. Calculating $cnt[]$ array is pretty easy: if $freq[x]$ is the number of occurrences of x , then $cnt[d] = \sum_{d|x} freq[x]$. Since we will be iterating over multiples for each d , it will take $O(n \log n)$ time. Hence we get the final answer:

$$\sum_{d=1}^{MAXV} \mu(d) \binom{cnt[d]}{3}$$

Problem 8. (CF Mike and Foam) You are given an array a and q queries. ($|a|, q \leq 2 \times 10^5, a[i] \leq 5 \times 10^5$). You also have a list L , which is initially empty. In each query, you will be given an integer x . If $x \in L$, then you will remove x from L , otherwise add x to L . Then you need to output the number of pairs (i, j) where $i < j$ and $i, j \in L$ such that $\gcd(a_i, a_j) = 1$.

Solution. Exercise. Check my code if needed [here](#).

References

- [Multiplicative Function, Wikipedia](#)
- Beginning Number Theory, Neville Robbins
- [\[Tutorial\] Math note — Möbius inversion](#)