

# Impact of Glacial Lake Outburst Floods on Flood Frequency Statistics of the Simme River

Thierry Zehnder

Tim Renggli

Group 4 · ETH Zürich · Prof. Passalacqua

9 June 2026

## Abstract

Eight Glacial Lake Outburst Floods (GLOFs) from the Plaine Morte glacier (2011–2018) affected the Simme catchment before a supraglacial bypass channel ended the outburst regime in 2019. We test whether these events distort flood frequency statistics using two long-term discharge records at the Simme gauge in Lenk: daily maximum discharge (BAFU/FOEN, 1944–2025) and daily mean discharge (GRDC, 1944–2020). Under daily maxima, GLOFs produced the annual maximum in 4 of 8 GLOF years; under daily means in 6 of 8, because sustained outflows elevate the 24-hour average above brief storm peaks. The 2018 outburst ( $\hat{T} \approx 789$  yr) dominates the statistical signal. Kolmogorov–Smirnov and Rank-Sum tests reject homogeneity between the GLOF-inclusive record and the pre-GLOF reference in both datasets ( $p \leq 0.002$ ). After removing direct GLOF contributions with a  $\pm 5$ -day masking window, both tests fail to reject homogeneity ( $p \geq 0.17$ ), indicating the natural-flood regime remained stationary. A consistent but non-significant upward tendency in the masked peaks suggests possible gradual hydroclimatic change that warrants monitoring as post-bypass data accumulate. Naive inclusion of GLOF peaks overstates the 100-year design flood by approximately 15 %.

# 1 Introduction

Flood frequency analysis (FFA) rests on the assumption of stationarity: that annual maximum discharges are drawn from a single, time-invariant distribution. When a recurring process inflates extreme flows in a subset of years, this assumption is violated and fitted design quantiles become misleading.

The Simme river (canton Bern, Switzerland) provides a compelling case study.



Study area overview. Lac des Faverges sits on the Glacier de la Plaine Morte at  $\sim 2800$  masl. The GLOF route (blue) follows the Ischer torrent to the Simme, reaching the BAFU 2219 gauge at Oberried ( $\sim 1105$  masl) after  $\sim 12$  km. Orange arrow: uncontrolled subsurface drainage path active 2011–2018. Green arrow: supraglacial bypass channel operational since 2019.

The Plaine Morte glacier hosts a supraglacial lake (Lac des Faverges) whose drainage produced eight documented Glacial Lake Outburst Floods (GLOFs) between 2011 and 2018. Since GLOFs are mechanistically distinct from the rainfall- and snowmelt-driven floods that dominate the pre-2011 record, their inclusion in a stationary FFA is conceptually problematic. Following the largest recorded GLOF in 2018, which caused approximately CHF 2.5 million in damage [GEOTEST AG, 2019], a supraglacial bypass channel was constructed in 2019 [GEOPRÆVENT AG, 2022], ending the uncontrolled outburst regime. The period 2011–2018 is therefore a closed historical episode whose statistical imprint on the long-term flood frequency record warrants retrospective investigation.

Both research questions are addressed in parallel using daily maximum instantaneous discharge (*Tagesmaxima*, BAFU 2219, 1944–2025) and daily mean discharge (GRDC 6935331, 1944–2020), making the choice of flow-aggregation statistic a key methodological variable.

## 2 Research Questions

1. Do recurring Glacial Lake Outburst Floods (GLOFs) from the Plaine Morte glacier systematically distort the flood frequency distribution of the Simme river at Lenk, and if so, by

how much do GLOF-inclusive design quantiles exceed those from a homogeneous reference period?

2. Are the annual maximum discharges recorded during the GLOF period (2011–2018) statistically consistent with the pre-GLOF reference distribution once the direct GLOF contribution is removed, or does the post-2010 natural-flood regime show evidence of additional change?

### 3 Hypotheses

The research questions are operationalised as two testable statistical hypotheses.

**Hypothesis 1 — GLOF inflation:** GLOF-inclusive annual maxima from the 2011–2018 period are statistically incompatible with the pre-GLOF reference GEV distribution.

$H_0^{(1)}$ : The 2011–2018 full annual maxima belong to the same population as the pre-GLOF reference period (no distributional shift).

$H_1^{(1)}$ : The 2011–2018 full annual maxima differ from the pre-GLOF reference distribution (the GLOF signal shifts the tail).

**Hypothesis 2 — Regime stationarity:** Once the direct GLOF contribution is removed by masking, the 2011–2018 annual maxima are statistically indistinguishable from the pre-GLOF reference, indicating that the underlying natural-flood regime remained stationary.

$H_0^{(2)}$ : The 2011–2018 GLOF-masked annual maxima belong to the same population as the pre-GLOF reference period.

$H_1^{(2)}$ : The 2011–2018 GLOF-masked annual maxima differ from the pre-GLOF reference distribution (evidence of additional hydroclimatic change beyond GLOFs).

### 4 Data

Daily maximum discharge (*Tagesmaxima*) for the Simme at Oberried/Lenk (BAFU/FOEN station 2219) spans 1974–2025. Combined with *Monatsmaxima* for 1944–1973 from the same station, this yields an extended Annual Maximum Series (AMS) of  $n = 82$  years (1944–2025). The monthly and daily sub-series agree exactly over the overlap period (maximum difference =  $0.000 \text{ m}^3/\text{s}$ ), so the two sub-series are spliced at 1974 without bias. Both records were provided upon request by the Swiss Federal Office for the Environment [Swiss Federal Office for the Environment (BAFU/FOEN), 2026].

The supplementary analysis uses daily mean discharge from GRDC station 6935331 [Global Runoff Data Centre (GRDC), 2026] (Simme at Lenk, 1944–2020,  $n = 77$ ), providing a 67-year pre-GLOF reference.

GLOF event dates were determined from the Lac des Faverges lake-level time series provided by Ogier et al. [2021], who instrumented the lake from 2012 to 2019 at 10-minute resolution (water level, inferred volume, computed discharge). The GLOF date is defined as the timestamp of peak lake outflow: maximum discharge during the outburst phase, identified by requiring (i) lake volume to be decreasing ( $\Delta V < 0$ ) and (ii) computed lake discharge to exceed  $5 \text{ m}^3/\text{s}$ . The 2011 event pre-dates the logger installation and its date is sourced from Gletschersee Lenk [2026]. Uncontrolled outbursts occurred in each year from 2011 to 2018; from 2019 a bypass channel eliminates spontaneous outbursts [GEOTEST AG, 2019]. The eight confirmed event dates are listed in Table 1.

Table 1: Confirmed GLOF event dates, Plaine Morte glacier (2011–2018). Dates are peak lake outflow from the Lac des Faverges time series; 2011 from Gletschersee Lenk [2026].

| Year | GLOF date      |
|------|----------------|
| 2011 | 27 July 2011   |
| 2012 | 16 July 2012   |
| 2013 | 2 August 2013  |
| 2014 | 7 August 2014  |
| 2015 | 1 August 2015  |
| 2016 | 28 August 2016 |
| 2017 | 19 July 2017   |
| 2018 | 27 July 2018   |

## 5 Methods

### 5.1 Series Construction and GLOF Masking

Annual block maxima are extracted from each daily discharge record. Three annual maximum series (AMS) are constructed: (i) the **full** series spanning the complete record, (ii) a **GLOF-masked** series in which values within  $\pm 5$  days of each confirmed GLOF date are set to missing before the block maximum is taken, and (iii) a **reference** series restricted to the pre-GLOF period (1944–2010). The five-day symmetric window captures the typical GLOF recession limb at this gauge. Hypothesis tests are applied exclusively to the GLOF period 2011–2018; post-2018 years are excluded because the bypass channel altered the outburst mechanism.

From these series, the test groups summarised in Table 2 are derived.

Table 2: Annual maximum series and test groups used in the homogeneity tests. GRDC data extend to 2020 only;  $n$  values for GRDC reflect this shorter record.

| Label           | Description                                     | Period               | $n$ (BAFU 2219) | $n$ (GRDC) |
|-----------------|-------------------------------------------------|----------------------|-----------------|------------|
| Reference       | Pre-GLOF annual maxima; basis for GEV fit       | 1944–2010            | 67              | 67         |
| Full AMS        | Complete record, GLOF-inclusive                 | 1944–2025            | 82              | 77         |
| GLOF-masked AMS | Full AMS with $\pm 5$ -day GLOF windows removed | 1944–2025            | 82              | 77         |
| GLOF-inclusive  | Full annual maxima, GLOF period only            | 2011–2018            | 8               | 8          |
| GLOF-masked     | Masked annual maxima, GLOF period only          | 2011–2018            | 8               | 8          |
| Non-GLOF years  | All years outside 2011–2018                     | 1944–2010, 2019–2025 | 74              | 69         |

### 5.2 GEV Flood Frequency Analysis

A Generalised Extreme Value (GEV) distribution is fitted to each AMS by maximum likelihood estimation. The GEV encompasses Gumbel ( $\xi = 0$ ), Fréchet ( $\xi > 0$ ), and Weibull ( $\xi < 0$ ) families, allowing the data to determine the appropriate tail behaviour. Design quantiles for  $T = 2, 5, 10, 20, 50, 100, 200$  yr are derived from the fitted parameters with delta-method confidence intervals [Wilks, 2011]. Gringorten plotting positions are used for empirical return periods, and model fit is compared using the Akaike Information Criterion.

### 5.3 Homogeneity Tests

Two complementary tests compare the 2011–2018 annual maxima against the pre-GLOF reference, applied identically to both datasets.

**KS test:** each 2011–2018 peak is transformed to a uniform variate via the reference GEV CDF (probability integral transform). The Kolmogorov–Smirnov statistic tests departure from

Uniform(0,1) with critical value  $C_\alpha = K_\alpha/(\sqrt{n} + 0.12 + 0.11/\sqrt{n})$ ,  $K_{0.05} = 1.358$  (Module 1, Eq. 3.3; Passalacqua 2026). Tests run separately for GLOF-inclusive ( $n = 8$ ) and GLOF-masked ( $n = 8$ ).

**Rank-Sum test:** the non-parametric Wilcoxon Rank-Sum test (Module 1, §2.4.1; Passalacqua 2026) runs four comparisons: (5a) pre-GLOF reference vs. GLOF-inclusive as a direct test of  $H_0^{(1)}$ , (5b) pre-GLOF reference vs. GLOF-masked as a direct test of  $H_0^{(2)}$ , (5c) GLOF-inclusive vs. all non-GLOF years as additional corroboration of  $H_0^{(1)}$ , and (5d) GLOF-masked vs. all non-GLOF years as additional corroboration of  $H_0^{(2)}$ .

## 6 Results

### 6.1 Descriptive Statistics

Table 3 summarises the BAFU 2219 annual maxima. GLOF years show a mean 70% above non-GLOF years ( $29.35 \text{ m}^3/\text{s}$  vs.  $17.28 \text{ m}^3/\text{s}$ ), driven largely by the 2018 event. The relative gap is smaller in the GRDC daily-mean series (39%), consistent with daily averaging dampening brief storm peaks.

Table 3: Descriptive statistics of the annual maximum series (1944–2025, BAFU 2219 extended AMS combining Monatsmaxima 1944–1973 and Tagesmaxima 1974–2025).

|                                     | Full series  | Non-GLOF years | GLOF years (2011–2018) |
|-------------------------------------|--------------|----------------|------------------------|
| $n$                                 | 82           | 74             | 8                      |
| Mean [ $\text{m}^3/\text{s}$ ]      | 18.46        | 17.28          | 29.35                  |
| Std [ $\text{m}^3/\text{s}$ ]       | 7.51         | 5.55           | 13.45                  |
| Median [ $\text{m}^3/\text{s}$ ]    | 16.41        | 16.09          | 27.13                  |
| Min / Max [ $\text{m}^3/\text{s}$ ] | 9.89 / 56.90 | 9.89 / 34.67   | 15.55 / 56.90          |

### 6.2 GLOF Contribution to Annual Maxima

Table 4 reports the annual maximum, its implied return period under the reference GEV, and whether the  $\pm 5$ -day masking window captures the annual maximum for each GLOF year, for both discharge records.

Table 4: Annual maxima during the GLOF period 2011–2018 under daily maximum discharge (BAFU 2219) and daily mean discharge (GRDC). The implied return period  $\hat{T}$  is computed under the respective pre-GLOF reference GEV (1944–2010 for both datasets). “GLOF = max?” indicates whether the  $\pm 5$ -day masking window contains the annual maximum.

| Year                            | GLOF date | Daily max. (BAFU 2219)                  |                   |               | Daily mean (GRDC)                       |                   |               |
|---------------------------------|-----------|-----------------------------------------|-------------------|---------------|-----------------------------------------|-------------------|---------------|
|                                 |           | $Q_{\text{ann}}$<br>[m <sup>3</sup> /s] | $\hat{T}$<br>[yr] | GLOF<br>=max? | $Q_{\text{ann}}$<br>[m <sup>3</sup> /s] | $\hat{T}$<br>[yr] | GLOF<br>=max? |
| 2011                            | 27 Jul    | 27.09                                   | 17.0              | No            | 15.08                                   | 17.4              | No            |
| 2012                            | 16 Jul    | 19.12                                   | 3.8               | Yes           | 14.56                                   | 13.6              | Yes           |
| 2013                            | 2 Aug     | 20.59                                   | 5.0               | Yes           | 14.99                                   | 16.6              | Yes           |
| 2014                            | 7 Aug     | 40.48                                   | 126.7             | No            | 21.09                                   | 220.0             | Yes           |
| 2015                            | 1 Aug     | 15.55                                   | 1.9               | No            | 10.11                                   | 1.6               | Yes           |
| 2016                            | 28 Aug    | 27.16                                   | 17.2              | Yes           | 13.97                                   | 10.2              | No            |
| 2017                            | 19 Jul    | 27.95                                   | 19.7              | No            | 13.94                                   | 10.0              | Yes           |
| 2018                            | 27 Jul    | 56.90                                   | 788.8             | Yes           | 19.24                                   | 107.1             | Yes           |
| GLOF = ann. max (of 8 GLOF yrs) |           | 4 / 8                                   |                   |               | 6 / 8                                   |                   |               |

Under daily maxima, the GLOF produced the annual maximum in 4 of 8 years (2012, 2013, 2016, 2018); in the remaining four, storm floods dominated, including a  $\hat{T} \approx 127$ -year event in 2014. Under daily means, GLOFs dominated in 6 of 8 years, because their sustained outflows elevate the 24-hour average above brief convective peaks.

### 6.3 GEV Parameter Estimates

Table 5 shows the MLE parameter estimates. Including GLOF years shifts  $\hat{\xi}$  by +0.091 relative to the reference for BAFU 2219 and by +0.069 for GRDC; masking reduces this residual shift to +0.043 and  $-0.008$  respectively, recovering near-reference tail behaviour. The AIC is substantially lower for the masked than the full-series fit.

Table 5: GEV MLE parameter estimates and AIC for both discharge datasets.

| Dataset                                            | Series                | $n$ | $\hat{\mu}$ | $\hat{\sigma}$ | $\hat{\xi}$ | AIC    |
|----------------------------------------------------|-----------------------|-----|-------------|----------------|-------------|--------|
| <i>BAFU 2219 — daily maximum [m<sup>3</sup>/s]</i> |                       |     |             |                |             |        |
|                                                    | Full 1944–2025        | 82  | 14.94       | 3.95           | 0.251       | 513.66 |
|                                                    | GLOF-masked 1944–2025 | 82  | 14.76       | 3.67           | 0.203       | 497.56 |
|                                                    | Reference 1944–2010   | 67  | 14.52       | 3.55           | 0.160       | 399.73 |
| <i>GRDC — daily mean [m<sup>3</sup>/s]</i>         |                       |     |             |                |             |        |
|                                                    | Full 1944–2020        | 77  | 10.211      | 1.711          | 0.170       | 346.93 |
|                                                    | GLOF-masked 1944–2020 | 77  | 10.133      | 1.580          | 0.093       | 328.16 |
|                                                    | Reference 1944–2010   | 67  | 10.038      | 1.542          | 0.101       | 283.54 |

### 6.4 Design Quantiles

Design quantiles are compared in Tables 6 and 7. The full AMS 100-year quantile exceeds the GLOF-masked estimate by 15.0% for BAFU 2219 and 15.3% for GRDC, growing to 18–20% at 200 years. Masked and reference fits are closely aligned in both datasets.

Table 6: GEV design quantiles [ $\text{m}^3/\text{s}$ ] for selected return periods — BAFU 2219 extended AMS (1944–2025), with 95 % delta-method confidence intervals [Wilks, 2011].

| $T$ [yr]              | 2     | 5     | 10    | 20    | 50    | 100   | 200   |
|-----------------------|-------|-------|-------|-------|-------|-------|-------|
| Full 1944–2025        | 16.47 | 22.12 | 26.81 | 32.38 | 41.08 | 48.99 | 58.68 |
| GLOF-masked 1944–2025 | 16.17 | 21.19 | 25.18 | 29.75 | 36.61 | 42.61 | 49.71 |
| Reference 1944–2010   | 15.87 | 20.53 | 24.09 | 28.04 | 33.77 | 38.61 | 44.17 |

Table 7: GEV design quantiles [ $\text{m}^3/\text{s}$ ] for selected return periods — GRDC daily mean discharge.

| $T$ [yr]              | 2     | 5     | 10    | 20    | 50    | 100   | 200   |
|-----------------------|-------|-------|-------|-------|-------|-------|-------|
| Full 1944–2020        | 10.86 | 13.13 | 14.90 | 16.82 | 19.68 | 22.14 | 24.90 |
| GLOF-masked 1944–2020 | 10.72 | 12.68 | 14.09 | 15.54 | 17.57 | 19.21 | 20.95 |
| Reference 1944–2010   | 10.61 | 12.54 | 13.93 | 15.38 | 17.41 | 19.07 | 20.83 |




Figure 1: Return period plot for BAFU 2219 extended AMS (1944–2025): full (blue), GLOF-masked 1944–2025 (green), and reference 1944–2010 (red) GEV fits with 95 % delta-method confidence bands [Wilks, 2011]. Black crosses mark GLOF-year annual maxima from the full series with year labels. The 2018 event at  $\hat{T} \approx 789$  yr dominates the upper tail. The GLOF-masked curve lies slightly above the reference across all return periods.

## 6.5 Homogeneity Tests

### 6.5.1 KS Test

Table 8 reports the KS statistics ( $n = 8$ ,  $C_{0.05} = 0.455$ ). The GLOF-inclusive reject  $H_0^{(1)}$  in both datasets (BAFU 2219:  $D_n = 0.610$ ,  $p = 0.002$ ; GRDC:  $D_n = 0.776$ ,  $p < 0.001$ ). The GLOF-masked fail to reject  $H_0^{(2)}$  in both ( $D_n < 0.37$ ,  $p > 0.19$ ).

Table 8: KS test results: 2011–2018 peaks vs. pre-GLOF reference GEV (1944–2010) for both discharge datasets ( $n = 8$ ;  $C_{0.05} = 0.455$ ).

| Dataset                | Group          | $D_n$ | $C_{0.05}$ | $p$ -value | Decision                   |
|------------------------|----------------|-------|------------|------------|----------------------------|
| BAFU 2219 (daily max.) | GLOF-inclusive | 0.610 | 0.455      | 0.002      | Reject $H_0^{(1)}$ at 5%   |
|                        | GLOF-masked    | 0.360 | 0.455      | 0.195      | Fail to reject $H_0^{(2)}$ |
| GRDC (daily mean)      | GLOF-inclusive | 0.776 | 0.455      | <0.001     | Reject $H_0^{(1)}$ at 5%   |
|                        | GLOF-masked    | 0.318 | 0.455      | 0.321      | Fail to reject $H_0^{(2)}$ |

### 6.5.2 Rank-Sum Test

Table 9 shows the Rank-Sum results. Comparison 5a directly tests  $H_0^{(1)}$  and rejects in both datasets (BAFU 2219:  $p = 0.002$ ; GRDC:  $p < 0.001$ ). Comparison 5b directly tests  $H_0^{(2)}$  and fails to reject in both (BAFU 2219:  $p = 0.172$ ; GRDC:  $p = 0.233$ ). Comparisons 5c and 5d corroborate these findings using the broader non-GLOF reference group.

Table 9: Rank-Sum test results (two-sided,  $\alpha = 0.05$ ) for both discharge datasets. Comparisons 5a and 5b directly test  $H_0^{(1)}$  and  $H_0^{(2)}$  against the pre-GLOF reference; comparisons 5c and 5d use all non-GLOF years as additional corroboration.

| Dataset   |    | Comparison                            | $W$   | $p$ -value | Decision                   |
|-----------|----|---------------------------------------|-------|------------|----------------------------|
| BAFU 2219 | 5a | Pre-GLOF reference vs. GLOF-inclusive | 83.0  | 0.002      | Reject $H_0^{(1)}$         |
|           | 5b | Pre-GLOF reference vs. GLOF-masked    | 188.0 | 0.172      | Fail to reject $H_0^{(2)}$ |
|           | 5c | GLOF-inclusive vs. non-GLOF years     | 499.0 | 0.002      | Reject                     |
|           | 5d | GLOF-masked vs. non-GLOF years        | 382.0 | 0.182      | Fail to reject             |
| GRDC      | 5a | Pre-GLOF reference vs. GLOF-inclusive | 65.0  | <0.001     | Reject $H_0^{(1)}$         |
|           | 5b | Pre-GLOF reference vs. GLOF-masked    | 198.0 | 0.233      | Fail to reject $H_0^{(2)}$ |
|           | 5c | GLOF-inclusive vs. non-GLOF years     | 486.0 | <0.001     | Reject                     |
|           | 5d | GLOF-masked vs. non-GLOF years        | 349.0 | 0.226      | Fail to reject             |






Figure 2: Left: 2011–2018 BAFU 2219 daily-maximum peaks on the reference GEV return-period curve (1944–2010). The 2018 GLOF-inclusive peak (crimson,  $\hat{T} \approx 789$  yr) dominates; after masking it drops to  $\hat{T} \approx 1.5$  yr (blue triangle). Right: probability plot (Module 1, §3.2; Passalacqua 2026). GLOF-inclusive lie far above the diagonal, driven by 2018. GLOF-masked lie systematically above the 1:1 line, indicating peaks slightly higher than the reference predicts.

## 7 Discussion

**Statistical signal driven by the 2018 event.** The 2018 outburst ( $Q = 56.90 \text{ m}^3/\text{s}$ ,  $\hat{T} \approx 789$  yr) is the primary driver of the KS rejection. The causal attribution rests on the masking experiment: after the  $\pm 5$ -day window, 2018’s annual maximum collapses to  $14.28 \text{ m}^3/\text{s}$  ( $\hat{T} \approx 1.5$  yr), confirming the extreme peak was attributable to the outburst, not concurrent meteorological forcing. With  $n = 8$  the formal KS test carries limited inferential weight in isolation; the masking experiment provides the more compelling mechanistic evidence.

**Hypothesis 1 confirmed.** The KS test and Rank-Sum comparison 5a both reject  $H_0^{(1)}$  in both datasets (BAFU 2219:  $p = 0.002$ ; GRDC:  $p < 0.001$ ). The stronger rejection for GRDC reflects GLOFs dominating 6 of 8 years under daily means versus 4 of 8 under daily maxima. The full AMS GEV overstates the 100-year design flood by 15.0 % for BAFU 2219 and 15.3 % for GRDC relative to the GLOF-masked estimate, driven by an upward shift in the shape parameter ( $\hat{\xi} : 0.160 \rightarrow 0.251$  for BAFU 2219) and dominated by the 2018 event.

**Hypothesis 2 supported.** The KS test and Rank-Sum comparison 5b both fail to reject  $H_0^{(2)}$  in both datasets (BAFU 2219: KS  $p = 0.195$ , Rank-Sum  $p = 0.172$ ; GRDC: KS  $p = 0.321$ , Rank-Sum  $p = 0.233$ ). Once GLOF contributions are removed, the natural-flood regime during 2011–2018 is statistically indistinguishable from the pre-GLOF reference. However, Figs. 1 and 2 reveal a consistent upward tendency: the masked GEV lies above the reference at all return periods and all eight GLOF-masked annual maxima plot above the reference diagonal. This could reflect gradual hydroclimatic change — for instance, increasing precipitation intensity under a warmer climate — but with  $n = 8$  the tests cannot distinguish this from sampling variability. A longer post-bypass record is needed to assess this tendency.

**Implications of the bypass intervention.** The bypass channel constructed in 2019 [GEOTEST AG, 2019] has ended the uncontrolled GLOF episode. The GLOF-masked or pre-GLOF reference GEV is now the appropriate design distribution for natural flood hazard at Lenk. Naive inclu-

sion of unmasked peaks overstates the 100-year design flood by 15 % and the 200-year flood by 18–20 %, leading to unnecessarily conservative standards in any post-bypass hazard reassessment.

**Limitations.** The principal limitation is sample size: only  $n = 8$  GLOF-period years constrain the test groups, making results sensitive to a single outlier. Failure to reject  $H_0^{(2)}$  cannot be interpreted as confirmation of stationarity. Future work should extend the analysis as post-bypass data accumulate and apply mixture models that explicitly separate the GLOF and natural-flood populations [Kite, 1977].

## 8 Conclusion

This study examines the influence of eight Plaine Morte GLOFs (2011–2018) on flood frequency statistics at the Simme gauge in Lenk, using daily maximum (BAFU 2219, 1944–2025,  $n = 82$ ) and daily mean discharge (GRDC 6935331, 1944–2020). The key findings are:

1. **GLOF dominance depends critically on the discharge statistic.** Under daily maxima (BAFU 2219), the GLOF produced the annual maximum in 4 of 8 GLOF years; under daily means (GRDC), in 6 of 8, because sustained outflows elevate the daily average above brief storm peaks. The GLOF period also included an approximately 127-year natural storm flood (2014), two approximately 17–20-year events (2011, 2017), and one minor storm year (2015,  $\hat{T}_{pprox} \approx 2$  yr).
2. **One extreme event dominates the signal.** The 2018 outburst ( $Q = 56.90 \text{ m}^3/\text{s}$ ,  $\hat{T} \approx 789$  yr) drives the KS rejection. After masking, its return period collapses to  $\hat{T} \approx 1.5$  yr, confirming the GLOF as the mechanistic cause.
3. **Tests reject homogeneity for GLOF-inclusive.** KS and Rank-Sum comparison 5a (pre-GLOF reference vs. GLOF-inclusive) both reject  $H_0^{(1)}$  in both datasets (BAFU 2219: KS  $p = 0.002$ , Rank-Sum  $p = 0.002$ ; GRDC: KS  $p < 0.001$ , Rank-Sum  $p < 0.001$ ). With only  $n = 8$  test observations, results are sensitive to a single extreme event.
4. **Masking restores homogeneity.** After the  $\pm 5$ -day mask, neither the KS test nor Rank-Sum comparison 5b (pre-GLOF reference vs. GLOF-masked, direct test of  $H_0^{(2)}$ ) rejects in either dataset (BAFU 2219: KS  $p = 0.195$ , Rank-Sum  $p = 0.172$ ; GRDC: KS  $p = 0.321$ , Rank-Sum  $p = 0.233$ ). Comparison 5d (GLOF-masked vs. all non-GLOF years) corroborates (BAFU 2219:  $p = 0.182$ ; GRDC:  $p = 0.226$ ), confirming the natural-flood regime is statistically unchanged across periods.
5. **Design flood inflation is substantial but event-driven.** Unmasked GEV fitting overstates the 100-year design quantile by approximately 15–16 % in both datasets, driven by the upward shift in shape parameter ( $\hat{\xi} : 0.160 \rightarrow 0.251$  for BAFU 2219).

For engineering applications, the GLOF-masked or pre-GLOF reference GEV is recommended as the design distribution for natural flood hazard at Lenk. The 2019 bypass channel makes this recommendation directly applicable to current planning.

## References

- GEOPRÆVENT AG. Plaine morte: Overall monitoring system for glacial lake drainage channel and runoff. <https://www.geopraevent.ch/plaine-morte-overall-monitoring-system-for-glacial-lake-drainage-channel-and-runoff/>, 2022. Second parallel bypass channel constructed June 2022; accessed May 2026.
- GEOTEST AG. Handling of spontaneous glacial lake drainage — Lac des Faverges, Plaine Morte. <https://www.geotest.ch/en/references/geoinformatics-and-3d-modelling/handling-of-spontaneous-glacial-lake-drainage>, 2019. Supraglacial bypass channel completed summer 2019; accessed May 2026.
- Gletschersee Lenk. Gletschersee lenk — outburst flood event archive. <https://www.gletschersee-lenk.ch>, 2026. Accessed: May 2026.
- Global Runoff Data Centre (GRDC). Daily discharge record, station 6935331 — Simme at Oberried/Lenk. <https://www.bafg.de/GRDC>, 2026. Downloaded April 2026; Federal Institute of Hydrology, Koblenz, Germany.
- Geoffrey W. Kite. *Frequency and Risk Analyses in Hydrology*. Water Resources Publications, Fort Collins, Colorado, 1977.
- Christophe Ogier, Mauro A. Werder, Matthias Huss, Isabelle Kull, Dario Hodel, and Daniel Farinotti. Drainage of an ice-dammed lake through a supraglacial stream: hydraulics and thermodynamics. *The Cryosphere*, 15:5133–5150, 2021. doi: 10.5194/tc-15-5133-2021.
- Paola Passalacqua. Stochastic hydrology — lecture notes, modules 1 & 2. ETH Zürich, 2026.
- Swiss Federal Office for the Environment (BAFU/FOEN). Tagesmaxima und monatsmaxima, station 2219 — Simme at Oberried/Lenk, 2026. Data provided upon request, May 2026.
- Daniel S. Wilks. *Statistical Methods in the Atmospheric Sciences*. Academic Press, Oxford, 3 edition, 2011.