

Adverse Drug Reaction (ADR) Reporting Form

Pharmacovigilance Documentation — Individual Case Safety Report

1. Patient Information

Patient Initials	_____
Age	50 years
Sex	Male
Weight	_____

2. Suspected Drug

Drug Name	Ciprofloxacin
Dose	500 mg
Route	Oral
Frequency	Twice daily (BID)
Indication	Bacterial infection
Start Date	_____
Stop Date	Same day as onset (drug withdrawn)

3. Adverse Event Description

Patient developed generalized urticaria (hives) on the same day as starting ciprofloxacin therapy for a bacterial infection. The reaction presented as widespread, itchy, raised wheals consistent with an acute hypersensitivity reaction to the suspected drug.

Onset	Same day after first dose
Seriousness	Non-serious (no hospitalization/anaphylaxis reported)
Outcome	Recovered

4. Action Taken

Ciprofloxacin was discontinued immediately upon onset of the reaction. Symptomatic treatment was administered, and the patient recovered fully.

5. Causality Assessment — Naranjo Adverse Drug Reaction Probability Scale

#	Question	Answer	Score
1	Are there previous conclusive reports on this reaction?	Yes	+1

2	Did the adverse event appear after the suspected drug was given?	Yes	+2
3	Did the reaction improve when the drug was discontinued?	Yes	+1
4	Did the reaction reappear when the drug was re-administered?	Not done	0
5	Are there alternative causes that could explain the reaction?	No	+2
6	Did the reaction reappear with a placebo?	Not done	0
7	Was the drug detected in blood/fluids at toxic levels?	Not done	0
8	Was the reaction more severe with a higher dose, or less with a lower dose?	Not done	0
9	Did the patient have a similar reaction to the same/similar drug before?	Unknown	0
10	Was the event confirmed by objective evidence?	Yes (clinical observation)	+1

Total Naranjo Score: 7 → Category: Probable

6. WHO-UMC Causality Category

Probable — There is a reasonable time relationship between drug intake and reaction onset. The reaction is unlikely to be explained by the patient's underlying disease or other concomitant medications. Clinical response to withdrawal of the drug was reasonable. Rechallenge was not performed, so the reaction cannot be classified as "Certain."

7. Conclusion

Based on the temporal relationship between drug administration and reaction onset, resolution of symptoms following drug withdrawal, and the absence of alternative explanations, **ciprofloxacin is assessed as the probable causative agent** for the generalized urticaria observed in this patient.

8. Reporter Information

Reporter Name	_____
Role/Designation	_____
Date of Report	_____