



Eötvös Loránd University  
Faculty of Informatics  
Department of Numerical Analysis

# Color image analysis and recognition using Zernike moments

TDK thesis

*Supervisor:*

Zsolt Németh

Title title title

*Author:*

Gergely Nagy

Computer Science MSc

1<sup>st</sup> year

Budapest

## **Abstract**

Image moments and moment invariants are widely used for image analysis and pattern recognition. Rotation, scaling and translation invariant Quaternion Zernike Moment Invariants (QZMIs) have been constructed for use on color images [1]. These moment invariants utilize the orthogonal Zernike polynomials.

In this thesis we introduce a method of transforming a digital RGB image inside the unit circle onto a discrete orthogonal points system. Using this points system for the discretization of the QZMIs, we have achieved significant improvements in the image recognition and reconstruction capabilities of the method.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Contributions . . . . .	3
1.2	Structure of the paper . . . . .	3
<b>2</b>	<b>Background</b>	<b>4</b>
2.1	Image moments . . . . .	4
2.2	Zernike moments . . . . .	5
2.3	Quaternion Zernike moments . . . . .	6
2.4	Quaternion Zernike Moment Invariants . . . . .	7
2.5	Applications . . . . .	7
<b>3</b>	<b>Math?</b>	<b>8</b>
<b>4</b>	<b>Implementation?</b>	<b>9</b>
<b>5</b>	<b>Tests/improvements?</b>	<b>10</b>
5.1	Invariance . . . . .	11
5.2	Image reconstruction . . . . .	11

5.3	Image recognition . . . . .	11
5.4	Template matching . . . . .	11
<b>6</b>	<b>Conclusion</b>	<b>12</b>
	<b>Bibliography</b>	<b>13</b>

# Chapter 1

## Introduction

TODO Intro

Something about the topic, why is it relevant, important...

### 1.1 Contributions

What have we achieved, what is new?

### 1.2 Structure of the paper

Structure

## Chapter 2

# Background

This chapter contains a summary and overview of the concepts used in this thesis and previous results this work is based on, such as image moments and their relevance in image analysis, and more specifically Zernike moments and the state-of-the-art of their application for both grayscale and color image analysis. Furthermore, some examples are provided to show the relevance and use cases of such image moments.

### 2.1 Image moments

In general, image moments are certain descriptive values calculated using the pixel intensities of an image. Different moments can be used to extract certain properties from a picture, for example the centroid of a greyscale image can be calculated as

$$\{\bar{x}, \bar{y}\} = \left\{ \frac{M_{10}}{M_{00}}, \frac{M_{01}}{M_{00}} \right\}$$

where  $M_{ij}$  are the regular image moments defined as

$$M_{ij} = \sum_x \sum_y x^i y^j I(x, y)$$

with  $I(x, y)$  being the pixel value at the coordinates  $(x, y)$ .

#### Moment invariants

Using the image moments, moment invariants can be defined, which are invariant to certain transformations, such as rotation, scaling, and translation.

These moment invariants are widely used in applications for pattern matching and image recognition [2, 3, 4]. In particular, moment invariants can be used in medical applications, such as solving the Pathological Brain Detection problem [5].

## 2.2 Zernike moments

Zernike moments are image moments, defined for greyscale images inside the unit circle, using the Zernike polynomials [6].

Zernike polynomials are complex polynomials, which are orthogonal over the unit circle. In polar coordinates:

$$n \in \mathbb{N}, m \in \mathbb{Z}, n \geq |m|, n - |m| \text{ is even}$$

$$V_{n,m}(\rho, \theta) = R_{n,m}(\rho)e^{im\theta}$$

where  $R_{n,m}(\rho)$  are the radial polynomials defined as

$$R_{n,m}(\rho) = \sum_{k=0}^{\frac{n-|m|}{2}} \frac{(-1)^k (n-k)!}{k! \left(\frac{n+|m|}{2} - k\right)! \left(\frac{n-|m|}{2} - k\right)!} \rho^{n-2k}$$

The Zernike moment of order  $n$  and repetition  $m$  of a greyscale, continuous image function  $f(\rho, \theta)$  given in polar coordinates, is defined as

$$Z_{n,m}(f) = \frac{n+1}{\pi} \int_0^1 \int_0^{2\pi} f(\rho, \theta) \overline{V_{n,m}(\rho, \theta)} \rho d\rho d\theta$$

Because the Zernike polynomials are orthogonal, the following reconstruction of the image function is possible, using a finite  $M$  number of Zernike moments:

$$f(\rho, \theta) \approx \sum_{n=0}^M \sum_{m=-n}^n Z_{n,m}(f) V_{n,m}(\rho, \theta)$$

Since digital images are not represented in polar coordinates, a transformation of the image is needed in order to calculate its Zernike moments. The most commonly used transformation is a linear transformation from the image coordinates to a suitable square

inside the unit circle. After this linear transformation, the following discrete approximation is used to calculate the Zernike moments of a digital image  $f(x, y)$

$$Z_{n,m}(f) = \frac{n+1}{\pi(N-1)^2} \sum_{x=1}^N \sum_{y=1}^N f(x, y) \overline{V_{n,m}(\rho_{x,y}, \theta_{x,y})}$$

where  $N$  is the size of the image, and  $(\rho_{x,y}, \theta_{x,y})$  are the polar coordinates corresponding to the  $(x, y)$  image coordinates.

## 2.3 Quaternion Zernike moments

The previously defined Zernike moments can only be used for greyscale images. For color images, quaternion Zernike moments (QZMs) can be used [7].

Quaternions are a generalization of complex numbers, consisting of one real and three imaginary parts. A quaternion  $q$  can be represented in the form  $q = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$ , where  $a, b, c, d \in \mathbb{R}$  and  $\mathbf{i}, \mathbf{j}, \mathbf{k}$  are the imaginary units, defined by the multiplication rules  $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{ijk} = -1$ . The set of quaternions is denoted by  $\mathbb{H}$ .

Let  $f(\rho, \theta)$  be a pure quaternion valued, continuous RGB image function, defined in polar coordinates on the unit circle. Each color component corresponds to one of the imaginary units. Let  $\boldsymbol{\mu} = \frac{\mathbf{i}+\mathbf{j}+\mathbf{k}}{\sqrt{3}}$  be a unit pure quaternion.

Since the multiplication of quaternions is not commutative, right-side and left-side quaternion Zernike moments can also be defined. The right-side QZM of order  $n$  and repetition  $m$  is defined as

$$Z_{n,m}^R(f) = \frac{n+1}{\pi} \int_0^1 \int_0^{2\pi} R_{n,m}(\rho) f(\rho, \theta) e^{-\boldsymbol{\mu} m \theta} \rho \, d\rho \, d\theta,$$

$$n \geq |m| \text{ and } n - |m| \text{ is even}$$

The left-side QZMs are defined as

$$Z_{n,m}^L(f) = \frac{n+1}{\pi} \int_0^1 \int_0^{2\pi} R_{n,m}(\rho) e^{-\boldsymbol{\mu} m \theta} f(\rho, \theta) \rho \, d\rho \, d\theta$$

Similarly to non-quaternion Zernike moments, the original image can be approximated



by using only a finite  $M$  number of QZMs.

$$f(\rho, \theta) \approx \sum_{n=0}^M \sum_{m=-n}^n Z_{n,m}^R(f) R_{n,m}(\rho) e^{\mu m \theta}$$

$$f(\rho, \theta) \approx \sum_{n=0}^M \sum_{m=-n}^n e^{\mu m \theta} Z_{n,m}^L(f) R_{n,m}(\rho)$$

## Discretization

## 2.4 Quaternion Zernike Moment Invariants

## 2.5 Applications

TODO Background

Image moments in general

Explain Zernike moments

QZMI, QZMRI: greyscale, color, etc

examples, use cases

## Chapter 3

### Math?

## Chapter 4

# Implementation?

This chapter presents the tools and methods used for implementing a program for calculating the QZMIs of an image using both methods of discretization.

Methods for calculating roots of Legendre poly

Quadratures

Program

## Chapter 5

# Tests/improvements?

In this chapter we present the different tests that were performed to compare the capabilities of the old and new methods.

In total, four kinds of tests were conducted:

- Invariance
- Image reconstruction
- Image recognition
- Template matching

All of these tests were run for both the old and new methods and the results have been compared.

**5.1 Invariance**

**5.2 Image reconstruction**

**5.3 Image recognition**

**5.4 Template matching**

## Chapter 6

# Conclusion

TODO conclusion

Future work, hogyan tovább ... (akar saját fejezetebe is)

# Acknowledgment

TODO EFOP

# Bibliography

- [1] Beijing Chen, Huazhong Shu, Hui Zhang, Gang Chen, Christine Toumoulin, Jean-Louis Dillenseger, and Limin Luo. Quaternion zernike moments and their invariants for color image analysis and object recognition. *Signal Processing*, 92:308–318, 02 2012. doi: 10.1016/j.sigpro.2011.07.018.
- [2] Feng Zhang, Shang-qian Liu, Da-bao Wang, and Wei Guan. Aircraft recognition in infrared image using wavelet moment invariants. *Image and Vision Computing*, 27: 313–318, 03 2009. doi: 10.1016/j.imavis.2008.08.007.
- [3] Yi Hsien Lin and Chin-Hsing Chen. Template matching using the parametric template vector with translation, rotation and scale invariance. *Pattern Recognition*, 41 (7):2413–2421, July 2008. ISSN 0031-3203. doi: 10.1016/j.patcog.2008.01.017.
- [4] Florica Mindru, Tinne Tuytelaars, Luc Van Gool, and Theo Moons. Moment invariants for recognition under changing viewpoint and illumination. *Computer Vision and Image Understanding*, 94(1):3 – 27, 2004. ISSN 1077-3142. doi: <https://doi.org/10.1016/j.cviu.2003.10.011>. URL <http://www.sciencedirect.com/science/article/pii/S1077314203001930>. Special Issue: Colour for Image Indexing and Retrieval.
- [5] Yu-Dong Zhang, Shui-Hua Wang, Xiao-Jun Yang, Zheng-Chao Dong, Ge Liu, Preetha Phillips, and Ti-Fei Yuan. Pathological brain detection in mri scanning by wavelet packet tsallis entropy and fuzzy support vector machine. *Springer-Plus*, 4:716, 2015. ISSN 2193-1801. doi: 10.1186/s40064-015-1523-4. URL <https://europepmc.org/articles/PMC4656268>.
- [6] A. Khotanzad and Y. H. Hong. Invariant image recognition by zernike moments. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 12(5):489–497, 1990.
- [7] B. Chen, H. Shu, H. Zhang, G. Chen, and L. Luo. Color image analysis by quaternion zernike moments. In *2010 20th International Conference on Pattern Recognition*, pages 625–628, 2010.