

M624 HOMEWORK – SPRING 2026

Prof. Andrea R. Nahmod

SET 1 - DUE 02/17/2026 BY 5PM

From Chapter 4 :

Additional Problem 1: Prove the Auxiliary Lemma we stated in Lecture 2. Namely prove the following:

A normed vector space $(X, \|\cdot\|)$ is *complete* if and only if every absolutely convergent series in X converges in X .

Recall; i) A series $\{x_n\}_n$ is said to be absolutely convergent if $\sum_{n=1}^{\infty} \|x_n\| < \infty$.

ii) A series $\sum_{n=1}^{\infty} x_n$ converges in X if there exists an x in X such that $\lim_{N \rightarrow \infty} \sum_{n=1}^N x_n = x$ in X ; that is if $\|\sum_{n=1}^N x_n - x\| \rightarrow 0$ as $N \rightarrow \infty$

Additional Problem 2: For any $1 \leq p < \infty$ consider the space

$$L^p(\mathbb{R}^d) := \{f : \mathbb{R}^d \rightarrow \mathbb{C}, \text{ measurable, } \|f\|_{L^p(\mathbb{R}^d)} := \left(\int_{\mathbb{R}^d} |f(x)|^p dm \right)^{\frac{1}{p}} < \infty\}.$$

Assume that $\|f\|_{L^p(\mathbb{R}^d)}$ is a norm[†] whence $d_p(f, g) := \|f - g\|_{L^p(\mathbb{R}^d)}$ defines a metric and L^p is a metric space. Use the proof of completeness that I gave in class from Folland (Lecture 3) when $p = 2$ to **prove** that $L^p(\mathbb{R}^d)$ is *complete*.

(†) Question: can you guess what you would need to prove the triangle inequality when $p \neq 2$?. Justify your answer.

From Chapter 4 (pp 193-194): 1, 2*, 3, 4 (show completeness only), 5, 6, 7.

* to show that $f - g$ is *orthogonal* to g you need to show that $\langle f - g, g \rangle = 0$.

SET 2 - DUE 02/24/2026 BY 5PM

From Chapter 4 (pp 195-197): 8a)

Pb. I. (This is an undergrad. problem but a very useful property to remember)
Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be a periodic function with period p ; that is $\phi(x + p) = \phi(x)$, $\forall x \in \mathbb{R}$. Assume that ϕ is integrable on any finite interval.

(a) Prove that for any $a, b \in \mathbb{R}$

$$\int_a^b \phi(x) dx = \int_{a+p}^{b+p} \phi(x) dx = \int_{a-p}^{b-p} \phi(x) dx$$

(b) Prove that for any $a \in \mathbb{R}$

$$\int_{-p/2}^{p/2} \phi(x+a) dx = \int_{-p/2}^{p/2} \phi(x) dx = \int_{-p/2+a}^{p/2+a} \phi(x) dx$$

In particular we have that $\int_a^{a+p} \phi(x) dx$ does not depend on a , as we discussed in class.

Pb.II. Consider $f \in L^2([-\pi, \pi])$ and assume that $\sum_{n \in \mathbb{Z}} a_n e^{inx} = f(x)$ a.e. x . Show that on any subinterval $[a, b] \subset [-\pi, \pi]$,

$$\int_a^b f(x) dx = \sum_n \int_a^b a_n e^{inx} dx.$$

In particular if $g(x) = \int_a^x f(y) dy$, the Fourier coefficients and series of $g(x)$ can be obtained from a_n , the Fourier coefficients of f .

Pb. III. For $0 < \alpha < 1$, we say that a function f is C^α -Hölder continuous with exponent α if there exists a constant $c = c_\alpha > 0$ such that $|f(x) - f(y)| \leq c|x - y|^\alpha$ for all x, y . For $k \in \mathbb{N}$, we can also define the space $C^{k, \alpha}$ to be that of functions which are k -th times differentiable and whose k -th derivative is C^α -Hölder continuous (we could relabel C^α as $C^{0, \alpha}$). Consider now f a 2π -periodic $C^{k, \alpha}$ function. If a_n are the Fourier coefficients of f , show that for some $C > 0$ independent of n ,

$$|a_n| \leq \frac{C}{|n|^{k+\alpha}}$$

(Hint Integrate by parts and use periodicity.

Bonus Problem (do but do not turn in): 2*a)b) from Chapter 4, pp 202.