



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Hive Simulation

Stefan Gugler, Elias Huwyler, Fabian Tschopp

December 16, 2013

Outline

Introduction

Description of the Standard Model

Environmental Influences

Advanced model: Environment simulation

Simulation results and analysis

Summary

Introduction




Figure 1: *Diagram of our work.*

Description of the Standard Model

It consists of four differential equations:

- Change of brood number
- Change of hive bee number
- Change of forager bee number
- Change of food

Description of the Standard Model

Let's look into one equation: **Change of brood number**

$$\frac{dB}{dt} = LS(H, f) - \phi B$$

- L is the laying rate of the queen
- S is the survival rate
- H is the amount of hive bees
- f is the amount of food
- ϕ is the adult bee emerging factor

Environmental Influences




Figure 2: *The laying rate of the bee queen plotted over a year.*

Environmental Influences

Comparison between the environment dependent equation and the standard one:

- static equation: $\frac{dB}{dt} = LS(H, f) - \phi B$
- dynamic equation: $\frac{dB}{dt} = L_t S(H, f) - \phi B$

→ small changes in the formulae have significant effects (cf. Discussion).

Advanced model: Environment simulation




Figure 3: *Honey bee social dynamics/influences covered by our model.*

Agents: Assigning jobs




Figure 4: *Assigning jobs to unemployed bees. Scouts and foragers are possible.*

Agents: Scout bees




Figure 5: *Scouting behaviour until a flower patch is found or the maximum distance is reached.*

Scouts' random walk

The path a scout bee walks is recorded in a vector of x and y coordinates:

$$\begin{pmatrix} x_0 & x_1 & \dots & x_n \\ y_0 & y_1 & \dots & y_n \end{pmatrix}$$




Figure 6: *Example of a random walk executed by a scout bee.*

Agents: Forager bees




Figure 7: *Foraging behaviour.*

Path optimization




Figure 8: *Path optimization used to short cut the path to flower patches.*

Agents: Returning to the hive




Figure 9: *Forager bee, returning from foraging.*


Agent based model: recorded sample clips

- Day 158, recorded sample with scouts displayed
- Day 158, recorded sample without scouts displayed
- Two different runs, not the same flower patches are being selected


Simulation results and analysis

- Evolution of the model
- Missing flower season comparison
- Critical points in the fall season

Evolution of the model




Daily simulation




Missing flower season comparison

- Eliminate non critical seasons
- Study effects of missing season
- Observe the hives compensation measures

Spring



Summer




Critical points in the fall season


Death criteria:

- Less than 1000 bees at day 400
- Less than 20 kg of stored food at day 400


Stored food variation before fall



Peak value influence



Delay influence around breaking point



Summary

- Standard Model after *D.S. Khoury*
- Advanced Model: Environment simulation
- Complex model of assigning jobs
- Method: random walk and optimization
- Evolution of the model
- Results

The background of the slide features a honeycomb pattern of yellow hexagons with black outlines. A horizontal band of dark green, semi-transparent material covers the middle section of the slide.

Questions?