

23MAT204

Adithya Nair

August 14, 2024

Contents

I	Mathematics	2
1	Revision Linear Algebra	3
1.1	Ways To Calculate Whether A Matrix Is Positive Definite	3
1.2	Large Eigenvalue Computation	3
1.3	Spectral Decomposition	3
1.4	Singular Value Decomposition	3
1.5	The Geometry of SVD	4
1.6	Polar Decomposition	4
1.7	Principal Component Analysis	4
1.8	Norms Of Vectors And Matrices	4
1.9	Best Low Rank Matrix	6
2	Multi variable Optimization	7
2.1	Contour Curves	7
2.2	Multi variable Calculus	7
2.3	Newton's Method	7
2.3	How solution to a linear system can be found as a solution of an optimization problem. .	8
2.4	Numerical Algorithm Of Gauss-Jacobi Method	8
2.5	Gauss-Siedel Iteration Method	9
3	Unidirectional/Line search	10
4	Euler's Method	11
II	Overview	12
5	Calculating Eigenvalues and Eigenvectors	13
6	Norms Of Vectors	14
6.1	L_n norm	14
6.2	S-norm	14
7	Norms Of Matrices	15
7.1	Spectral Norm	15
7.2	Frobenius Norm	15
7.3	Nuclear Norm	15
8	Optimization	16
8.1	Types Of Optimization Problems	16

Part I

Mathematics

Chapter 1

Revision Linear Algebra

Every matrix satisfies its own characteristic matrix.

What this means is

$$\begin{aligned}\lambda^3 - 2\lambda^2 + \lambda - 4 &= 0 \\ A^3 - 2A^2 + A - 4I &= 0 \\ A^2 - 2A + I - 4A^{-1} &= 0 \\ A^{-1} &= \frac{1}{4}[A^2 - 2A + I] \\ X = A^{-1}b &= \frac{1}{4}[A^2 - 2A + I]b\end{aligned}$$

There are some implications behind this. Here's a problem, generate a random 3x3 matrix, find the ranks of $(A - \lambda_1 I)$ and then $(A - \lambda_1 I)(A - \lambda_2 I)$

What you will see is that the rank reduces upon multiplying roots of the characteristic equation

1.1 Ways To Calculate Whether A Matrix Is Positive Definite

1. The minors are positive then negative alternating.
2. The eigenvalues are positive.

1.2 Large Eigenvalue Computation

Large eigenvalues are computed by using a matrix multiplication method. This method involves:

1.3 Spectral Decomposition

$$S = Q\Lambda Q^T = \lambda_1 q_1 q_1^T + \lambda_2 q_2 q_2^T + \cdots + \lambda_n q_n q_n^T$$

This allows us to represent a matrix by the sum of n rank 1 matrices. This is used in square symmetric matrices.

This is the basis behind Principal Component Analysis.

1.4 Singular Value Decomposition

For rectangular matrices.

$$A = U\Sigma V^T = \sigma_1 u_1 v_1^T + \sigma_2 u_2 v_2^T + \cdots + \sigma_r u_r v_r^T$$

This is mainly used for getting useful properties about the matrix without needing to perform significant operations on this matrix, such as the orthogonality.

Singular values are the eigenvalues of a matrix which are **non-zero**

So a rectangular matrix $m \times n$ would have AA^T which is $m \times m$ and $A^T A$ which is $n \times n$. The number of singular values is whether m or n is lower.

The trace of $A^T A$ equal to the sum of all a_{ij}^2 . The trace is the sum of all eigenvalues of $A^T A$, and For $A_{m \times n}$ that's the sum of the eigenvalues squared.

This is known as the **Frobenius Norm**

This method can also be used to generate orthonormal bases for the 4 Fundamental Subspaces

1.5 The Geometry of SVD

Since we have an orthogonal matrix, diagonal and orthogonal matrix... It's a rotation step followed by a stretching step, finally ended by another rotation step.

1.6 Polar Decomposition

This decomposition is a special case of the Singular Value Decomposition. Where you decompose the elements into the polar form with an orthogonal matrix and a positive semi-definite matrix.

For 2D,

$$\vec{x} = r(\cos(\theta) + \sin(\theta))$$

$$\begin{aligned} A &= U\Sigma V^T \\ &= U(V^T V)\Sigma V^T \\ &= (UV^T)(V\Sigma V^T) \\ &= Q \times S \end{aligned}$$

Here S is the scaling matrix, and Q is the orthogonal matrix.

Although my assumptions might be wrong, this might be a way to generate rotation matrices for higher-dimension spaces. Might be useful in exploring the semantic spaces of LLMs.

1.7 Principal Component Analysis

Why do we use normalization while working with data? We use normalization to ensure that the units are eliminated while working with that given data. **This normalization is done by subtracting by the average and divide by the standard deviation.**

Matrix where the row and columns sums are equal.

1.8 Norms Of Vectors And Matrices

Norms are a way to measure the size of a vector/matrix. The norm of a vector which calculates the magnitude is known as the L^2 norm or the Euclidean norm $\|v\|_2$. This number gives us the length of the vector.

In general, the $\|u\|_p$ or the p-norm of a vector is $[|u_1|^p + |u_2|^p + \dots + |u_n|^p]^{\frac{1}{p}}$.

Now, the L_1 norm would be the sum of the components of the vector. This is the sum of the projections to each corresponding axis.

The L_3 norm would be $[|u_1|^3 + |u_2|^3 + \dots + |u_n|^3]^{\frac{1}{3}}$

The L_∞ norm would be $[|u_1|^\infty + |u_2|^\infty + \dots + |u_n|^\infty]^\frac{1}{\infty}$. This returns the maximum vector component of the vector.

$$x + y \&= 1$$

Let's take the equation, $\|v\|_1 = 1 \setminus [^x - y \&= 1$

$$x + y \&= 1$$

$$-x - y = 1$$

\setminus

[./figures/l1eq1](#)

The S-norm of a vector \vec{x} is $\vec{x}^T S \vec{x}$. When S is a symmetric positive definite matrix, this S-norm is known as the energy of vector \vec{v}

There are three types of Matrix norms:

1. Spectral Norm

2. Frobenius Norm $= \sqrt{\sigma_1^2 + \sigma_2^2 + \dots + \sigma_r^2}$

3. Nuclear Norm

1. Spectral Norm We know that the vector norm for a vector \vec{x} is nothing but $\vec{x}^T \vec{x}$. We take this property.

$$\begin{aligned} \text{Max} \|A\|_2^2 &= \text{Max} \frac{\|Ax\|_2^2}{\|x\|_2^2} \\ &= \text{Max} \frac{x^T A^T A x}{x^T x} \\ &= \text{Max} \{\lambda_i(S)\} = \lambda_1 = \sigma_1^2 \end{aligned}$$

2. Frobenius Norm The Frobenius norm for a matrix M, $\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$ is the equation $\sqrt{a_{11}^2 + a_{12}^2 + a_{21}^2 + a_{22}^2}$

3. Nuclear Norm The nuclear norm is the sum of the singular values of a matrix A.

For an identity matrix,

- The Spectral Norm is 1
- The Frobenius Norm is \sqrt{n}
- The Nuclear Norm is n

For an orthogonal matrix,

- The Spectral Norm is 1
- The Frobenius Norm is \sqrt{n}
- The Nuclear Norm is n

1.9 Best Low Rank Matrix

We say that a matrix is the best approximation of another matrix, based on the Frobenius Norm.. For a singular value decomposition $A = U\Sigma V^T$, if we assume that the singular values are arranged in descending order... We can select the singular value range where the values are significant contributors to the final matrix.

We can then reduce the size of U, Σ and V^T into a smaller matrix B , based on the number of singular values chosen which would give the best approximation.

Let $A = U\Sigma V^T$ where $\Sigma : \sigma_1 \geq \sigma_2 \geq \dots \sigma_n$, then $B = U_{m \times m} \Sigma V^T_{n \times n}$ is a best rank- k approx. to A . Where, S is a diagonal matrix of $n \times n$ where $s_i = \sigma_i (i = 1 \dots k)$ else $s_i = 0$, by best B is a solution to $\min_B \|A - B\|_F$ where $\text{rank}(B) = k$

Chapter 2

Multi variable Optimization

We minimize $f(x_1, x_2, \dots, x_3)$.

2.1 Contour Curves

2.2 Multi variable Calculus

The points where $\nabla f = \vec{0}$ are called the stationary points. The Hessian matrix should be positive definite for a minima and negative definite for a maxima. A point where there is no change, is known as a saddle point.

$$f(x, y) = x^3 + y^3 + 2x^2 + 4y^2 + 6$$

$$\frac{\partial f}{\partial x} = 3x^2 + 4x$$

$$\frac{\partial^2 f}{\partial x} = 6x + 4$$

$$\frac{\partial f}{\partial y} = 3y^2 + 8y$$

$$\frac{\partial f}{\partial y \partial x} = 0$$

$$\frac{\partial^2 f}{\partial y} = 6y + 8$$

$$x = 0, \frac{-4}{3}$$

$$y = 0, \frac{-8}{3}$$

2.3 Newton's Method

Newton's Method is a numerical method to find a minimum of a function.

Iterative formula:

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)}$$

Find the minimum of $f(x) = x^2 + \frac{54}{x}$ using Newton's method.

Newton's method can be practically done for all functions by evaluating the first and second derivatives numerically and then apply the formula

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \text{ to find root of } f(x) = 0$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \text{ to find minimum of } f(x) \text{ (root of } f'(x) = 0) \text{ to find minimum of } f(x) \text{ (root of } f'(x) = 0) \text{ to find minimum}$$

Find an objective function whose solution is specified as $\vec{x} : A\vec{x} = b$

Take for example, $\frac{1}{2}x^T Ax - b^T x + c$, $x \in R^n$, $A = A^T$

2.4 Numerical Algorithm Of Gauss-Jacobi Method

Input A = [a_{ij}], X0 = x⁽⁰⁾, tolerance TOL, maximum number of iterations,

$$Ax = b$$

We separate A, A = D - L - U

$$(D - L - U)x = b$$

$$(Dx = b + (L + U)x$$

$$(x = D^{-1}b + D^{-1}(L + U)x$$

$$x^{(k)} = Tx^{(k-1)} + c$$

The code to implement the method:

```
A = [5,-2,3;-3,9,1;2,-1,-7];
```

```
b = [-1;2;3];
```

```
n = 100; % No. of iterations
```

```
D = diag(diag(A));
```

```
L = -tril(A,-1);
```

```
U = -triu(A,1);
```

```
X = zeros(size(A)(1),n);
```

```
T = inv(D)*(L+U);
```

```
c = inv(D)*b;
```

```
for i = 2:n
```

```
    X(:,i) = T*X(:,i-1) + c;
```

```
    if(X(:,i) == X(:,i-1))
```

```
        i-1
```

```
        break;
```

```
    end
```

```
end
```

2.5 Gauss-Siedel Iteration Method

The iterative method is somewhat similar to <Gauss-Jacobi method>.

The only difference is that the new values are computed by already existing new values.

```
A = [5,-2,3;-3,9,1;2,-1,-7];
b = [-1;2;3];

n = 100; % No. of iterations
D = diag(diag(A));
L = -tril(A,-1);
U = -triu(A,1);

X = zeros(size(A)(1),n);
T = inv(D-L)*U;
c = inv(D-L)*b;

for i = 2:n
    X(:,i) = T*X(:,i-1) + c;
    if(X(:,i) == X(:,i-1))
        display("The iteration it converges at:")
        i-1
        break;
    end
end
end
```

$$(D - L)x^k = Ux^{k-1} + b$$
$$x^k = (D - L)^{-1}\vec{b} + (D - L)^{-1}(U\vec{x})$$

If the matrix is 'diagonally dominant', where the magnitude of the diagonal elements should be greater than the sum of the magnitude of the other elements(absolute value) in the same row.

Chapter 3

Unidirectional/Line search

For a given function,

Find the minima of $f(x) = (x - 1)^2 + (y - 2)^2$ starting from $(0, 0)$ along the direction of x-axis.

How to perform uni-directional search:

- Write the parametric representation of the line search,

$$\vec{s}(t) = \vec{a} + t\vec{b}$$

- Write the function in terms of t , $f(\vec{s}(t))$. this is a single variable function.
- Find the minimum of $f(\vec{s}(t))$, t^*
- Obtain the solution for unidirectional search by substituting t^* in $\vec{s}(t)$

Find the minimum of the function $f(x) = (x - 2)^2 - y$, starting from the point $(-1, 0)$ along $(1, 1)^T$

Chapter 4

Euler's Method

Part II

Overview

Chapter 5

Calculating Eigenvalues and Eigenvectors

Chapter 6

Norms Of Vectors

6.1 L_n norm

The L_n norm of a vector $x_{m \times 1}$ is,

$$L_n = \sqrt[n]{x_1^n + x_2^n + \cdots + x_m^n}$$
$$L_\infty = \max |x_1|, |x_2|, \cdots, |x_n|$$

6.2 S-norm

Where S is a symmetric positive matrix

$$\|v\|_s = v^T S v$$

Chapter 7

Norms Of Matrices

7.1 Spectral Norm

It is the maximum of the ratio,

$$\frac{\|Ax\|}{\|x\|}$$

The maximum is σ_1 at the vector $x = v_1$

7.2 Frobenius Norm

$$\|A\|_F = \sqrt{\sum_i \sum_j a_{ij}^2}$$

It's the same as the vector norm of the vector of singular values of A

7.3 Nuclear Norm

$$\|A\|_N = \sigma_1 + \sigma_2 + \cdots + \sigma_r$$

Chapter 8

Optimization

8.1 Types Of Optimization Problems

Single variable and multi variable

1. Constrained and unconstrained
2. Linear and nonlinear
3. Continuous and discrete
4. oadsfjdsoifjdsoifj
5. Single-objective and multi-objective
6. Stochastic and deterministic