

Development of a BLM National Aquatic Monitoring Program

Scott Miller, Dan Damman, Melissa Dickard, Mark Gonzalez, Justin Jimenez, Ed Rimbold, Steve Smith, Karl Stein, Jason Taylor, Gordon Toevs and Matt Varner



Background

BLM's mission:

To sustain the health, diversity, and productivity of the public lands for the use and enjoyment of present and future generations



Background

Management objectives: Fisheries, Riparian & SWA




Background

BLM

AIM - Assessment, Inventory, and Monitoring

Management objectives: Fisheries, Riparian & SWA



Background

Management objectives: Fisheries, Riparian & SWA



Background


Common goal of Fisheries, Riparian and SWA Programs:

Protect, restore and maintain the chemical, physical and biological integrity of freshwater resources for present and future generations




Background

- BLM oversees > 250,000 km of perennial streams and rivers (12 Western states including AK)
 - Linear stream km = 5 x Earth's equatorial circumference



Background

Management objectives driven by policy and achieved through the planning process:



Background

- Monitoring (status, trend and effectiveness) is critical because it provides us with information regarding:
 - Resource extent (inventory)
 - Condition and trend
 - Compliance
 - Stressor identification
 - Treatment/BMP effectiveness



Background

- Example tools to ensure compliance with FLPMA and CWA mandates:
 - PFC
 - MIM
 - Macroinvertebrates
 - Water quality monitoring
- But...
 - PFC: qualitative and not a monitoring tool
 - MIM: not inclusive of all management elements (chemical, physical and biological)
 - Macroinvertebrates and water quality: disparate sampling methodologies and interpretation challenges




Background

- Problem:
 - BLM has no interdisciplinary monitoring strategy to determine what, how, when or where different aquatic or riparian indicators are measured and used to make status and trend determinations
- Consequences:
 - Data collection methodologies not standardized
 - Data cannot be integrated within or among agencies
 - Limited scope of inference – largely reach based
 - Frequently not making data-driven mngt. decisions



Background

- Consequences:
 - Are we compliant with federal and state regulations...?
 - What is the effectiveness of BMP...?
 - Collaboration and partnerships...



BLM – AIM Strategy

- Motivation for change:
 - Non-compliance with federal regulations
 - Increasing litigation
 - Landscape-scale management challenges and the advent of various landscape-scale initiatives
 - OMB report regarding BLM's terrestrial monitoring
- BLM's Assessment, Inventory and Monitoring Strategy coming to fruition!




BLM – AIM Strategy

- Central tenants of the AIM Strategy:
 - Quantitative core indicators and methodologies
 - Supplemental indicators
 - Statistically valid sampling frameworks
 - Integration of remote sensing technologies
 - Data acquisition and management plans
- Fundamental objectives:
 - Document the distribution and abundance of natural resources on public lands
 - Determine resource conditions
 - Identify natural resource trend or change



BLM – AIM Strategy

- Terrestrial core indicators, sampling methodologies and survey design established and actively implemented in numerous demonstration projects




BLM - National Aquatic Monitoring Program

- Aquatic corollary to AIM:
 - Oct. 2011: BLM working group established
 - External science advisory team (USGS, EPA, NPS, USFS, academics)
- Objectives:
 - Quantitative aquatic core indicators and methodologies
 - Statistically valid sampling frameworks
 - Integration of remote sensing technologies...?
 - Data acquisition and management plans
- Will be developed through application of a six step process



Road map for developing BLM's NAMP



NAMP - Step 1: Scope, objectives and questions

- Objective(s): to provide the BLM and its partners with quantitative, defensible information needed to understand the condition and trend of lotic ecosystems
- Scope/extent: multiple scales (e.g., stream segment, allotment, field office, state, national), but greatest extent is national level inference
- Target population: all perennial, wadeable streams and associated riparian areas on BLM lands
- Type of monitoring: status, trend and effectiveness




NAMP - Step 1: Scope, objectives and questions


- Monitoring questions:
 - Are key processes (hydrologic, geomorphic, biogeochemical, biological) that sustain functioning riparian and stream systems intact?
 - What stressors are present and what are their relative risks to perennial stream and riparian systems?
 - Has the state or condition of key aquatic indicators shifted in a direction indicating improved or degraded condition through time?



NAMP - Step 2: Conceptual model



NAMP - Step 3: Ecosystem attributes and stressors




NAMP - Step 4: Select aquatic core indicators

- Aquatic core indicators: structural or functional measures that either directly or indirectly quantify the condition of critical ecosystem processes and/or attributes
- Would like to directly measure key processes/attributes, but their inherent complexity = need surrogates
- Developed four phase process for identifying scientifically defensible ACI



NAMP - Step 4: Select aquatic core indicators



NAMP - Step 4: Select aquatic core indicators

- Phase 1 – prioritize ecosystem attributes: survey of state leads and FO personnel
- Phase 2 – ACI selection: survey of indicators used in existing monitoring programs; FO feedback; workshop to prioritize indicators based on selection criteria:
 - Minimal observer bias
 - Documented relationship to attribute(s)/processes
 - Responsive at relevant time-scales
 - Cost
 - Used by multiple agencies
- Phase 3 – Status determinations: workshop with science advisory team and others




NAMP - Step 4: Select aquatic core indicators

- Phase 4 – Field- and GIS-derived covariates: how to account for natural spatial gradients; changing chemical, physical and biological potential of sites
 - ACI working group, science advisory team and FO feedback




Road map for developing BLM's NAMP




NAMP - Step 5: Statistical framework


- Statistically valid survey design:
 - Unbiased: all sites have some known, non-zero probability of being sampled
 - Data integration among spatial scales
 - Known degree of confidence
 - Spatially balanced
- Targeted site selection




NAMP - Step 5: Statistical framework




Road map for developing BLM's NAMP



NAMP – Example: Benthic macroinvertebrates




NAMP – Example: Benthic macroinvertebrates



NAMP – Example: Benthic macroinvertebrates

- BLM use of macroinvertebrates:
 - ~2300 unique stations
 - Only ~1200 of the ~2300 stations contain reliable data
 - Only ~58% of FO personnel use data to make management decisions




NAMP – Example: Benthic macroinvertebrates

- BMI good core indicator because:
 - Can be measured with minimal observed bias
 - Documented relationship to the health of lotic system
 - Responds to disturbance at relevant time-scales
 - Used by regulatory agencies for status determinations
 - Actively used by multiple stated, federal and tribal entities
- Standardized collection methodologies: 2011 Instructional memorandum (IM OC-2011-044)
- Analytical framework for making status determinations



NAMP – Example: Benthic macroinvertebrates

- Analytical framework for making status determinations
 - Use of state specific MMI or OE models
 - Facilitates integration of state-based reference conditions



NAMP – Example: Benthic macroinvertebrates

Is O/E score $> 2 \times \text{S.D.}$ from reference average?

Yes

No

Were replicate samples
collected (>3)?

Site comparable to reference
= meeting beneficial uses

Yes

No

Site Impaired

O/E $> 3 \text{ S.D.}$?

Yes

No

Site impaired

More data needed




NAMP – Example: Benthic macroinvertebrates

- Site selection methodology (statistical design):

Reach-scale: targeted
(historic focus)




Landscape-scale: prob. based
(moving towards)




NAMP – Example: Benthic macroinvertebrates

- What is the biological condition of lotic wadeable, perennial streams on BLM lands throughout Utah?
 - UT: 5,200 km of perennial streams and rivers



NAMP – Example: Benthic macroinvertebrates

- Landscape-scale status and trend determinations, as well as stressor prioritization
- Data integration with existing state and federal programs



NAMP – Next steps

- Survey to prioritize ecosystem attributes (June, 2012)
 - Currently recruiting participants
- Workshop to select preliminary core indicators (Summer 2012)
- Demonstration projects:
 - Utah (ongoing): probabilistic state-wide surveys stratified by BLM district
 - CA (2012 start): probabilistic surveys of lentic and lotic resources for two large grazing allotments
 - Owyhee uplands (2013 start): integration of terrestrial and aquatic core indicators



Thank you...

Questions?

