

Numerical Analysis & Scientific Computing II

Module 1

Introduction

1.1 Computing vs scientific computing?

1.2 Pre-requisites



Akash Anand
MATH, IIT KANPUR

Numerical Analysis & Scientific Computing II

Module 1

Introduction

1.1 Computing vs scientific computing?

1.2 Pre-requisites



Introduction: Computing vs scientific computing



What is scientific computing?

Introduction: Computing vs scientific computing



What is scientific computing?


Scientific method



Introduction: Computing vs scientific computing

What is scientific computing?


Scientific method



Introduction: Computing vs scientific computing

What is scientific computing?


Scientific method



Introduction: Computing vs scientific computing

What is scientific computing?


Scientific method



Introduction: Computing vs scientific computing

What is scientific computing?


Scientific method



Introduction: Computing vs scientific computing

What is scientific computing?


Scientific method



Introduction: Computing vs scientific computing

What is scientific computing?

Scientific method




Introduction: Computing vs scientific computing

What is scientific computing?

Scientific method


*We can do things with
computation that we could
not do with experiments ...*



Introduction: Computing vs scientific computing

What is scientific computing?

Scientific method




*We can do things with
computation that we could
not do with experiments ...*

*... can go in inaccessible
scales*

Introduction: Computing vs scientific computing

What is scientific computing?

Scientific method



We can do things with computation that we could not do with experiments ...


... can go in inaccessible scales

... can go to environments that are impossible to recreate or are too dangerous to create

Introduction: Computing vs scientific computing

What is scientific computing?

Scientific method



We can do things with computation that we could not do with experiments ...

... can go in inaccessible scales

... can go to environments that are impossible to recreate or are too dangerous to create

... cost advantage


...

Introduction: Computing vs scientific computing

What is scientific computing?

Example

A numerical simulation showing the gravitational radiation emitted by the violent merger of two black holes



Source:
Approaching the Black by Numerical Simulations
by Christian Fendt

We can do things with computation that we could not do with experiments ...

... can go in inaccessible scales

... can go to environments that are impossible to recreate or are too dangerous to create

... cost advantage

...


Introduction: Computing vs scientific computing



What is scientific computing?

Example

Visualization of the
instantaneous vortical
structures around the
golf ball



Source:

*Numerical Investigation of the Flow Past a Rotating Golf Ball
and its comparison with a rotating smooth sphere by Jing Li,
Makoto Tsubokura, Masaya Tsunoda*

*We can do things with
computation that we could
not do with experiments ...*

*... can go in inaccessible
scales*

*... can go to environments
that are impossible to
recreate or are too
dangerous to create*

... cost advantage

...


Introduction: Computing vs scientific computing



What is scientific computing?

Example

Abdominal aortic
aneurysm



Source:

Team for Advanced Flow Simulation and Modeling
(<https://www.tafsm.org/PROJ/CVFSI/PSCMADBF>)

*We can do things with
computation that we could
not do with experiments ...*

*... can go in inaccessible
scales*

*... can go to environments
that are impossible to
recreate or are too
dangerous to create*

... cost advantage

...


Introduction: Computing vs scientific computing



What is scientific computing?

Example

Updraft in a
hypothetical
supercell simulation



Source:

Texas Advanced Computer Center, University of Texas at Austin

*We can do things with
computation that we could
not do with experiments ...*

*... can go in inaccessible
scales*

*... can go to environments
that are impossible to
recreate or are too
dangerous to create*

... cost advantage

...


Introduction: Computing vs scientific computing



What is scientific computing?

Example

Wave-satellite
interaction



Source: Anand et al.

*We can do things with
computation that we could
not do with experiments ...*

*... can go in inaccessible
scales*

*... can go to environments
that are impossible to
recreate or are too
dangerous to create*


... cost advantage

...

Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?


The basic paradigm



Introduction: Computing vs scientific computing

What is scientific computing?

The basic paradigm



Introduction: Computing vs scientific computing



A simple example

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by $1/10$ km/h for each kilometer travels.

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by $1/10$ km/h for each kilometer travels.

Introduction: Computing vs scientific computing

A simple example



Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t) \qquad \frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “**first discretization**” of differential equations is due to Euler in 1768.

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “**first discretization**” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives

195.39647 km

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives

195.39647 km

Recomputing using 600 intervals of 1 minute gives

189.72820 km

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives

195.39647 km

Recomputing using 600 intervals of 1 minute gives

189.72820 km

36,000 intervals of 1 second gives

189.63770 km

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives

195.39647 km

Recomputing using 600 intervals of 1 minute gives

189.72820 km

36,000 intervals of 1 second gives

189.63770 km

36,000,000 intervals of 1 millisecond gives

189.63617 km

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives	195.39647 km
Recomputing using 600 intervals of 1 minute gives	189.72820 km
36,000 intervals of 1 second gives	189.63770 km
36,000,000 intervals of 1 millisecond gives	189.63617 km
36,000,000,000 intervals of 1 microsecond gives	189.63617 km

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.
How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives
Recomputing using 600 intervals of 1 minute gives
36,000 intervals of 1 second gives
36,000,000 intervals of 1 millisecond gives
36,000,000,000 intervals of 1 microsecond gives

195.39647 km
189.72820 km
189.63770 km
189.63617 km
189.63617 km

Numerical solution

Introduction: Computing vs scientific computing

A simple example

Physical system

A robotic vehicle departs at 30 km/h, but, due to battery drain, its speed decreases by 1/10 km/h for each kilometer travels.

How far does it go in 10 hours?

If t time is the time since departure, $v(t)$ is the speed (in km/h) at time t , and $x(t)$ is the distance (in km) travelled at time t , then, the mathematical system/model is given by

$$v(t) = 30 - \frac{1}{10}x(t)$$

$$\frac{dx(t)}{dt} = 30 - \frac{1}{10}x(t)$$

Mathematical system

The “*first discretization*” of differential equations is due to Euler in 1768.

Discrete system

Euler’s method partitions the 10 hour time interval into many short intervals and successively computes the distance travelled in each interval using the speed at the start.

Initial speed is 30 km/h, so vehicle goes about 30 kilometers in the first hour. By then speed has dropped to 27 km/h, so it travels 27 kilometers in the second hour. Then, 24.3 kilometers in the third hour, etc.

Using 10 intervals of 1 hour gives
Recomputing using 600 intervals of 1 minute gives
36,000 intervals of 1 second gives
36,000,000 intervals of 1 millisecond gives
36,000,000,000 intervals of 1 microsecond gives

195.39647 km
189.72820 km
189.63770 km
189.63617 km
189.63617 km

Numerical solution

Reliability/accuracy?

Speed of calculation?

Numerical Analysis & Scientific Computing II

Module 1

Introduction

1.1 Computing vs scientific computing?

1.2 Pre-requisites



Akash Anand
MATH, IIT KANPUR

Introduction: Pre-requisites



*In the first course on **Numerical Analysis & Scientific Computing**, you should have seen the following topics:*

- ***Approximation** of*
 - *functions,*
 - *derivatives,*
 - *integrals.*

- ***Solution** of (system of)*
 - *linear equations and*
 - *non-linear equations.*

Introduction: Pre-requisites

In the first course on *Numerical Analysis & Scientific Computing*, you should have seen the following topics:

- *Approximation* of
 - *functions,*
 - *derivatives,*
 - *integrals.*

- *Solution* of (system of)
 - *linear equations* and
 - *non-linear equations.*

In addition, it is useful to know the *well-posedness of initial and/or boundary value problems for ODEs and PDEs*, a subject matter of theoretical courses on ODE/PDE.

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Initial Value Problems: First order system of ODE



If we learn to solve first order system of ODEs, then we can also solve a high order ODE provided it is explicit.

Initial Value Problems: First order system of ODE

If we learn to solve first order system of ODEs, then we can also solve a high order ODE provided it is explicit.

Recall that a k th order ODE is said to be explicit if it can be written in the form

$$u^{(k)} = f\left(t, u, u', u'', \dots, u^{(k-1)}\right)$$

where $f : \mathbb{R}^{kn+1} \rightarrow \mathbb{R}^n$.

Initial Value Problems: First order system of ODE

If we learn to solve first order system of ODEs, then we can also solve a high order ODE provided it is explicit.

Recall that a k th order ODE is said to be explicit if it can be written in the form

$$u^{(k)} = f\left(t, u, u', u'', \dots, u^{(k-1)}\right)$$

where $f : \mathbb{R}^{kn+1} \rightarrow \mathbb{R}^n$.

Introduce new variables

$$y_1(t) = u(t), y_2(t) = u'(t), \dots, y_k(t) = u^{(k-1)}(t)$$

so that the original k th order system becomes a system of kn first order equations

$$y' = \begin{bmatrix} y_1' \\ \vdots \\ y_{k-1}' \\ y_k' \end{bmatrix} = \begin{bmatrix} y_2 \\ \vdots \\ y_k \\ f(t, y_1, y_2, \dots, y_k) \end{bmatrix}$$

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Well-posedness



We want to study the methods for numerical solution of a first order system of ordinary differential equations with initial conditions,

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $y: [a, b] \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$.

Initial Value Problems: Well-posedness

We want to study the methods for numerical solution of a first order system of ordinary differential equations with initial conditions,

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $y: [a, b] \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$.

Before we study the numerical solution, we need to investigate the well-posedness of the problem, that is, we study if the problem has following three properties:

- (i) existence of a solution (**existence**),
- (ii) uniqueness of the solution (**uniqueness**), and
- (iii) continuous dependence of the solution of the data (**conditioning**).

Initial Value Problems: Well-posedness



We want to study the methods for numerical solution of a first order system of ordinary differential equations with initial conditions,

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $y: [a, b] \rightarrow \mathbb{R}^n$ and $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$.

Before we study the numerical solution, we need to investigate the well-posedness of the problem, that is, we study if the problem has following three properties:

- (i) existence of a solution (**existence**),
- (ii) uniqueness of the solution (**uniqueness**), and
- (iii) continuous dependence of the solution of the data (**conditioning**).

This investigation, however, is the main subject matter of the course on Ordinary Differential Equations (ODE) and we, in this course, will only recall the relevant discussions.

Initial Value Problems: Well-posedness



Initial Value Problems: Well-posedness



Let $D = [a, b] \times \Omega \subseteq \mathbb{R}^{n+1}$ be a closed and bounded set.

Suppose that $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is a **Lipschitz continuous function in y** on D , that is, there is a constant L such that for any $t \in [a, b]$ and for any y and $\hat{y} \in \Omega$,

$$\|f(t, \hat{y}) - f(t, y)\| \leq L \|\hat{y} - y\|.$$

Initial Value Problems: Well-posedness



Let $D = [a, b] \times \Omega \subseteq \mathbb{R}^{n+1}$ be a closed and bounded set.

Suppose that $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is a **Lipschitz continuous function in y** on D , that is, there is a constant L such that for any $t \in [a, b]$ and for any y and $\hat{y} \in \Omega$,

$$\|f(t, \hat{y}) - f(t, y)\| \leq L \|\hat{y} - y\|.$$

Recall, from your ODE course, that for such functions, the following Initial Value Problem (IVP)

$$y' = f(t, y), \quad y(t_0) = y_0,$$

has a unique solution in a subinterval of $[a, b]$ containing y_0 .

Initial Value Problems: Well-posedness

Let $D = [a, b] \times \Omega \subseteq \mathbb{R}^{n+1}$ be a closed and bounded set.

Suppose that $f: \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ is a **Lipschitz continuous function in y** on D , that is, there is a constant L such that for any $t \in [a, b]$ and for any y and $\hat{y} \in \Omega$,

$$\|f(t, \hat{y}) - f(t, y)\| \leq L \|\hat{y} - y\|.$$

Recall, from your ODE course, that for such functions, the following Initial Value Problem (IVP)

$$y' = f(t, y), \quad y(t_0) = y_0,$$

has a unique solution in a subinterval of $[a, b]$ containing y_0 .

Example: If f is differentiable, then f is Lipschitz continuous with

$$L = \max_{(t,y) \in D} \|f'(t, y)\|,$$

where f' is the $n \times n$ Jacobian matrix of f with respect to y , $[f'(t, y)]_{ij} = \frac{\partial f_i(t, y)}{\partial y_j}$.

Initial Value Problems: Well-posedness



Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Initial Value Problems: Well-posedness



Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Then, it can be shown that for any $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

Initial Value Problems: Well-posedness



Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Then, it can be shown that for any $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

If the function is also perturbed so that we have $\hat{y}' = \hat{f}(t, \hat{y})$, then ...

Initial Value Problems: Well-posedness

Let $\hat{y}(t)$ be the solution of the following IVP with a perturbed initial condition

$$y' = f(t, y), \quad y(t_0) = \hat{y}_0.$$

Then, it can be shown that for any $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

If the function is also perturbed so that we have $\hat{y}' = \hat{f}(t, \hat{y})$, then for $t \geq t_0$,

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\| + \frac{e^{L(t-t_0)} - 1}{L} \|\hat{f} - f\|,$$

where

$$\|\hat{f} - f\| = \max_{(t,y) \in D} \|\hat{f}(t, y) - f(t, y)\|.$$

These perturbation bounds show that the unique solution to the IVP is a continuous function of the problem data, and hence the problem is well-posed.

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Stability



*We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.*

Initial Value Problems: Stability



We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.

Also recall that a solution of the ODE $y' = f(t, y)$ is said to be **stable** if for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $\hat{y}(t)$ satisfied the ODE and $\|\hat{y}(t_0) - y(t_0)\| \leq \delta$, then $\|\hat{y}(t) - y(t)\| \leq \epsilon$ for all $t \geq t_0$.



We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.

Also recall that a solution of the ODE $y' = f(t, y)$ is said to be **stable** if for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $\hat{y}(t)$ satisfied the ODE and $\|\hat{y}(t_0) - y(t_0)\| \leq \delta$, then $\|\hat{y}(t) - y(t)\| \leq \epsilon$ for all $t \geq t_0$.

Thus, for a **stable solution**, if the initial value is perturbed, then the perturbed solution remains close to the original solution, which **rules out the exponential divergence of perturbed solution** allowed by the perturbation bound

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$



We will see that the stability of the solution plays a significant role in the propagation of error, introduced by a numerical method, over time; they can either get **amplified** or **diminished** depending on the stability.

Also recall that a solution of the ODE $y' = f(t, y)$ is said to be **stable** if for every $\epsilon > 0$, there exists a $\delta > 0$ such that if $\hat{y}(t)$ satisfied the ODE and $\|\hat{y}(t_0) - y(t_0)\| \leq \delta$, then $\|\hat{y}(t) - y(t)\| \leq \epsilon$ for all $t \geq t_0$.

Thus, for a **stable solution**, if the initial value is perturbed, then the perturbed solution remains close to the original solution, which **rules out the exponential divergence of perturbed solution** allowed by the perturbation bound

$$\|\hat{y}(t) - y(t)\| \leq e^{L(t-t_0)} \|\hat{y}_0 - y_0\|.$$

A stable solution is said to be **asymptotically stable** if $\|\hat{y}(t) - y(t)\| \rightarrow 0$ as $t \rightarrow \infty$. This stronger form means that the original and perturbed solution not only remain close to each other, but they converge toward each other over time.

Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are ...

Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

Initial Value Problems: Stability



Examples:


- (i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.
- (ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

Initial Value Problems: Stability



Examples:

- (i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.
- (ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.




Initial Value Problems: Stability

Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\text{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.





Initial Value Problems: Stability

Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\text{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.



Initial Value Problems: Stability



Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\text{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\text{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.





Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\operatorname{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\operatorname{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.

If $\operatorname{Re}(\lambda) = 0$, then ...



Initial Value Problems: Stability

Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\operatorname{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\operatorname{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.

If $\operatorname{Re}(\lambda) = 0$, then ... (**Exercise**)



Initial Value Problems: Stability

Examples:

(i) Solutions of $y' = b$, for a given b , are stable but not asymptotically stable.

(ii) Consider the IVP $y' = \lambda y$, for a given constant λ , $y(0) = y_0$. The solution is $y(t) = y_0 e^{\lambda t}$.

If $\operatorname{Re}(\lambda) > 0$, then all non-zero solutions grow exponentially so that any two solutions diverge away, and hence every solution is unstable.

If $\operatorname{Re}(\lambda) < 0$, then all non-zero solutions decay exponentially so that any two solutions converge toward each other, and hence, every solution is not only stable but asymptotically stable.

If $\operatorname{Re}(\lambda) = 0$, then ... (**Exercise**)

(iii) A linear homogeneous system of ODEs with constant coefficients has the form

$$y' = Ay,$$

where A is an $n \times n$ matrix. Suppose we have the initial condition $y(0) = y_0$. Discuss the stability of the solutions if

- (a) A is diagonalizable, and
- (b) A is not diagonalizable.

(**Exercise**)

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.0 First-order system of ODE

2.1 Well-posedness

2.2 Stability

2.3 Euler's method



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds$$

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

Initial Value Problems: Euler's method



Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule,

Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE


$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that


$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(T) = y(t_0) + \int_{t_0}^T y'(s) ds = y(t_0) + \int_{t_0}^T f(s, y(s)) ds$$



The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$



The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$

The simplest approximation for the integral is by using the left end-point rule, that is,



$$\int_a^b g(s) ds \approx (b - a)g(a)$$

$$f(t_k, y(t_k))$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$



The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$


yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



$$y_{k+1} = y_k + (t_{k+1} - t_k)f(t_k, y_k)$$



Initial Value Problems: Euler's method

Starting at time t_0 with the given initial value y_0 , we would like to track the solution trajectory governed by the ODE

$$y' = f(t, y)$$

The fundamental theorem of calculus tells us that

$$y(t_{k+1}) = y(t_k) + \int_{t_k}^{t_{k+1}} y'(s) ds = y(t_k) + \int_{t_k}^{t_{k+1}} f(s, y(s)) ds$$


The simplest approximation for the integral is by using the left end-point rule, that is,

$$\int_a^b g(s) ds \approx (b - a)g(a)$$

yielding the method

$$y(T) = y(t_0) + (T - t_0)f(t_0, y_0)$$

To have more control over the error, we can also do this in multiple steps as follows:



Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.1 Well-posedness

2.2 Stability

2.3 Euler's method

- Derivations



Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Euler's method



Derivation using Taylor series

Initial Value Problems: Euler's method



Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$



Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$

The Euler's method results from taking $y_k = y(t_k)$, $y'(t_k) = f(t_k, y_k)$ and dropping terms of second and higher orders!

Initial Value Problems: Euler's method

Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$

The Euler's method results from taking $y_k = y(t_k)$, $y'(t_k) = f(t_k, y_k)$ and dropping terms of second and higher orders!

Derivation using finite difference approximation

Initial Value Problems: Euler's method

Derivation using Taylor series

Consider the Taylor series

$$y(t_{k+1}) = y(t_k) + (t_{k+1} - t_k)y'(t_k) + \frac{(t_{k+1} - t_k)^2}{2}y''(t_k) + \dots$$

The Euler's method results from taking $y_k = y(t_k)$, $y'(t_k) = f(t_k, y_k)$ and dropping terms of second and higher orders!

Derivation using finite difference approximation

Replacing the $y'(t)$ in the ODE $y' = f(t, y)$ by a first order forward difference approximation, we obtain an algebraic equation

$$\frac{y_{k+1} - y_k}{t_{k+1} - t_k} = f(t_k, y_k)$$

that yields the Euler's method.

Initial Value Problems: Euler's method



Derivation using polynomial interpolation

Initial Value Problems: Euler's method



Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

Initial Value Problems: Euler's method



Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$



Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} .

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined.

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined. To find the coefficients, we enforce that the predictor should be exact for first two monomials, that is, for $y(t) = e$ and $y(t) = te$, where e is the vector whose components are all ones.

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined. To find the coefficients, we enforce that the predictor should be exact for first two monomials, that is, for $y(t) = e$ and $y(t) = te$, where e is the vector whose components are all ones. Therefore, we have

$$e = \alpha e + \beta 0, \quad t_{k+1}e = \alpha t_k e + \beta e$$

Initial Value Problems: Euler's method

Derivation using polynomial interpolation

One point Hermite polynomial $p(t)$ that matches the function and derivative data at $t = t_k$, that is,

$$p(t_k) = y(t_k) (= y_k), \quad p'(t) = y'(t_k) (= f(t_k, y_k)),$$

is given by

$$p(t) = y_k + (t - t_k)f(t_k, y_k)$$

The Euler's method results from evaluating this polynomial at $t = t_{k+1}$.

Derivation using the method of undetermined coefficients

At $t = t_k$, we know the values y_k and y'_k , and based on these values we want to predict the value y_{k+1} . A predictor based on their linear combination has the form

$$y_{k+1} = \alpha y_k + \beta y'_k$$

where α and β are coefficients to be determined. To find the coefficients, we enforce that the predictor should be exact for first two monomials, that is, for $y(t) = e$ and $y(t) = te$, where e is the vector whose components are all ones. Therefore, we have

$$e = \alpha e + \beta 0, \quad t_{k+1}e = \alpha t_k e + \beta e$$

implying $\alpha = 1$ and $\beta = t_{k+1} - t_k$ resulting in the Euler's method.

Numerical Analysis & Scientific Computing II

Module 2

Initial Value Problems

2.1 Well-posedness

2.2 Stability

2.3 Euler's method

**- Errors and error
propagation**




Akash Anand
MATH, IIT KANPUR

Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method



Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$




Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – *one step error* or *single step error*.



Initial Value Problems: Euler's method


Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – *one step error* or *single step error*.

These local errors accumulate over time to produce a *global error*.



Initial Value Problems: Euler's method

Example


Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – **one step error** or **single step error**.

These local errors accumulate over time to produce a **global error**.

Note that how local errors accumulate is not always clear and it depends on the underlying solution.



Initial Value Problems: Euler's method

Example


Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – **one step error** or **single step error**.

These local errors accumulate over time to produce a **global error**.

Note that how local errors accumulate is not always clear and it depends on the underlying solution.



With each step, we hop from one solution of the ODE to another.

Initial Value Problems: Euler's method

Example


Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$

An error is introduced at each step of the method – **one step error** or **single step error**.

These local errors accumulate over time to produce a **global error**.

Note that how local errors accumulate is not always clear and it depends on the underlying solution.



With each step, we hop from one solution of the ODE to another.

Local errors get amplified if the solutions to the ODE are unstable.

Initial Value Problems: Euler's method

Example

Let us solve $y' = y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = h_k = t_{k+1} - t_k = 0.5$.

$$y_1 = 1 + h, \quad y_2 = y_1 + hy_1 = (1 + h)^2, \quad y_3 = y_2 + hy_2 = (1 + h)^3, \dots, \quad y_k = (1 + h)^k, \dots$$



With each step, we hop from one solution of the ODE to another.

Local errors get amplified if the solutions to the ODE are unstable.

Source: <https://youtu.be/714HgSO0h7g>

Initial Value Problems: Euler's method



Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.

Initial Value Problems: Euler's method



Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.


$$y_1 = 1 - h, \quad y_2 = y_1 - hy_1 = (1 - h)^2, \quad y_3 = y_2 - hy_2 = (1 - h)^3, \dots, \quad y_k = (1 - h)^k, \dots$$

Initial Value Problems: Euler's method

Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.

$$y_1 = 1 - h, \quad y_2 = y_1 - hy_1 = (1 - h)^2, \quad y_3 = y_2 - hy_2 = (1 - h)^3, \dots, \quad y_k = (1 - h)^k, \dots$$




Initial Value Problems: Euler's method

Example

Let us solve $y' = -y$, $y(0) = 1$ using the Euler's method taking the uniform step size $h = 0.5$.

$$y_1 = 1 - h, \quad y_2 = y_1 - hy_1 = (1 - h)^2, \quad y_3 = y_2 - hy_2 = (1 - h)^3, \dots, \quad y_k = (1 - h)^k, \dots$$



For an equation with stable solutions, the errors in the numerical solution do not grow, and for equations with asymptotically stable solutions, the errors diminish with time.