

Lesson 4

Numerical Solution of PDE

4.2 Parabolic PDE

- *Semi-discretization*
- *Full finite difference discretization*
- ***Fourier Analysis***



Numerical Methods for PDE: Parabolic PDE

Another useful way to analyze is to use Fourier analysis.

Recall that, on $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$ and $\varphi_m(x) = \sin \pi m x$, $m = 1, \dots, N-1$, form an orthogonal basis.

Also,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (\cos \pi m h - 1) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2},$$

where the eigenvalues satisfy

$$0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < \frac{4}{h^2}.$$

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Thus,

$$\frac{da_m^h}{dt} = -c a_m^h \lambda_m \quad \text{so} \quad a_m^h(t) = a_m^h(0) e^{-c \lambda_m t}.$$

Numerical Methods for PDE: Parabolic PDE

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Thus,

$$\frac{da_m^h}{dt} = -c a_m^h \lambda_m \quad \text{so} \quad a_m^h(t) = a_m^h(0) e^{-c \lambda_m t}.$$

The numbers $a_m^h(0)$ are the coefficients in the discrete Fourier sine transform of the initial data:

$$u_0(x) = \sum_{m=1}^{N-1} a_m^h(0) \varphi_m(x), \quad x \in \bar{I}_h.$$

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c \lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

Numerical Methods for PDE: Parabolic PDE

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c\lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution at time $t = jk$ as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

The difference equation then gives

$$\sum_{m=1}^{N-1} A_m^{j+1} \varphi_m(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x) - ck \sum_{m=1}^{N-1} A_m^j \lambda_m \varphi_m(x)$$

yielding

$$A_m^{j+1} = (1 - ck\lambda_m) A_m^j.$$

It follows that

$$A_m^j = (1 - ck\lambda_m)^j a_m^h(0), \quad u^j = \sum_{m=1}^{N-1} (1 - ck\lambda_m)^j a_m^h(0) \varphi_m.$$

If we assume $ck/h^2 \leq 1/2$, then $ck\lambda_m \leq ck(4/h^2) \leq 2$ and hence $|1 - ck\lambda_m| \leq 1$ for all m and the solution remains bounded. On the other hand, if $|1 - ck\lambda_m| > 1$ for some m , the initial data will increase exponentially.

Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k\|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m|\right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h.$$

*Because of the condition $ck/h^2 \leq 1/2$, which we know is not only sufficient but necessary for stability, forward-centered difference method is called **conditionally stable**.*

Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k\|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m|\right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h.$$

Because of the condition $ck/h^2 \leq 1/2$, which we know is not only sufficient but necessary for stability, forward-centered difference method is called **conditionally stable**.

From stability, we obtain the convergence result in the same way as earlier. Let $e_n^j = u_n^j - u(nh, jk)$, we have

$$\begin{aligned} \frac{e_n^{j+1} - e_n^j}{k} &= c \frac{e_{n+1}^j - 2e_n^j + e_{n-1}^j}{h^2} - \ell_n^j, \quad 0 < n < N, j = 0, 1, \dots, M-1, \\ e_0^j &= e_N^j = 0, \quad j = 0, 1, \dots, M-1, \\ e_n^0 &= 0, \quad 0 < n < N. \end{aligned}$$

Numerical Methods for PDE: Parabolic PDE

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k \|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m| \right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h$$

Because of the condition $ck/h^2 \leq 1/2$, which we know is not only sufficient but necessary for stability, forward-centered difference method is called **conditionally stable**.

From stability, we obtain the convergence result in the same way as earlier. Let $e_n^j = u_n^j - u(nh, jk)$, we have

$$\begin{aligned} \frac{e_n^{j+1} - e_n^j}{k} &= c \frac{e_{n+1}^j - 2e_n^j + e_{n-1}^j}{h^2} - \ell_n^j, & 0 < n < N, j = 0, 1, \dots, M-1, \\ e_0^j &= e_N^j = 0, & j = 0, 1, \dots, M-1, \\ e_n^0 &= 0, & 0 < n < N. \end{aligned}$$

The stability result then yields

$$\|e^j\|_h \leq \left(\max_m |1 - ck\lambda_m| \right)^j \|e^0\|_h + Mk \max_j \|\ell^j\|_h \leq T \max_j \|\ell^j\|_h.$$

Lesson 4

Numerical Solution of PDE

4.2 Parabolic PDE

- *Semi-discretization*
- *Full finite difference discretization*
- *Fourier Analysis*
- ***Unconditional Stability***



Numerical Methods for PDE: Parabolic PDE



Can we design a scheme that is unconditionally stable?

Numerical Methods for PDE: Parabolic PDE



Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

Numerical Methods for PDE: Parabolic PDE

Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

The backward-centered difference method for the heat equation leads to

$$\frac{u_n^{j+1} - u_n^j}{k} = c \frac{u_{n+1}^{j+1} - 2u_n^{j+1} + u_{n-1}^{j+1}}{h^2} + f_n^{j+1}, \quad 0 < n < N, j = 0, 1, \dots, M-1,$$
$$u_0^j = u_N^j = 0, \quad j = 0, 1, \dots, M,$$
$$u_n^0 = u_0(nh), \quad 0 < n < N.$$

Numerical Methods for PDE: Parabolic PDE

Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

The backward-centered difference method for the heat equation leads to

$$\frac{u_n^{j+1} - u_n^j}{k} = c \frac{u_{n+1}^{j+1} - 2u_n^{j+1} + u_{n-1}^{j+1}}{h^2} + f_n^{j+1}, \quad 0 < n < N, j = 0, 1, \dots, M-1,$$

$$u_0^j = u_N^j = 0, \quad j = 0, 1, \dots, M,$$

$$u_n^0 = u_0(nh), \quad 0 < n < N.$$

Thus, u_n^{j+1} can be obtained by solving a tridiagonal system

$$-\lambda u_{n+1}^{j+1} + (1 + 2\lambda)u_n^{j+1} - \lambda u_{n-1}^{j+1} = u_n^j + k f_n^{j+1}, \quad u_0^{j+1} = u_N^{j+1} = 0.$$

Numerical Methods for PDE: Parabolic PDE

Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

The backward-centered difference method for the heat equation leads to

$$\frac{u_n^{j+1} - u_n^j}{k} = c \frac{u_{n+1}^{j+1} - 2u_n^{j+1} + u_{n-1}^{j+1}}{h^2} + f_n^{j+1}, \quad 0 < n < N, j = 0, 1, \dots, M-1,$$

$$u_0^j = u_N^j = 0, \quad j = 0, 1, \dots, M,$$

$$u_n^0 = u_0(nh), \quad 0 < n < N.$$

Thus, u_n^{j+1} can be obtained by solving a tridiagonal system

$$-\lambda u_{n+1}^{j+1} + (1 + 2\lambda)u_n^{j+1} - \lambda u_{n-1}^{j+1} = u_n^j + k f_n^{j+1}, \quad u_0^{j+1} = u_N^{j+1} = 0.$$

It is easy to see that the local truncation error for the scheme is again $O(k + h^2)$.

Numerical Methods for PDE: Parabolic PDE

Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

The backward-centered difference method for the heat equation leads to

$$\frac{u_n^{j+1} - u_n^j}{k} = c \frac{u_{n+1}^{j+1} - 2u_n^{j+1} + u_{n-1}^{j+1}}{h^2} + f_n^{j+1}, \quad 0 < n < N, j = 0, 1, \dots, M-1,$$

$$u_0^j = u_N^j = 0, \quad j = 0, 1, \dots, M,$$

$$u_n^0 = u_0(nh), \quad 0 < n < N.$$

Thus, u_n^{j+1} can be obtained by solving a tridiagonal system

$$-\lambda u_{n+1}^{j+1} + (1 + 2\lambda)u_n^{j+1} - \lambda u_{n-1}^{j+1} = u_n^j + k f_n^{j+1}, \quad u_0^{j+1} = u_N^{j+1} = 0.$$

It is easy to see that the local truncation error for the scheme is again $O(k + h^2)$.

We again use the Fourier analysis to determine the stability of the scheme. In operator form, the method reads

$$\frac{u^{j+1} - u^j}{k} = c D_h^2 u^{j+1} + f^{j+1},$$

Numerical Methods for PDE: Parabolic PDE

Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

The backward-centered difference method for the heat equation leads to

$$\frac{u_n^{j+1} - u_n^j}{k} = c \frac{u_{n+1}^{j+1} - 2u_n^{j+1} + u_{n-1}^{j+1}}{h^2} + f_n^{j+1}, \quad 0 < n < N, j = 0, 1, \dots, M-1,$$

$$u_0^j = u_N^j = 0, \quad j = 0, 1, \dots, M,$$

$$u_n^0 = u_0(nh), \quad 0 < n < N.$$

Thus, u_n^{j+1} can be obtained by solving a tridiagonal system

$$-\lambda u_{n+1}^{j+1} + (1 + 2\lambda)u_n^{j+1} - \lambda u_{n-1}^{j+1} = u_n^j + k f_n^{j+1}, \quad u_0^{j+1} = u_N^{j+1} = 0.$$

It is easy to see that the local truncation error for the scheme is again $O(k + h^2)$.

We again use the Fourier analysis to determine the stability of the scheme. In operator form, the method reads

$$\frac{u^{j+1} - u^j}{k} = c D_h^2 u^{j+1} + f^{j+1},$$

that is,

$$u^{j+1} = (I - ck D_h^2)^{-1} (u^j + k f^{j+1}).$$

Numerical Methods for PDE: Parabolic PDE

Can we design a scheme that is unconditionally stable?

Try backward Euler method in time!

The backward-centered difference method for the heat equation leads to

$$\frac{u_n^{j+1} - u_n^j}{k} = c \frac{u_{n+1}^{j+1} - 2u_n^{j+1} + u_{n-1}^{j+1}}{h^2} + f_n^{j+1}, \quad 0 < n < N, j = 0, 1, \dots, M-1,$$

$$u_0^j = u_N^j = 0, \quad j = 0, 1, \dots, M,$$

$$u_n^0 = u_0(nh), \quad 0 < n < N.$$

Thus, u_n^{j+1} can be obtained by solving a tridiagonal system

$$-\lambda u_{n+1}^{j+1} + (1 + 2\lambda)u_n^{j+1} - \lambda u_{n-1}^{j+1} = u_n^j + k f_n^{j+1}, \quad u_0^{j+1} = u_N^{j+1} = 0.$$

It is easy to see that the local truncation error for the scheme is again $O(k + h^2)$.

We again use the Fourier analysis to determine the stability of the scheme. In operator form, the method reads

$$\frac{u^{j+1} - u^j}{k} = c D_h^2 u^{j+1} + f^{j+1},$$

that is,

$$u^{j+1} = (I - ck D_h^2)^{-1} (u^j + k f^{j+1}).$$

The operator $I - ck D_h^2$ has eigenvalues $1 + ck\lambda_m$ which are all greater than one, so $\|(I - ck D_h^2)^{-1} v\|_h \leq \|v\|_h$.



We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$



Numerical Methods for PDE: Parabolic PDE

We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$

Hence, we have

$$\|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h$$

Numerical Methods for PDE: Parabolic PDE

We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$

Hence, we have

$$\|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h$$

and the stability result

$$\max_{1 \leq j \leq M} \|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h.$$

Numerical Methods for PDE: Parabolic PDE

We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$

Hence, we have

$$\|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h$$

and the stability result

$$\max_{1 \leq j \leq M} \|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h.$$

As we did not have to make any assumption on the relation between k and h , the backward-centered difference scheme is **unconditionally stable**.

Numerical Methods for PDE: Parabolic PDE

We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$

Hence, we have

$$\|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h$$

and the stability result

$$\max_{1 \leq j \leq M} \|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h.$$

As we did not have to make any assumption on the relation between k and h , the backward-centered difference scheme is **unconditionally stable**.

Combining the stability estimate with the local truncation error, we obtain the error estimate

$$\max_{1 \leq j \leq M} \|e^j\|_h = O(k + h^2)$$

for the method.

Numerical Methods for PDE: Parabolic PDE

We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$

Hence, we have

$$\|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h$$

and the stability result

$$\max_{1 \leq j \leq M} \|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h.$$

As we did not have to make any assumption on the relation between k and h , the backward-centered difference scheme is **unconditionally stable**.

Combining the stability estimate with the local truncation error, we obtain the error estimate

$$\max_{1 \leq j \leq M} \|e^j\|_h = O(k + h^2)$$

for the method.

If we use trapezoidal rule to discretize in time, we get the **Crank-Nicolson method**:

$$\frac{u^{j+1} - u^j}{k} = \frac{c}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{1}{2} (f^j + f^{j+1}).$$

Numerical Methods for PDE: Parabolic PDE

We, therefore, have

$$\|u^{j+1}\|_h = \|(I - ckD_h^2)^{-1}(u^j + kf^{j+1})\|_h \leq \|u^j\|_h + k\|f^j\|_h, \quad j = 0, 1, \dots, M-1.$$

Hence, we have

$$\|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h$$

and the stability result

$$\max_{1 \leq j \leq M} \|u^j\|_h \leq \|u^0\|_h + T \max_{1 \leq j \leq M} \|f^j\|_h.$$

As we did not have to make any assumption on the relation between k and h , the backward-centered difference scheme is **unconditionally stable**.

Combining the stability estimate with the local truncation error, we obtain the error estimate

$$\max_{1 \leq j \leq M} \|e^j\|_h = O(k + h^2)$$

for the method.

If we use trapezoidal rule to discretize in time, we get the **Crank-Nicolson method**:

$$\frac{u^{j+1} - u^j}{k} = \frac{c}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{1}{2} (f^j + f^{j+1}).$$

Using the Taylor expansion about the point $(nh, (j + 1/2)k)$, it is straightforward to show that the local truncation error is $O(k^2 + h^2)$, so the Crank-Nicolson method is second order in both space and time.

Numerical Methods for PDE: Parabolic PDE

If we use trapezoidal rule to discretize in time, we get the **Crank-Nicolson method**:

$$\frac{u^{j+1} - u^j}{k} = \frac{c}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{1}{2} (f^j + f^{j+1}).$$

Using the Taylor expansion about the point $(nh, (j + 1/2)k)$, it is straightforward to show that the local truncation error is $O(k^2 + h^2)$, so the Crank-Nicolson method is second order in both space and time.

To get the operator equation for the method, we write

$$u^{j+1} = u^j + \frac{ck}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{k}{2} (f^j + f^{j+1}),$$

that is

$$u^{j+1} - \frac{ck}{2} D_h^2 u^{j+1} = u^j + \frac{ck}{2} D_h^2 u^j + \frac{k}{2} (f^j + f^{j+1}).$$

Numerical Methods for PDE: Parabolic PDE

If we use trapezoidal rule to discretize in time, we get the **Crank-Nicolson method**:

$$\frac{u^{j+1} - u^j}{k} = \frac{c}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{1}{2} (f^j + f^{j+1}).$$

Using the Taylor expansion about the point $(nh, (j + 1/2)k)$, it is straightforward to show that the local truncation error is $O(k^2 + h^2)$, so the Crank-Nicolson method is second order in both space and time.

To get the operator equation for the method, we write

$$u^{j+1} = u^j + \frac{ck}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{k}{2} (f^j + f^{j+1}),$$

that is

$$u^{j+1} - \frac{ck}{2} D_h^2 u^{j+1} = u^j + \frac{ck}{2} D_h^2 u^j + \frac{k}{2} (f^j + f^{j+1}).$$

Thus

$$u^{j+1} = \left(I - \frac{1}{2} ck D_h^2\right)^{-1} \left(I + \frac{1}{2} ck D_h^2\right) u^j + \frac{k}{2} \left(I - \frac{1}{2} ck D_h^2\right)^{-1} (f^j + f^{j+1}).$$

Numerical Methods for PDE: Parabolic PDE

If we use trapezoidal rule to discretize in time, we get the **Crank-Nicolson method**:

$$\frac{u^{j+1} - u^j}{k} = \frac{c}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{1}{2} (f^j + f^{j+1}).$$

Using the Taylor expansion about the point $(nh, (j + 1/2)k)$, it is straightforward to show that the local truncation error is $O(k^2 + h^2)$, so the Crank-Nicolson method is second order in both space and time.

To get the operator equation for the method, we write

$$u^{j+1} = u^j + \frac{ck}{2} (D_h^2 u^j + D_h^2 u^{j+1}) + \frac{k}{2} (f^j + f^{j+1}),$$

that is

$$u^{j+1} - \frac{ck}{2} D_h^2 u^{j+1} = u^j + \frac{ck}{2} D_h^2 u^j + \frac{k}{2} (f^j + f^{j+1}).$$

Thus

$$u^{j+1} = \left(I - \frac{1}{2} ck D_h^2 \right)^{-1} \left(I + \frac{1}{2} ck D_h^2 \right) u^j + \frac{k}{2} \left(I - \frac{1}{2} ck D_h^2 \right)^{-1} (f^j + f^{j+1}).$$

The eigenvalues of $\left(I - \frac{1}{2} ck D_h^2 \right)^{-1} \left(I + \frac{1}{2} ck D_h^2 \right)$ is $(1 - ck\lambda_m/2)/(1 + ck\lambda_m/2)$ which is less than 1. Therefore, we get unconditional stability. The Crank-Nicolson method converges with $O(k^2 + h^2)$.

Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.3 Hyperbolic PDE



Akash Anand
MATH, IIT KANPUR

Numerical Methods for PDE: Hyperbolic PDE



For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

Numerical Methods for PDE: Hyperbolic PDE



For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.



For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.

For the solution of the pure initial value problem, if u is the exact solution, consider $U(t) = u(x_0 + ct, t)$.

Numerical Methods for PDE: Hyperbolic PDE



For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.

For the solution of the pure initial value problem, if u is the exact solution, consider $U(t) = u(x_0 + ct, t)$. Then,

$$\frac{dU}{dt} = \frac{\partial u}{\partial x} \frac{d(x_0 + ct)}{dt} + \frac{\partial u}{\partial t} \equiv 0.$$

Numerical Methods for PDE: Hyperbolic PDE



For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.

For the solution of the pure initial value problem, if u is the exact solution, consider $U(t) = u(x_0 + ct, t)$. Then,

$$\frac{dU}{dt} = \frac{\partial u}{\partial x} \frac{d(x_0 + ct)}{dt} + \frac{\partial u}{\partial t} \equiv 0.$$

So $U(t) = U(0) = u(x_0, 0) = u_0(x_0)$ where u_0 is the initial data.

Numerical Methods for PDE: Hyperbolic PDE

For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.

For the solution of the pure initial value problem, if u is the exact solution, consider $U(t) = u(x_0 + ct, t)$. Then,

$$\frac{dU}{dt} = \frac{\partial u}{\partial x} \frac{d(x_0 + ct)}{dt} + \frac{\partial u}{\partial t} \equiv 0.$$

So $U(t) = U(0) = u(x_0, 0) = u_0(x_0)$ where u_0 is the initial data.

Thus, for $x \in \mathbb{R}$, $t > 0$, we have

$$u(x, t) = u((x - ct) + ct, t) = u_0(x - ct).$$

Numerical Methods for PDE: Hyperbolic PDE

For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.

For the solution of the pure initial value problem, if u is the exact solution, consider $U(t) = u(x_0 + ct, t)$. Then,

$$\frac{dU}{dt} = \frac{\partial u}{\partial x} \frac{d(x_0 + ct)}{dt} + \frac{\partial u}{\partial t} \equiv 0.$$

So $U(t) = U(0) = u(x_0, 0) = u_0(x_0)$ where u_0 is the initial data.

Thus, for $x \in \mathbb{R}$, $t > 0$, we have

$$u(x, t) = u((x - ct) + ct, t) = u_0(x - ct).$$

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$).

Numerical Methods for PDE: Hyperbolic PDE

For the first example, we take the advection equation

$$\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} = 0,$$

where c is a constant. We consider an initial value problem so that the function $u = u(x, t)$ is given when $t = 0$ and is to be found for $t > 0$.

The spatial domain can be an interval in which case we will need to impose boundary conditions to obtain a well posed initial value-boundary value problem, or the entire real line (no boundaries) for which the pure initial value problem is well-posed.

For the solution of the pure initial value problem, if u is the exact solution, consider $U(t) = u(x_0 + ct, t)$. Then,

$$\frac{dU}{dt} = \frac{\partial u}{\partial x} \frac{d(x_0 + ct)}{dt} + \frac{\partial u}{\partial t} \equiv 0.$$

So $U(t) = U(0) = u(x_0, 0) = u_0(x_0)$ where u_0 is the initial data.

Thus, for $x \in \mathbb{R}$, $t > 0$, we have


$$u(x, t) = u((x - ct) + ct, t) = u_0(x - ct).$$

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the **characteristics of the equation**.

Numerical Methods for PDE: Hyperbolic PDE




We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.



Numerical Methods for PDE: Hyperbolic PDE




We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.



Numerical Methods for PDE: Hyperbolic PDE




We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.



Numerical Methods for PDE: Hyperbolic PDE




We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.



Numerical Methods for PDE: Hyperbolic PDE




We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.



Numerical Methods for PDE: Hyperbolic PDE

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.


Suppose that the spatial domain is an interval, say $(0,1)$.



Numerical Methods for PDE: Hyperbolic PDE

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.


Suppose that the spatial domain is an interval, say $(0,1)$.



Numerical Methods for PDE: Hyperbolic PDE

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.

Suppose that the spatial domain is an interval, say $(0,1)$.




Numerical Methods for PDE: Hyperbolic PDE

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.

Suppose that the spatial domain is an interval, say $(0,1)$.

For determining solution at (x, t) with $x - ct < 0$, we need boundary condition at $x = 0$. This can be specified for all $t > 0$, and reads $u(0, t) = g(t)$ for a given g .



Numerical Methods for PDE: Hyperbolic PDE


We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.

Suppose that the spatial domain is an interval, say $(0,1)$.

For determining solution at (x, t) with $x - ct < 0$, we need boundary condition at $x = 0$. This can be specified for all $t > 0$, and reads $u(0, t) = g(t)$ for a given g .

Thus, to obtain a well-posed problem, we need

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, 0 < x < 1, t > 0, \\ u(x, 0) &= u_0(x), 0 < x < 1, \\ u(0, t) &= g(t), t > 0.\end{aligned}$$



Numerical Methods for PDE: Hyperbolic PDE

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.


Suppose that the spatial domain is an interval, say $(0,1)$.

For determining solution at (x, t) with $x - ct < 0$, we need boundary condition at $x = 0$. This can be specified for all $t > 0$, and reads $u(0, t) = g(t)$ for a given g .

Thus, to obtain a well-posed problem, we need

$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, 0 < x < 1, t > 0, \\ u(x, 0) &= u_0(x), 0 < x < 1, \\ u(0, t) &= g(t), t > 0.\end{aligned}$$

The only boundary condition needed is on the left or **inflow boundary**. It is not necessary or permissible to impose a boundary condition on the right or **outflow boundary**.



Numerical Methods for PDE: Hyperbolic PDE

We, therefore, see that the pure initial value problem for advection equation can be solved analytically/exactly. The initial data simply travels to the right with speed c (for $c > 0$). The solution u is constant along the lines $\{(x_0 + ct, t): t > 0\}$ called the characteristics of the equation.

Suppose that the spatial domain is an interval, say $(0,1)$.

For determining solution at (x, t) with $x - ct < 0$, we need boundary condition at $x = 0$. This can be specified for all $t > 0$, and reads $u(0, t) = g(t)$ for a given g .

Thus, to obtain a well-posed problem, we need


$$\begin{aligned}\frac{\partial u}{\partial t} + c \frac{\partial u}{\partial x} &= 0, 0 < x < 1, t > 0, \\ u(x, 0) &= u_0(x), 0 < x < 1, \\ u(0, t) &= g(t), t > 0.\end{aligned}$$

The only boundary condition needed is on the left or **inflow boundary**. It is not necessary or permissible to impose a boundary condition on the right or **outflow boundary**.

Now consider a system of the form

$$\frac{\partial u}{\partial t} + C \frac{\partial u}{\partial x} = 0,$$

where C is an $n \times n$ symmetric matrix (that is, has real eigenvalues). Suppose $C = S^{-1}DS$ with S invertible and D a diagonal matrix.



Numerical Methods for PDE: Hyperbolic PDE



Now consider a system of the form

$$\frac{\partial u}{\partial t} + C \frac{\partial u}{\partial x} = 0,$$

where C is an $n \times n$ symmetric matrix (that is, has real eigenvalues). Suppose $C = S^{-1}DS$ with S invertible and D a diagonal matrix.

If we change dependent variable by $v = Su$, then we get

$$\frac{\partial v}{\partial t} + D \frac{\partial v}{\partial x} = 0,$$

Numerical Methods for PDE: Hyperbolic PDE

Now consider a system of the form

$$\frac{\partial u}{\partial t} + C \frac{\partial u}{\partial x} = 0,$$

where C is an $n \times n$ symmetric matrix (that is, has real eigenvalues). Suppose $C = S^{-1}DS$ with S invertible and D a diagonal matrix.

If we change dependent variable by $v = Su$, then we get

$$\frac{\partial v}{\partial t} + D \frac{\partial v}{\partial x} = 0,$$

that is,

$$\frac{\partial v_i}{\partial t} + d_i \frac{\partial v_i}{\partial x} = 0, \quad i = 1, \dots, n.$$

Numerical Methods for PDE: Hyperbolic PDE

Now consider a system of the form

$$\frac{\partial u}{\partial t} + C \frac{\partial u}{\partial x} = 0,$$

where C is an $n \times n$ symmetric matrix (that is, has real eigenvalues). Suppose $C = S^{-1}DS$ with S invertible and D a diagonal matrix.

If we change dependent variable by $v = Su$, then we get

$$\frac{\partial v}{\partial t} + D \frac{\partial v}{\partial x} = 0,$$

that is,

$$\frac{\partial v_i}{\partial t} + d_i \frac{\partial v_i}{\partial x} = 0, \quad i = 1, \dots, n.$$

Thus, the system decouples into n advection equations whose speeds are the eigenvalues of the original coefficient matrix.

Numerical Methods for PDE: Hyperbolic PDE

Now consider a system of the form

$$\frac{\partial u}{\partial t} + C \frac{\partial u}{\partial x} = 0,$$

where C is an $n \times n$ symmetric matrix (that is, has real eigenvalues). Suppose $C = S^{-1}DS$ with S invertible and D a diagonal matrix.

If we change dependent variable by $v = Su$, then we get

$$\frac{\partial v}{\partial t} + D \frac{\partial v}{\partial x} = 0,$$

that is,

$$\frac{\partial v_i}{\partial t} + d_i \frac{\partial v_i}{\partial x} = 0, \quad i = 1, \dots, n.$$

Thus, the system decouples into n advection equations whose speeds are the eigenvalues of the original coefficient matrix.

Example:

Consider the wave equation

$$\frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2}.$$

Numerical Methods for PDE: Hyperbolic PDE

Now consider a system of the form

$$\frac{\partial u}{\partial t} + C \frac{\partial u}{\partial x} = 0,$$

where C is an $n \times n$ symmetric matrix (that is, has real eigenvalues). Suppose $C = S^{-1}DS$ with S invertible and D a diagonal matrix.

If we change dependent variable by $v = Su$, then we get

$$\frac{\partial v}{\partial t} + D \frac{\partial v}{\partial x} = 0,$$

that is,

$$\frac{\partial v_i}{\partial t} + d_i \frac{\partial v_i}{\partial x} = 0, \quad i = 1, \dots, n.$$

Thus, the system decouples into n advection equations whose speeds are the eigenvalues of the original coefficient matrix.

Example:

Consider the wave equation

$$\frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2}.$$

Let $u_1 = \partial w / \partial t$ and $u_2 = \partial w / \partial x$. Then,



Example:

Consider the wave equation

$$\frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2}.$$

Let $u_1 = \partial w / \partial t$ and $u_2 = \partial w / \partial x$. Then,

$$\frac{\partial}{\partial t} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \frac{\partial}{\partial x} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = 0.$$

Numerical Methods for PDE: Hyperbolic PDE

Example:

Consider the wave equation

$$\frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2}.$$

Let $u_1 = \partial w / \partial t$ and $u_2 = \partial w / \partial x$. Then,

$$\frac{\partial}{\partial t} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \frac{\partial}{\partial x} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = 0.$$

Thus,

$$C = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} = S^{-1}DS, \quad S = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad D = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Numerical Methods for PDE: Hyperbolic PDE

Example:

Consider the wave equation

$$\frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2}.$$

Let $u_1 = \partial w / \partial t$ and $u_2 = \partial w / \partial x$. Then,

$$\frac{\partial}{\partial t} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \frac{\partial}{\partial x} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = 0.$$

Thus,

$$C = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} = S^{-1}DS, \quad S = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad D = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Finally, we find the solution (u_1, u_2) from $u_1 + u_2$ which is a wave moving from right to left and from $u_1 - u_2$, a wave moving to the right.