

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.1 Well-posedness

3.2 Shooting Method

3.3 Finite Difference Method



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Finite Difference Method



In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

Boundary Value Problems: Finite Difference Method



In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Boundary Value Problems: Finite Difference Method

In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Boundary Value Problems: Finite Difference Method

In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Introduce the mesh points $t_i = a + ih, i = 0, \dots, n + 1$, where $h = (b - a)/(n + 1)$.

Boundary Value Problems: Finite Difference Method

In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Introduce the mesh points $t_i = a + ih, i = 0, \dots, n + 1$, where $h = (b - a)/(n + 1)$.



Boundary Value Problems: Finite Difference Method

In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Introduce the mesh points $t_i = a + ih, i = 0, \dots, n + 1$, where $h = (b - a)/(n + 1)$.

We seek the approximations u_i to $u(t_i)$, that is, $u_i \approx u(t_i)$.



Boundary Value Problems: Finite Difference Method

In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Introduce the mesh points $t_i = a + ih, i = 0, \dots, n + 1$, where $h = (b - a)/(n + 1)$.

We seek the approximations u_i to $u(t_i)$, that is, $u_i \approx u(t_i)$.

We replace the derivatives appearing in the ODE by finite difference approximations

$$u'(t_i) \approx \frac{u_{i+1} - u_{i-1}}{2h}, \quad u''(t_i) \approx \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2}.$$



Boundary Value Problems: Finite Difference Method

In the shooting method, we start by approximately satisfying the ODE (using an IVP solver) and iterate until the boundary conditions are satisfied.

Alternatively, we can start by satisfying the boundary conditions and iterate until the ODE is satisfied approximately.

In the finite difference approach, we convert a BVP into a system of algebraic equations rather than a sequence of IVPs as in the shooting method.

Consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Introduce the mesh points $t_i = a + ih, i = 0, \dots, n + 1$, where $h = (b - a)/(n + 1)$.

We seek the approximations u_i to $u(t_i)$, that is, $u_i \approx u(t_i)$.

We replace the derivatives appearing in the ODE by finite difference approximations

$$u'(t_i) \approx \frac{u_{i+1} - u_{i-1}}{2h}, \quad u''(t_i) \approx \frac{u_{i+1} - 2u_i + u_{i-1}}{h^2}.$$

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), \quad i = 1, \dots, n,$$

to be solved for the unknowns $u_i, i = 1, \dots, n$.

Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns $u_i, i = 1, \dots, n$.

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \cdots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix}$$

Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns $u_i, i = 1, \dots, n$.

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & & \vdots \\ & & & & 1 & -2 & 1 \\ 0 & \cdots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns $u_i, i = 1, \dots, n$.

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \cdots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

which is denoted as

$$\frac{1}{h^2} Au = F(u) + g.$$

Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns $u_i, i = 1, \dots, n$.

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & \cdots & 0 \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \cdots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

which is denoted as

$$\frac{1}{h^2} Au = F(u) + g.$$

Thus, the Newton's method for solving the system of algebraic equations is given by

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} Au^{(m)} - F(u^{(m)}) - g \right]$$

where the Jacobian matrix is given by $[F(u)]_{ij} = [\partial f(t_i, u_i, (u_{i+1} - u_{i-1})/(2h))/\partial u_j]$.

Boundary Value Problems: Finite Difference Method

This yields a system of algebraic equations

$$\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} = f\left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h}\right), i = 1, \dots, n,$$

to be solved for the unknowns $u_i, i = 1, \dots, n$.

In the matrix form, we have

$$\frac{1}{h^2} \begin{bmatrix} -2 & 1 & 0 & & & \\ 1 & -2 & 1 & & & \\ \vdots & \vdots & & \ddots & \vdots & \\ & & & & 1 & -2 & 1 \\ 0 & \dots & & & 0 & 1 & -2 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} = \begin{bmatrix} f\left(t_1, u_1, \frac{u_2 - \alpha}{2h}\right) \\ f\left(t_2, u_2, \frac{u_3 - u_1}{2h}\right) \\ \vdots \\ f\left(t_n, u_n, \frac{\beta - u_n}{2h}\right) \end{bmatrix} - \begin{bmatrix} \frac{\alpha}{h^2} \\ \vdots \\ \frac{\beta}{h^2} \end{bmatrix}$$

which is denoted as

$$\frac{1}{h^2} Au = F(u) + g.$$

Thus, the Newton's method for solving the system of algebraic equations is given by

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} Au^{(m)} - F(u^{(m)}) - g \right]$$

where the Jacobian matrix is given by $[F(u)]_{ij} = [\partial f(t_i, u_i, (u_{i+1} - u_{i-1})/(2h))/\partial u_j]$.

Boundary Value Problems: Finite Difference Method

In the matrix form, we have

$$\frac{1}{h^2} Au = F(u) + g.$$

Thus, the Newton's method for solving the system of algebraic equations is given by

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} Au^{(m)} - F(u^{(m)}) - g \right]$$

where the Jacobian matrix is given by $[F'(u)]_{ij} = [\partial f(t_i, u_i, (u_{i+1} - u_{i-1})/(2h))/\partial u_j]$.

In particular,

$$\begin{aligned} [F'(u)]_{ii} &= f_2 \left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h} \right), & 1 \leq i \leq n, \\ [F'(u)]_{i,i-1} &= -\frac{1}{2h} f_3 \left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h} \right), & 2 \leq i \leq n, \\ [F'(u)]_{i,i+1} &= \frac{1}{2h} f_3 \left(t_i, u_i, \frac{u_{i+1} - u_{i-1}}{2h} \right), & 1 \leq i \leq n-1, \end{aligned}$$

where all other entries of $F'(u)$ are 0 and $f_2(t, u, v)$, $f_3(t, u, v)$ denote the partial derivatives of f with respect to u and v respectively.

Boundary Value Problems: Finite Difference Method

Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} =$$

$$[F'(u)]_{i,i-1} =$$

$$[F'(u)]_{i,i+1} =$$

and

$$g =$$

Boundary Value Problems: Finite Difference Method

Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left(\frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} =$$

$$[F'(u)]_{i,i+1} =$$

and

$$g =$$

Boundary Value Problems: Finite Difference Method

Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left(\frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} = -\frac{1}{2h} \left(4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

$$[F'(u)]_{i,i+1} =$$

and

$$g =$$

Boundary Value Problems: Finite Difference Method

Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left(\frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} = -\frac{1}{2h} \left(4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

$$[F'(u)]_{i,i+1} = \frac{1}{2h} \left(4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

and

$$g =$$

Boundary Value Problems: Finite Difference Method

Example

Consider the two-point BVP

$$u'' = -u + \frac{2(u')^2}{u}, \quad -1 < t < 1,$$

$$u(-1) = u(1) = (e + e^{-1})^{-1}.$$

The iterative solution via Newton's method satisfies

$$u^{(m+1)} = u^{(m)} - \left[\frac{1}{h^2} A - F'(u^{(m)}) \right]^{-1} \left[\frac{1}{h^2} A u^{(m)} - F(u^{(m)}) - g \right],$$

where

$$[F'(u)]_{ii} = -1 - 2 \left(\frac{u_{i+1} - u_{i-1}}{2hu_i} \right)^2, \quad [F'(u)]_{i,i-1} = -\frac{1}{2h} \left(4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

$$[F'(u)]_{i,i+1} = \frac{1}{2h} \left(4 \frac{u_{i+1} - u_{i-1}}{2hu_i} \right),$$

and

$$g = \frac{-1}{h^2} \begin{bmatrix} (e + e^{-1})^{-1} \\ 0 \\ \vdots \\ 0 \\ (e + e^{-1})^{-1} \end{bmatrix}.$$

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.1 Well-posedness

3.2 Shooting Method

3.3 Finite Difference Method

- Error Analysis



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Finite Difference Method



The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.



Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i)$$

Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Now, if $u \in C^4([a, b])$, then

$$u(t_i+h) = u(t_i) + hu'(t_i) + \frac{h^2}{2}u''(t_i) + \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_1), \quad \xi_1 \in (t_i, t_i+h),$$

$$u(t_i-h) = u(t_i) - hu'(t_i) + \frac{h^2}{2}u''(t_i) - \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_2), \quad \xi_2 \in (t_i-h, t_i).$$

Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Now, if $u \in C^4([a, b])$, then

$$u(t_i+h) = u(t_i) + hu'(t_i) + \frac{h^2}{2}u''(t_i) + \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_1), \quad \xi_1 \in (t_i, t_i+h),$$

$$u(t_i-h) = u(t_i) - hu'(t_i) + \frac{h^2}{2}u''(t_i) - \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_2), \quad \xi_2 \in (t_i-h, t_i).$$

Thus,

$$\ell(t_i, u) = \frac{h^2}{24} \left(u^{(4)}(\xi_1) + u^{(4)}(\xi_2) \right)$$

Boundary Value Problems: Finite Difference Method

The error analysis in full generality is too complicated to be discussed in this course. We will, therefore, look at simpler situation where the right hand side is independent of u and u' , that is, $f(t, u, u') = f(t)$.

The model BVP, therefore, reads

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta. \end{aligned}$$

We define

$$D_h^2 v(t) = \frac{v(t+h) - 2v(t) + v(t-h)}{h^2}.$$

Then, the local (truncation) error

$$\ell(t_i, u) = D_h^2 u(t_i) - f(t_i) = \frac{u(t_i+h) - 2u(t_i) + u(t_i-h)}{h^2} - u''(t_i)$$

Now, if $u \in C^4([a, b])$, then

$$u(t_i+h) = u(t_i) + hu'(t_i) + \frac{h^2}{2}u''(t_i) + \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_1), \quad \xi_1 \in (t_i, t_i+h),$$

$$u(t_i-h) = u(t_i) - hu'(t_i) + \frac{h^2}{2}u''(t_i) - \frac{h^3}{6}u'''(t_i) + \frac{h^4}{24}u^{(4)}(\xi_2), \quad \xi_2 \in (t_i-h, t_i).$$

Thus,

$$\ell(t_i, u) = \frac{h^2}{24} \left(u^{(4)}(\xi_1) + u^{(4)}(\xi_2) \right) = \frac{h^2}{12} u^{(4)}(\xi), \quad \xi \in (t_i-h, t_i+h).$$

Boundary Value Problems: Finite Difference Method

Theorem

If $v \in C^2([a, b])$, then

$$\lim_{h \rightarrow 0} \|D_h^2 v - v''\|_{\infty, h} = 0.$$

If $v \in C^4([a, b])$, then

$$\|D_h^2 v - v''\|_{\infty, h} \leq \frac{h^2}{12} \|v^{(4)}\|_{\infty, [a, b]}.$$

Boundary Value Problems: Finite Difference Method

Theorem

If $v \in C^2([a, b])$, then

$$\lim_{h \rightarrow 0} \|D_h^2 v - v''\|_{\infty, h} = 0.$$

If $v \in C^4([a, b])$, then

$$\|D_h^2 v - v''\|_{\infty, h} \leq \frac{h^2}{12} \|v^{(4)}\|_{\infty, [a, b]}.$$

Proof:

We have already seen the proof for the second part. For the first part, we use

$$v(t_i + h) = v(t_i) + hv'(t_i) + \frac{h^2}{2} v''(\xi_1), \quad \xi_1 \in (t_i, t_i + h),$$

$$v(t_i - h) = v(t_i) - hv'(t_i) + \frac{h^2}{2} v''(\xi_2), \quad \xi_2 \in (t_i - h, t_i),$$

yielding

$$D_h^2 v(t_i) - v''(t_i) = \frac{v''(\xi_1) + v''(\xi_2)}{2} - v''(t_i) = v''(\xi) - v''(t_i), \quad \xi \in (t_i - h, t_i + h).$$

The result follows!



Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.



Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} .



Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0})$$



Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1})$$

Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since $v(t_i) \leq v(t_{i_0})$.

Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since $v(t_i) \leq v(t_{i_0})$. Thus, equality holds throughout and v achieves its maximum at t_{i_0-1} and t_{i_0+1} as well.

Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since $v(t_i) \leq v(t_{i_0})$. Thus, equality holds throughout and v achieves its maximum at t_{i_0-1} and t_{i_0+1} as well. Applying the same arguments again to t_{i_0-1} and t_{i_0+1} (if in the interior), and then to their neighbors, etc., we conclude that v is constant.

Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since $v(t_i) \leq v(t_{i_0})$. Thus, equality holds throughout and v achieves its maximum at t_{i_0-1} and t_{i_0+1} as well. Applying the same arguments again to t_{i_0-1} and t_{i_0+1} (if in the interior), and then to their neighbors, etc., we conclude that v is constant.

Remark

The analogous discrete minimum principle, obtained by reversing the inequalities and replacing max by min holds.

Boundary Value Problems: Finite Difference Method

Theorem (Discrete Maximum Principle)

Let v be a function on $[a, b]$ satisfying $D_h^2 v \geq 0$ on $t_i, i = 1, \dots, n$. Then $\max_{1 \leq i \leq n} v(t_i) \leq \max\{v(t_0), v(t_{n+1})\}$. Equality holds if and only if v is constant.

Proof:

Suppose $\max_{1 \leq i \leq n} v(t_i) \geq \max\{v(t_0), v(t_{n+1})\}$ and the max is achieved at t_{i_0} . Then,

$$2v(t_{i_0}) = v(t_{i_0+1}) + v(t_{i_0-1}) - h^2 D_h^2 v(t_{i_0}) \leq v(t_{i_0+1}) + v(t_{i_0-1}) \leq 2v(t_{i_0})$$

since $v(t_i) \leq v(t_{i_0})$. Thus, equality holds throughout and v achieves its maximum at t_{i_0-1} and t_{i_0+1} as well. Applying the same arguments again to t_{i_0-1} and t_{i_0+1} (if in the interior), and then to their neighbors, etc., we conclude that v is constant.

Remark

The analogous discrete minimum principle, obtained by reversing the inequalities and replacing max by min holds.

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Boundary Value Problems: Finite Difference Method

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Proof:

It is sufficient to show that, if $D_h^2 u_h(t_i) = 0$ for $t_i, i = 1, \dots, n$, and $u_h(a) = 0$, $u_h(b) = 0$, then $u_h \equiv 0$.



Boundary Value Problems: Finite Difference Method

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Proof:

It is sufficient to show that, if $D_h^2 u_h(t_i) = 0$ for $t_i, i = 1, \dots, n$, and $u_h(a) = 0$, $u_h(b) = 0$, then $u_h \equiv 0$. But this follows from the discrete maximum principle and discrete minimum principle.

Boundary Value Problems: Finite Difference Method

Theorem

There is a unique solution to the discrete BVP

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta. \end{aligned}$$

Proof:

It is sufficient to show that, if $D_h^2 u_h(t_i) = 0$ for $t_i, i = 1, \dots, n$, and $u_h(a) = 0$, $u_h(b) = 0$, then $u_h \equiv 0$. But this follows from the discrete maximum principle and discrete minimum principle.

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

$$\text{Let } w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

therefore,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \max_{1 \leq i \leq n} (u_h(t_i) + w(t_i)\|f\|_{\infty, h}) \leq \max\{u_h(a) + w(a)\|f\|_{\infty, h}, (u_h(b) + w(b)\|f\|_{\infty, h})\}.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

therefore,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \max_{1 \leq i \leq n} (u_h(t_i) + w(t_i)\|f\|_{\infty, h}) \leq \max\{u_h(a) + w(a)\|f\|_{\infty, h}, (u_h(b) + w(b)\|f\|_{\infty, h})\}.$$

Hence,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{\alpha, \beta\}.$$

Boundary Value Problems: Finite Difference Method

Theorem

The solution u_h to

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), \quad t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, \quad u_h(b) = \beta, \end{aligned}$$

satisfies

$$\|u_h\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{|\alpha|, |\beta|\}.$$

Proof:

Let $w(t) = \frac{1}{2} \left(t - \frac{a+b}{2} \right)^2$. Then, $D_h^2 w(t_i) = w''(t_i) = 1, i = 1, \dots, n$, and $0 \leq w(t) \leq \frac{(b-a)^2}{8}, t \in [a, b]$.

Now,

$$D_h^2(u_h + w\|f\|_{\infty, h}) = D_h^2 u_h + \|f\|_{\infty, h} \geq 0,$$

therefore,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \max_{1 \leq i \leq n} (u_h(t_i) + w(t_i)\|f\|_{\infty, h}) \leq \max\{u_h(a) + w(a)\|f\|_{\infty, h}, (u_h(b) + w(b)\|f\|_{\infty, h})\}.$$

Hence,

$$\max_{1 \leq i \leq n} u_h(t_i) \leq \frac{(b-a)^2}{8} \|f\|_{\infty, h} + \max\{\alpha, \beta\}.$$

A similar argument applies to $-u_h$ giving the theorem.

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Proof:

The error estimate follows from the previous theorem applied to the discrete problem

$$\begin{aligned} D_h^2(u_h - u)(t_i) &= D_h^2 u_h(t_i) - D_h^2 u(t_i) = f(t_i) - D_h^2 u(t_i), & i = 1, \dots, n, \\ (u_h - u)(a) &= 0, & (u_h - u)(b) = 0. \end{aligned}$$

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Proof:

The error estimate follows from the previous theorem applied to the discrete problem

$$\begin{aligned} D_h^2(u_h - u)(t_i) &= D_h^2 u_h(t_i) - D_h^2 u(t_i) = f(t_i) - D_h^2 u(t_i), & i = 1, \dots, n, \\ (u_h - u)(a) &= 0, & (u_h - u)(b) = 0. \end{aligned}$$

Corollary

If $u \in C^2([a, b])$, then

$$\lim_{h \rightarrow 0} \|u_h - u\|_{\infty, h} = 0.$$

Boundary Value Problems: Finite Difference Method

Theorem

Let u be the solution to

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= \alpha, & u(b) = \beta, \end{aligned}$$

and u_h be the solution to the corresponding discrete problem

$$\begin{aligned} D_h^2 u_h(t_i) &= f(t_i), & t_i, i = 1, \dots, n, \\ u_h(a) &= \alpha, & u_h(b) = \beta. \end{aligned}$$

Then,

$$\|u_h - u\|_{\infty, h} \leq \frac{(b-a)^2}{8} \|f - D_h^2 u\|_{\infty, h} = \frac{(b-a)^2}{8} \|u'' - D_h^2 u\|_{\infty, h}.$$

Corollary

If $u \in C^2([a, b])$, then

$$\lim_{h \rightarrow 0} \|u_h - u\|_{\infty, h} = 0.$$

If $u \in C^4([a, b])$, then

$$\|u_h - u\|_{\infty, h} \leq \frac{h^2(b-a)^2}{96} \|u^{(4)}\|_{\infty, [a, b]}.$$



Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

An estimate of the form $\|v\| \leq C_h \|f\|$ whenever $D_h^2 v(t_i) = f(t_i), i = 1, \dots, n$, and $v(a) = 0, v(b) = 0$, is a **stability estimate** and if it holds with C_h independent of h , we say that the discretization is **stable**.

Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

An estimate of the form $\|v\| \leq C_h \|f\|$ whenever $D_h^2 v(t_i) = f(t_i), i = 1, \dots, n$, and $v(a) = 0, v(b) = 0$, is a **stability estimate** and if it holds with C_h independent of h , we say that the discretization is **stable**.

The previous result can be summarized as “**consistency + stability implies convergence**”.



Boundary Value Problems: Finite Difference Method

Remark

The quantity $\|f - D_h^2 u\| = \|u'' - D_h^2 u\|$ is the **consistency error** of the discretization and the statement $\lim_{h \rightarrow 0} \|u'' - D_h^2 u\| = 0$ means that the discretization is **consistent**.

An estimate of the form $\|v\| \leq C_h \|f\|$ whenever $D_h^2 v(t_i) = f(t_i)$, $i = 1, \dots, n$, and $v(a) = 0$, $v(b) = 0$, is a **stability estimate** and if it holds with C_h independent of h , we say that the discretization is **stable**.

The previous result can be summarized as “**consistency + stability implies convergence**”.

As a final remark, the finite difference method helped us find the solution values at the mesh points, but the solution at non-mesh points are not readily available from the method. If needed, one can obtain the solution at non-mesh points through interpolation or try other approximation approaches ...