

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- *Finite Difference Method*
- **Finite Element Method**



Numerical Methods for PDE: 2nd Order Elliptic PDE



Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.



Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$

Conversely, if u satisfies the second equation for all integrable functions v , then u satisfies the Poisson's equation. In fact, it is sufficient that the equation is satisfied for all C^∞ functions with compact support inside Ω .

Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$

Conversely, if u satisfies the second equation for all integrable functions v , then u satisfies the Poisson's equation. In fact, it is sufficient that the equation is satisfied for all C^∞ functions with compact support inside Ω .

In particular, if v is a C^1 function on Ω which vanishes on Γ , we have

$$-\int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f \, v.$$

This is the weak formulation of the boundary value problem.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Now, let's try to solve the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

using the finite element method where Ω is a bounded domain.

Assume that f is continuous on $\overline{\Omega}$. Then, solution u also satisfies,

$$\int_{\Omega} \Delta u \, v = \int_{\Omega} f \, v$$

Conversely, if u satisfies the second equation for all integrable functions v , then u satisfies the Poisson's equation. In fact, it is sufficient that the equation is satisfied for all C^∞ functions with compact support inside Ω .

In particular, if v is a C^1 function on Ω which vanishes on Γ , we have

$$-\int_{\Omega} \nabla u \cdot \nabla v = \int_{\Omega} f \, v.$$

This is the weak formulation of the boundary value problem.

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space.



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w)$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w)$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

To introduce the abstract framework, let V be a Hilbert space (complete inner product space), $B: V \times V \rightarrow \mathbb{R}$ be a bilinear form, and $F: V \rightarrow \mathbb{R}$ be a linear functional. Then, the weak formulation of a boundary value problem reads: find $u \in V$ such that

$$B(u, v) = F(v) \text{ for all } v \in V.$$

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

that is

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\|$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\| \leq (1 + C/\gamma) \|u - w\| \text{ for all } w \in V_h.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h,$$

so if $w \in V_h$ is an arbitrary element of V_h , then

$$B(u_h - w, v) = B(u - w, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\| \leq (1 + C/\gamma) \|u - w\| \text{ for all } w \in V_h. \text{ Therefore,}$$

$$\|u_h - u\| \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

In the Galerkin method, we choose a finite dimensional subspace V_h of V as an approximation space and same space as the test space. The Galerkin solution is defined as $u_h \in V_h$ such that

$$B(u_h, v) = F(v) \text{ for all } v \in V_h.$$

We then have

$$B(u_h, v) = B(u, v) \text{ for all } v \in V_h.$$

If the bilinear form B satisfies

$$|B(u, v)| \leq C \|u\| \|v\|, \quad B(v, v) \geq \gamma \|v\|^2$$

then

$$\gamma \|u_h - w\|^2 \leq B(u_h - w, u_h - w) = B(u - w, u_h - w) \leq C \|u - w\| \|u_h - w\|$$

then

$$\|u_h - w\| \leq (C/\gamma) \|u - w\|$$

Thus,

$$\|u_h - u\| \leq \|u_h - w\| + \|u - w\| \leq (1 + C/\gamma) \|u - w\| \text{ for all } w \in V_h. \text{ Therefore,}$$

$$\|u_h - u\| \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|.$$

We, therefore, see that the Galerkin approximation error is bounded by a constant multiple of the best approximation error for u by functions in V_h !

Lesson 4

Numerical Solution of PDE

4.1 BVP for 2nd Order Elliptic PDE

- *Finite Difference Method*
- **Finite Element Method**
 - **Construction of FEM Approximation Spaces**



Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

Numerical Methods for PDE: 2nd Order Elliptic PDE




Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.




Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.




Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.





Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.





Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.



Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure.



Numerical Methods for PDE: 2nd Order Elliptic PDE



Back to the Poisson's equation with homogeneous boundary condition



$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) =$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition



$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$

How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$



Numerical Methods for PDE: 2nd Order Elliptic PDE

Back to the Poisson's equation with homogeneous boundary condition

$$\begin{aligned}\Delta u &= f, & \text{in } \Omega, \\ u &= 0, & \text{on } \Gamma.\end{aligned}$$



How do we construct a finite element (that is, locally supported) basis for V_h ?

Recall, that in one dimensions, we partitioned Ω (an interval) into subintervals and constructed a basis for piecewise linear functions with respect to the partition.

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$

The functions φ_{mn} form a basis for subspace of piecewise linear functions with respect to the given partition/triangulation.




Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$



The functions φ_{mn} form a basis for subspace of piecewise linear functions with respect to the given partition/triangulation.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$


Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

For simplicity, we take $\Omega = (0,1) \times (0,1)$ and the triangulation shown in the figure. Then,

$$\varphi_{mn}(x_1, x_2) = \begin{cases} 1 - (x_1 - mh)/h \\ 1 - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h - (x_2 - nh)/h \\ 1 + (x_1 - mh)/h \\ 1 + (x_2 - nh)/h \\ 1 - (x_1 - mh)/h + (x_2 - nh)/h \end{cases}$$

The functions φ_{mn} form a basis for subspace of piecewise linear functions with respect to the given partition/triangulation.



Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

We see that matrix on the left hand side of the linear matrix (called stiffness matrix) for the piecewise linear finite elements for the Laplace operator on the unit square using a uniform mesh is exactly the matrix of the 5-point Laplacian.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

We see that matrix on the left hand side of the linear matrix (called stiffness matrix) for the piecewise linear finite elements for the Laplace operator on the unit square using a uniform mesh is exactly the matrix of the 5-point Laplacian.

Note that (*exercise*)

$$\int_{\Omega} \varphi_{mn} = h^2$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N.$$

We see that matrix on the left hand side of the linear matrix (called stiffness matrix) for the piecewise linear finite elements for the Laplace operator on the unit square using a uniform mesh is exactly the matrix of the 5-point Laplacian.

Note that (*exercise*)

$$\int_{\Omega} \varphi_{mn} = h^2$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N,$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Therefore, at vertices, we see that the finite element method converges with order 2 (*why?*).

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N,$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Therefore, at vertices, we see that the finite element method converges with order 2 (*why?*).

Recall that, from the general error analysis, we have

$$\|u_h - u\|_{H^1} \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|_{H^1}.$$

Numerical Methods for PDE: 2nd Order Elliptic PDE

Note that (*exercise*)

$$\int_{\Omega} \nabla \varphi_{mn} \cdot \nabla \varphi_{kl} = \begin{cases} 4, & m = k, n = l, \\ -1, & m = k \pm 1, n = l \text{ or } m = k, n = l \pm 1, \\ 0, & \text{otherwise.} \end{cases}$$

Thus, for $u_h = \sum u_{mn} \varphi_{mn}$, the linear system reads

$$\frac{u_{m-1,n} + u_{m+1,n} + u_{m,n-1} + u_{m,n+1} - 4u_{mn}}{h^2} = \frac{1}{h^2} \int_{\Omega} f \varphi_{mn} = \tilde{f}_{mn}, \quad 1 \leq m, n \leq N,$$

and if $f \in C^2$, then (*exercise*)

$$\tilde{f}_{mn} = \frac{1}{h^2} \int_{\Omega} \left(f(mh, nh) + f_{x_1}(mh, nh)(x_1 - mh) + f_{x_2}(mh, nh)(x_2 - nh) + O(h^2) \right) \varphi_{mn} = f_{mn} + O(h^2)$$

Therefore, at vertices, we see that the finite element method converges with order 2 (*why?*).

Recall that, from the general error analysis, we have

$$\|u_h - u\|_{H^1} \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|_{H^1}.$$

Exercise: Find C and γ for the piecewise linear finite elements.

Numerical Methods for PDE: 2nd Order Elliptic PDE

Recall that, from the general error analysis, we have

$$\|u_h - u\|_{H^1} \leq (1 + C/\gamma) \inf_{w \in V_h} \|u - w\|_{H^1}.$$

Exercise: Find C and γ for the piecewise linear finite elements.

From the approximation theory, we have the following result on the best approximation error.

Theorem

Let there be given a family of triangulations $\{\mathcal{T}_h\}$ of a polygonal domain Ω and let $h = \max_{T \in \mathcal{T}_h} \text{diam}(T)$. Let r be a positive integer. For each h let $P_h: C(\Omega) \rightarrow M_0^r(\mathcal{T}_h)$ denote the nodal interpolant, where $M_0^r(\mathcal{T}_h)$ is the space of continuous functions which restrict to polynomials of degree at most r when restricted to any triangle $T \in \mathcal{T}_h$. Then, there is a constant c such that

$$\begin{aligned} \|u - P_h u\|_{L^\infty(\Omega)} &\leq ch^{r+1} \|u^{(r+1)}\|_{L^\infty(\Omega)}, & u \in C^{r+1}(\overline{\Omega}), \\ \|u - P_h u\|_{L^2(\Omega)} &\leq ch^{r+1} \|u^{(r+1)}\|_{L^2(\Omega)}, & u \in H^{r+1}(\Omega). \end{aligned}$$

Moreover, if the family of triangulations are shape regular (the minimal angle of each triangulation is bounded below uniformly), then there is a constant C such that

$$\begin{aligned} \|\nabla(u - P_h u)\|_{L^\infty(\Omega)} &\leq Ch^r \|u^{(r+1)}\|_{L^\infty(\Omega)}, & u \in C^{r+1}(\overline{\Omega}), \\ \|\nabla(u - P_h u)\|_{L^2(\Omega)} &\leq Ch^r \|u^{(r+1)}\|_{L^2(\Omega)}, & u \in H^{r+1}(\Omega). \end{aligned}$$

Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.2 Parabolic PDE



Numerical Methods for PDE: Parabolic PDE



As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

Numerical Methods for PDE: Parabolic PDE



As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

To obtain a well-posed problem, we need to give boundary conditions

$$u = 0, \quad \mathbf{x} \in \Gamma, t \in [0, T]$$

and an initial condition

$$u = u_0, \quad \mathbf{x} \in \Omega, t = 0.$$

Numerical Methods for PDE: Parabolic PDE

As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

To obtain a well-posed problem, we need to give boundary conditions

$$u = 0, \quad \mathbf{x} \in \Gamma, t \in [0, T]$$

and an initial condition

$$u = u_0, \quad \mathbf{x} \in \Omega, t = 0.$$

Let's assume, for simplicity, that $\Omega = (0,1) \times (0,1)$.

Numerical Analysis & Scientific Computing II

Lesson 4

Numerical Solution of PDE

4.2 Parabolic PDE

- Semi-discretization



Akash Anand
MATH, IIT KANPUR

Numerical Methods for PDE: Parabolic PDE

As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

To obtain a well-posed problem, we need to give boundary conditions

$$u = 0, \quad \mathbf{x} \in \Gamma, t \in [0, T]$$

and an initial condition

$$u = u_0, \quad \mathbf{x} \in \Omega, t = 0.$$

Let's assume, for simplicity, that $\Omega = (0,1) \times (0,1)$. Then, we have the 5-point Laplacian Δ_h as the discretization of the Laplacian Δ . Thus, we the solution u_h satisfying

$$\begin{aligned} \frac{\partial u_h}{\partial t} &= c\Delta_h u_h + f, & \mathbf{x} \in \Omega_h, t \in [0, T] \\ u_h &= 0, & \mathbf{x} \in \Gamma_h, t \in [0, T] \\ u_h &= u_0, & \mathbf{x} \in \Omega_h, t = 0. \end{aligned}$$

Numerical Methods for PDE: Parabolic PDE

As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

To obtain a well-posed problem, we need to give boundary conditions

$$u = 0, \quad \mathbf{x} \in \Gamma, t \in [0, T]$$

and an initial condition

$$u = u_0, \quad \mathbf{x} \in \Omega, t = 0.$$

Let's assume, for simplicity, that $\Omega = (0,1) \times (0,1)$. Then, we have the 5-point Laplacian Δ_h as the discretization of the Laplacian Δ . Thus, we the solution u_h satisfying

$$\begin{aligned} \frac{\partial u_h}{\partial t} &= c\Delta_h u_h + f, & \mathbf{x} \in \Omega_h, t \in [0, T] \\ u_h &= 0, & \mathbf{x} \in \Gamma_h, t \in [0, T] \\ u_h &= u_0, & \mathbf{x} \in \Omega_h, t = 0. \end{aligned}$$

We may view this as an initial value problem for a system of $(N - 1)^2$ ODEs.

Numerical Methods for PDE: Parabolic PDE

As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

To obtain a well-posed problem, we need to give boundary conditions

$$u = 0, \quad \mathbf{x} \in \Gamma, t \in [0, T]$$

and an initial condition

$$u = u_0, \quad \mathbf{x} \in \Omega, t = 0.$$

Let's assume, for simplicity, that $\Omega = (0,1) \times (0,1)$. Then, we have the 5-point Laplacian Δ_h as the discretization of the Laplacian Δ . Thus, we the solution u_h satisfying

$$\begin{aligned} \frac{\partial u_h}{\partial t} &= c\Delta_h u_h + f, & \mathbf{x} \in \Omega_h, t \in [0, T] \\ u_h &= 0, & \mathbf{x} \in \Gamma_h, t \in [0, T] \\ u_h &= u_0, & \mathbf{x} \in \Omega_h, t = 0. \end{aligned}$$

We may view this as an initial value problem for a system of $(N - 1)^2$ ODEs.

The process of reducing the evolutionary PDE to a system of ODEs by using a finite difference approximation of the spatial operator is called **semi-discretization** or the **method of lines**.

Numerical Methods for PDE: Parabolic PDE

As a model problem, consider the heat equation on a spatial domain Ω for a time interval $[0, T]$. The solution u satisfies

$$\frac{\partial u}{\partial t} = c\Delta u + f, \quad \mathbf{x} \in \Omega, t \in [0, T], \quad c > 0.$$

To obtain a well-posed problem, we need to give boundary conditions

$$u = 0, \quad \mathbf{x} \in \Gamma, t \in [0, T]$$

and an initial condition

$$u = u_0, \quad \mathbf{x} \in \Omega, t = 0.$$

Let's assume, for simplicity, that $\Omega = (0,1) \times (0,1)$. Then, we have the 5-point Laplacian Δ_h as the discretization of the Laplacian Δ . Thus, we the solution u_h satisfying

$$\begin{aligned} \frac{\partial u_h}{\partial t} &= c\Delta_h u_h + f, & \mathbf{x} \in \Omega_h, t \in [0, T] \\ u_h &= 0, & \mathbf{x} \in \Gamma_h, t \in [0, T] \\ u_h &= u_0, & \mathbf{x} \in \Omega_h, t = 0. \end{aligned}$$

We may view this as an initial value problem for a system of $(N - 1)^2$ ODEs.

The process of reducing the evolutionary PDE to a system of ODEs by using a finite difference approximation of the spatial operator is called **semi-discretization** or the **method of lines**.

This is not a full discretization as we still have to choose a numerical method to solve the ODEs.

Lesson 4

Numerical Solution of PDE

4.2 Parabolic PDE

- *Semi-discretization*
- ***Finite difference discretization***



Numerical Methods for PDE: Parabolic PDE



To further simplify the presentation, let's drop down to one space dimension, that is, take $\Omega = (0,1)$.

Numerical Methods for PDE: Parabolic PDE



To further simplify the presentation, let's drop down to one space dimension, that is, take $\Omega = (0,1)$.

To solve the ODEs, we first employ the Euler method.



Numerical Methods for PDE: Parabolic PDE

To further simplify the presentation, let's drop down to one space dimension, that is, take $\Omega = (0,1)$.

To solve the ODEs, we first employ the Euler method.

Writing $u_n^j = u_h(nh, jk)$, the fully discrete system reads

$$\begin{aligned} \frac{u_n^{j+1} - u_n^j}{k} &= c \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} + f_n^j, & 0 < n < N, j = 0, 1, \dots, M-1, \\ u_0^j &= u_N^j = 0, & j = 0, 1, \dots, M-1, \\ u_n^0 &= u_0(nh), & 0 < n < N. \end{aligned}$$

Numerical Methods for PDE: Parabolic PDE

To further simplify the presentation, let's drop down to one space dimension, that is, take $\Omega = (0,1)$.

To solve the ODEs, we first employ the Euler method.

Writing $u_n^j = u_h(nh, jk)$, the fully discrete system reads

$$\begin{aligned} \frac{u_n^{j+1} - u_n^j}{k} &= c \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} + f_n^j, & 0 < n < N, j = 0, 1, \dots, M-1, \\ u_0^j &= u_N^j = 0, & j = 0, 1, \dots, M-1, \\ u_n^0 &= u_0(nh), & 0 < n < N. \end{aligned}$$

We call this the **forward-centered difference method** for the heat equation. Since the Euler's method is explicit, we don't need to solve a linear system:

$$u_n^{j+1} = (1 - 2\lambda)u_n^j + \lambda u_{n+1}^j + \lambda u_{n-1}^j + k f_n^j, \quad j = 0, 1, \dots, M-1,$$

where $\lambda = ck/h^2$.

Numerical Methods for PDE: Parabolic PDE

To further simplify the presentation, let's drop down to one space dimension, that is, take $\Omega = (0,1)$.

To solve the ODEs, we first employ the Euler method.

Writing $u_n^j = u_h(nh, jk)$, the fully discrete system reads

$$\begin{aligned} \frac{u_n^{j+1} - u_n^j}{k} &= c \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} + f_n^j, & 0 < n < N, j = 0, 1, \dots, M-1, \\ u_0^j &= u_N^j = 0, & j = 0, 1, \dots, M-1, \\ u_n^0 &= u_0(nh), & 0 < n < N. \end{aligned}$$

We call this the **forward-centered difference method** for the heat equation. Since the Euler's method is explicit, we don't need to solve a linear system:

$$u_n^{j+1} = (1 - 2\lambda)u_n^j + \lambda u_{n+1}^j + \lambda u_{n-1}^j + k f_n^j, \quad j = 0, 1, \dots, M-1,$$

where $\lambda = ck/h^2$.

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

Numerical Methods for PDE: Parabolic PDE

To further simplify the presentation, let's drop down to one space dimension, that is, take $\Omega = (0,1)$.

To solve the ODEs, we first employ the Euler method.

Writing $u_n^j = u_h(nh, jk)$, the fully discrete system reads

$$\begin{aligned} \frac{u_n^{j+1} - u_n^j}{k} &= c \frac{u_{n+1}^j - 2u_n^j + u_{n-1}^j}{h^2} + f_n^j, & 0 < n < N, j = 0, 1, \dots, M-1, \\ u_0^j &= u_N^j = 0, & j = 0, 1, \dots, M-1, \\ u_n^0 &= u_0(nh), & 0 < n < N. \end{aligned}$$

We call this the **forward-centered difference method** for the heat equation. Since the Euler's method is explicit, we don't need to solve a linear system:

$$u_n^{j+1} = (1 - 2\lambda)u_n^j + \lambda u_{n+1}^j + \lambda u_{n-1}^j + k f_n^j, \quad j = 0, 1, \dots, M-1,$$

where $\lambda = ck/h^2$.

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

By Taylor's theorem,

$$\ell_n^j = \frac{k}{2} \frac{\partial^2 u}{\partial t^2} - c \frac{h^2}{12} \frac{\partial^4 u}{\partial x^4}$$

where derivatives are evaluated at appropriate points.

Numerical Methods for PDE: Parabolic PDE

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

By Taylor's theorem,

$$\ell_n^j = \frac{k}{2} \frac{\partial^2 u}{\partial t^2} - c \frac{h^2}{12} \frac{\partial^4 u}{\partial x^4}$$

where derivatives are evaluated at appropriate points.

Let $e_n^j = u_n^j - u(nh, jk)$. Then,

$$e_n^{j+1} = (1 - 2\lambda)e_n^j + \lambda e_{n+1}^j + \lambda e_{n-1}^j - k\ell_n^j.$$

Numerical Methods for PDE: Parabolic PDE

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

By Taylor's theorem,

$$\ell_n^j = \frac{k}{2} \frac{\partial^2 u}{\partial t^2} - c \frac{h^2}{12} \frac{\partial^4 u}{\partial x^4}$$

where derivatives are evaluated at appropriate points.

Let $e_n^j = u_n^j - u(nh, jk)$. Then,

$$e_n^{j+1} = (1 - 2\lambda)e_n^j + \lambda e_{n+1}^j + \lambda e_{n-1}^j - k\ell_n^j.$$

If we let $E^j = \max_n |e_n^j|$ denote the maximum norm of the error at the j th time step, we get

$$E^{j+1} \leq |1 - 2\lambda|E^j + \lambda E^j + \lambda E^j + k \max_n |\ell_n^j|.$$

Numerical Methods for PDE: Parabolic PDE

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

By Taylor's theorem,

$$\ell_n^j = \frac{k}{2} \frac{\partial^2 u}{\partial t^2} - c \frac{h^2}{12} \frac{\partial^4 u}{\partial x^4}$$

where derivatives are evaluated at appropriate points.

Let $e_n^j = u_n^j - u(nh, jk)$. Then,

$$e_n^{j+1} = (1 - 2\lambda)e_n^j + \lambda e_{n+1}^j + \lambda e_{n-1}^j - k\ell_n^j.$$

If we let $E^j = \max_n |e_n^j|$ denote the maximum norm of the error at the j th time step, we get

$$E^{j+1} \leq |1 - 2\lambda|E^j + \lambda E^j + \lambda E^j + k \max_n |\ell_n^j|.$$

If $\lambda \leq 1/2$, and $\ell = \max_{n,j} |\ell_n^j|$, we get

$$E^{j+1} \leq E^j + k\ell.$$

Numerical Methods for PDE: Parabolic PDE

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

By Taylor's theorem,

$$\ell_n^j = \frac{k}{2} \frac{\partial^2 u}{\partial t^2} - c \frac{h^2}{12} \frac{\partial^4 u}{\partial x^4}$$

where derivatives are evaluated at appropriate points.

Let $e_n^j = u_n^j - u(nh, jk)$. Then,

$$e_n^{j+1} = (1 - 2\lambda)e_n^j + \lambda e_{n+1}^j + \lambda e_{n-1}^j - k\ell_n^j.$$

If we let $E^j = \max_n |e_n^j|$ denote the maximum norm of the error at the j th time step, we get

$$E^{j+1} \leq |1 - 2\lambda|E^j + \lambda E^j + \lambda E^j + k \max_n |\ell_n^j|.$$

If $\lambda \leq 1/2$, and $\ell = \max_{n,j} |\ell_n^j|$, we get

$$E^{j+1} \leq E^j + k\ell.$$

It follows that

$$E^j \leq E^0 + jk\ell$$

for all j .

Numerical Methods for PDE: Parabolic PDE

To analyze the method, consider the local (truncation) error

$$\ell_n^j = \frac{u(nh, (j+1)k) - u(nh, jk)}{k} - c \frac{u((n+1)h, jk) - 2u(nh, jk) + u((n-1)h, jk)}{h^2} - f_n^j.$$

By Taylor's theorem,

$$\ell_n^j = \frac{k}{2} \frac{\partial^2 u}{\partial t^2} - c \frac{h^2}{12} \frac{\partial^4 u}{\partial x^4}$$

where derivatives are evaluated at appropriate points.

Let $e_n^j = u_n^j - u(nh, jk)$. Then,

$$e_n^{j+1} = (1 - 2\lambda)e_n^j + \lambda e_{n+1}^j + \lambda e_{n-1}^j - k\ell_n^j.$$

If we let $E^j = \max_n |e_n^j|$ denote the maximum norm of the error at the j th time step, we get

$$E^{j+1} \leq |1 - 2\lambda|E^j + \lambda E^j + \lambda E^j + k \max_n |\ell_n^j|.$$

If $\lambda \leq 1/2$, and $\ell = \max_{n,j} |\ell_n^j|$, we get

$$E^{j+1} \leq E^j + k\ell.$$

It follows that

$$E^j \leq E^0 + jk\ell$$

for all j . As $jk \leq T$ and $k \leq h^2/(2c)$, we have

$$\max_{n,j} |e_n^j| \leq T\ell \leq C(k + h^2) \leq C(1 + 1/(2c))h^2.$$

Lesson 4

Numerical Solution of PDE

4.2 Parabolic PDE

- *Semi-discretization*
- *Full finite difference discretization*
- ***Fourier Analysis***



Numerical Methods for PDE: Parabolic PDE



Another useful way to analyze is to use Fourier analysis.

Numerical Methods for PDE: Parabolic PDE

Another useful way to analyze is to use Fourier analysis.

Recall that, on $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$ and $\varphi_m(x) = \sin \pi m x$, $m = 1, \dots, N-1$, form an orthogonal basis.

Also,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (\cos \pi m h - 1) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2},$$

where the eigenvalues satisfy

$$8 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < \frac{4}{h^2}.$$

Numerical Methods for PDE: Parabolic PDE

Another useful way to analyze is to use Fourier analysis.

Recall that, on $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$ and $\varphi_m(x) = \sin \pi m x$, $m = 1, \dots, N-1$, form an orthogonal basis.

Also,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (\cos \pi m h - 1) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2},$$

where the eigenvalues satisfy

$$8 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < \frac{4}{h^2}.$$

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Numerical Methods for PDE: Parabolic PDE

Another useful way to analyze is to use Fourier analysis.

Recall that, on $L(I_h)$, we define the inner product

$$\langle u, v \rangle_h = h \sum_{k=1}^{N-1} u(kh)v(kh)$$

with the corresponding norm $\|v\|_h$ and $\varphi_m(x) = \sin \pi m x$, $m = 1, \dots, N-1$, form an orthogonal basis.

Also,

$$D_h^2 \varphi_m = -\lambda_m \varphi_m, \quad \lambda_m = \frac{2}{h^2} (\cos \pi m h - 1) = \frac{4}{h^2} \sin^2 \frac{\pi m h}{2},$$

where the eigenvalues satisfy

$$0 < \lambda_1 < \lambda_2 < \dots < \lambda_{N-1} < \frac{4}{h^2}.$$

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Thus,

$$\frac{da_m^h}{dt} = -c a_m^h \lambda_m \quad \text{so} \quad a_m^h(t) = a_m^h(0) e^{-c \lambda_m t}.$$

Numerical Methods for PDE: Parabolic PDE

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Thus,

$$\frac{da_m^h}{dt} = -c a_m^h \lambda_m \quad \text{so} \quad a_m^h(t) = a_m^h(0) e^{-c \lambda_m t}.$$

The numbers $a_m^h(0)$ are the coefficients in the discrete Fourier sine transform of the initial data:

$$u_0(x) = \sum_{m=1}^{N-1} a_m^h(0) \varphi_m(x), \quad x \in \bar{I}_h.$$

Numerical Methods for PDE: Parabolic PDE

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Thus,

$$\frac{da_m^h}{dt} = -c a_m^h \lambda_m \quad \text{so} \quad a_m^h(t) = a_m^h(0) e^{-c \lambda_m t}.$$

The numbers $a_m^h(0)$ are the coefficients in the discrete Fourier sine transform of the initial data:

$$u_0(x) = \sum_{m=1}^{N-1} a_m^h(0) \varphi_m(x), \quad x \in \bar{I}_h.$$

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c \lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

Numerical Methods for PDE: Parabolic PDE

We write the semi-discrete solution $u_h(t, \cdot) = \sum_{m=1}^{N-1} a_m^h \varphi_m$. Then,

$$\frac{\partial u_h}{\partial t} = \sum_{m=1}^{N-1} \frac{da_m^h}{dt} \varphi_m, \quad D_h^2 u_h = - \sum_{m=1}^{N-1} a_m^h \lambda_m \varphi_m.$$

Thus,

$$\frac{da_m^h}{dt} = -c a_m^h \lambda_m \quad \text{so} \quad a_m^h(t) = a_m^h(0) e^{-c \lambda_m t}.$$

The numbers $a_m^h(0)$ are the coefficients in the discrete Fourier sine transform of the initial data:

$$u_0(x) = \sum_{m=1}^{N-1} a_m^h(0) \varphi_m(x), \quad x \in \bar{I}_h.$$

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c \lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

Numerical Methods for PDE: Parabolic PDE

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c\lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution at time $t = jk$ as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

The difference equation then gives

$$\sum_{m=1}^{N-1} A_m^{j+1} \varphi_m(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x) - ck \sum_{m=1}^{N-1} A_m^j \lambda_m \varphi_m(x)$$

yielding

$$A_m^{j+1} = (1 - ck\lambda_m) A_m^j.$$

Numerical Methods for PDE: Parabolic PDE

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c\lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution at time $t = jk$ as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

The difference equation then gives

$$\sum_{m=1}^{N-1} A_m^{j+1} \varphi_m(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x) - ck \sum_{m=1}^{N-1} A_m^j \lambda_m \varphi_m(x)$$

yielding

$$A_m^{j+1} = (1 - ck\lambda_m) A_m^j.$$

It follows that

$$A_m^j = (1 - ck\lambda_m)^j a_m^h(0)$$

Numerical Methods for PDE: Parabolic PDE

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c\lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution at time $t = jk$ as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

The difference equation then gives

$$\sum_{m=1}^{N-1} A_m^{j+1} \varphi_m(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x) - ck \sum_{m=1}^{N-1} A_m^j \lambda_m \varphi_m(x)$$

yielding

$$A_m^{j+1} = (1 - ck\lambda_m) A_m^j.$$

It follows that

$$A_m^j = (1 - ck\lambda_m)^j a_m^h(0), \quad u^j = \sum_{m=1}^{N-1} (1 - ck\lambda_m)^j a_m^h(0) \varphi_m.$$

Numerical Methods for PDE: Parabolic PDE

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c\lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution at time $t = jk$ as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

The difference equation then gives

$$\sum_{m=1}^{N-1} A_m^{j+1} \varphi_m(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x) - ck \sum_{m=1}^{N-1} A_m^j \lambda_m \varphi_m(x)$$

yielding

$$A_m^{j+1} = (1 - ck\lambda_m) A_m^j.$$

It follows that

$$A_m^j = (1 - ck\lambda_m)^j a_m^h(0), \quad u^j = \sum_{m=1}^{N-1} (1 - ck\lambda_m)^j a_m^h(0) \varphi_m.$$

If we assume $ck/h^2 \leq 1/2$, then $ck\lambda_m \leq ck(4/h^2) \leq 2$ and hence $|1 - ck\lambda_m| \leq 1$ for all m and the solution remains bounded.

Numerical Methods for PDE: Parabolic PDE

Thus, the solution of the semi-discrete system may be written as

$$u(x, t) = \sum_{m=1}^{N-1} a_m^h(0) e^{-c\lambda_m t} \varphi_m(x), \quad x \in \bar{I}_h.$$

For the fully discrete forward-centered scheme, we write the solution at time $t = jk$ as

$$u^j(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x).$$

The difference equation then gives

$$\sum_{m=1}^{N-1} A_m^{j+1} \varphi_m(x) = \sum_{m=1}^{N-1} A_m^j \varphi_m(x) - ck \sum_{m=1}^{N-1} A_m^j \lambda_m \varphi_m(x)$$

yielding

$$A_m^{j+1} = (1 - ck\lambda_m) A_m^j.$$

It follows that

$$A_m^j = (1 - ck\lambda_m)^j a_m^h(0), \quad u^j = \sum_{m=1}^{N-1} (1 - ck\lambda_m)^j a_m^h(0) \varphi_m.$$

If we assume $ck/h^2 \leq 1/2$, then $ck\lambda_m \leq ck(4/h^2) \leq 2$ and hence $|1 - ck\lambda_m| \leq 1$ for all m and the solution remains bounded. On the other hand, if $|1 - ck\lambda_m| > 1$ for some m , the initial data will increase exponentially.

Numerical Methods for PDE: Parabolic PDE



This idea can be used for rigorous stability analysis.



Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$



Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k\|f^{j-1}\|_h$$

Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k\|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m|\right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h.$$

Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k\|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m|\right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h.$$

*Because of the condition $ck/h^2 \leq 1/2$, which we know is not only sufficient but necessary for stability, forward-centered difference method is called **conditionally stable**.*

Numerical Methods for PDE: Parabolic PDE

This idea can be used for rigorous stability analysis.

In the matrix form, we write

$$u^{j+1} = (I + ckD_h^2)u^j + kf^j, \quad j = 0, 1, \dots, M-1,$$

where $(I + ckD_h^2)$ is a symmetric operator on $L(I_h)$. Thus,

$$\|(I + ckD_h^2)v\|_h \leq (\max_m |1 - ck\lambda_m|) \|v\|_h$$

and, therefore, we have

$$\|(I + ckD_h^2)\|_h \leq \max_m |1 - ck\lambda_m|.$$

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k\|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m|\right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h.$$

Because of the condition $ck/h^2 \leq 1/2$, which we know is not only sufficient but necessary for stability, forward-centered difference method is called **conditionally stable**.

From stability, we obtain the convergence result in the same way as earlier. Let $e_n^j = u_n^j - u(nh, jk)$, we have

$$\begin{aligned} \frac{e_n^{j+1} - e_n^j}{k} &= c \frac{e_{n+1}^j - 2e_n^j + e_{n-1}^j}{h^2} - \ell_n^j, \quad 0 < n < N, j = 0, 1, \dots, M-1, \\ e_0^j &= e_N^j = 0, \quad j = 0, 1, \dots, M-1, \\ e_n^0 &= 0, \quad 0 < n < N. \end{aligned}$$

Numerical Methods for PDE: Parabolic PDE

Finally,

$$\|u^j\|_h \leq \max_m |1 - ck\lambda_m| \|u^{j-1}\|_h + k \|f^{j-1}\|_h \leq \left(\max_m |1 - ck\lambda_m| \right)^j \|u^0\|_h + Mk \max_j \|f^j\|_h$$

Because of the condition $ck/h^2 \leq 1/2$, which we know is not only sufficient but necessary for stability, forward-centered difference method is called **conditionally stable**.

From stability, we obtain the convergence result in the same way as earlier. Let $e_n^j = u_n^j - u(nh, jk)$, we have

$$\begin{aligned} \frac{e_n^{j+1} - e_n^j}{k} &= c \frac{e_{n+1}^j - 2e_n^j + e_{n-1}^j}{h^2} - \ell_n^j, & 0 < n < N, j = 0, 1, \dots, M-1, \\ e_0^j &= e_N^j = 0, & j = 0, 1, \dots, M-1, \\ e_n^0 &= 0, & 0 < n < N. \end{aligned}$$

The stability result then yields

$$\|e^j\|_h \leq \left(\max_m |1 - ck\lambda_m| \right)^j \|e^0\|_h + Mk \max_j \|\ell^j\|_h \leq T \max_j \|\ell^j\|_h.$$