

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Variational Methods



One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.



Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

We seek the solution of the form

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \varphi_i(t),$$

where $\varphi_i(t)$ are basis functions defined on $[a, b]$ and y is an n -vector of parameters to be determined.

Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

We seek the solution of the form

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \varphi_i(t),$$

where $\varphi_i(t)$ are basis functions defined on $[a, b]$ and y is an n -vector of parameters to be determined.

Some popular finite dimensional approximation spaces are

- (1) polynomials of degree n or less.*
- (2) trigonometric polynomials of degree n or less.*
- (3) piecewise polynomials of degree n or less.*

Boundary Value Problems: Variational Methods

One way to rectify the non-availability of the solution at non-mesh points is to try to approximate the solution using functions coming from a finite dimensional space.

We again consider a scalar two-point boundary value problem

$$u'' = f(t, u, u'), \quad a < t < b,$$

with boundary conditions

$$u(a) = \alpha, \quad u(b) = \beta.$$

We seek the solution of the form

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \varphi_i(t),$$

where $\varphi_i(t)$ are basis functions defined on $[a, b]$ and y is an n -vector of parameters to be determined.

Some popular finite dimensional approximation spaces are

- (1) polynomials of degree n or less.*
- (2) trigonometric polynomials of degree n or less.*
- (3) piecewise polynomials of degree n or less.*

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods

- Collocation Method



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Variational Methods



The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points.



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$

that is,

$$\sum_{j=1}^n y_j \varphi_j''(t) = f\left(t_i, \sum_{j=1}^n y_j \varphi_j(t_i), \sum_{j=1}^n y_j \varphi_j'(t_i)\right), \quad i = 2, \dots, n - 1.$$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$

that is,

$$\sum_{j=1}^n y_j \varphi_j''(t) = f\left(t_i, \sum_{j=1}^n y_j \varphi_j(t_i), \sum_{j=1}^n y_j \varphi_j'(t_i)\right), \quad i = 2, \dots, n - 1.$$

In addition, we also enforce the boundary condition at the end-points:

$$\sum_{j=1}^n y_j \varphi_j(t_1) = \alpha, \quad \sum_{j=1}^n y_j \varphi_j(t_n) = \beta.$$



Boundary Value Problems: Variational Methods

The residual $r(t, y) := v''(t, y) - f(t, v(t, y), v'(t, y))$ measures how well the approximation satisfies the ODE. For an exact approximation, that is, $u(t) = v(t, y)$, we have $r(t, y) = 0$.

The **collocation method** seeks to enforce $r(t, y) = 0$ at a finitely many points in $[a, b]$.

We introduce the mesh points $t_i = a + (i - 1)h, i = 1, \dots, n$, where $h = (b - a)/(n - 1)$ and enforce $r(t_i, y) = 0$ at the interior mesh points. Thus, we require

$$v''(t_i, y) - f(t_i, v(t_i, y), v'(t_i, y)) = 0, \quad i = 2, \dots, n - 1,$$

that is,

$$\sum_{j=1}^n y_j \varphi_j''(t) = f\left(t_i, \sum_{j=1}^n y_j \varphi_j(t_i), \sum_{j=1}^n y_j \varphi_j'(t_i)\right), \quad i = 2, \dots, n - 1.$$

In addition, we also enforce the boundary condition at the end-points:

$$\sum_{j=1}^n y_j \varphi_j(t_1) = \alpha, \quad \sum_{j=1}^n y_j \varphi_j(t_n) = \beta.$$

This yields a system of n algebraic equation in n unknowns. This may be linear or non-linear depending on whether f is linear or non-linear.





Boundary Value Problems: Variational Methods

In particular, for the approximation space consisting of polynomials of degree $n - 1$ or less (i.e., \mathcal{P}_{n-1}), we have

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \ell_i(t) \in \mathcal{P}_{n-1},$$

where the Lagrange basis, given by

$$\ell_i(t) = \frac{\prod_{k=1, k \neq i}^n (t - t_k)}{\prod_{k=1, k \neq i}^n (t_i - t_k)} \in \mathcal{P}_{n-1}$$

is used.

Boundary Value Problems: Variational Methods

In particular, for the approximation space consisting of polynomials of degree $n - 1$ or less (i.e., \mathcal{P}_{n-1}), we have

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \ell_i(t) \in \mathcal{P}_{n-1},$$

where the Lagrange basis, given by

$$\ell_i(t) = \frac{\prod_{k=1, k \neq i}^n (t - t_k)}{\prod_{k=1, k \neq i}^n (t_i - t_k)} \in \mathcal{P}_{n-1}$$

is used. Then the collocation method yields the following system of algebraic equations: $Ay = F(y)$ where

$$Ay = \begin{bmatrix} \ell_1(t_1) & \ell_2(t_1) & \cdots & \ell_n(t_1) \\ \ell_1''(t_2) & \ell_2''(t_2) & & \\ \vdots & \vdots & \ddots & \vdots \\ \ell_1(t_n) & \cdots & \ell_{n-1}''(t_{n-1}) & \ell_n''(t_{n-1}) \\ & & \ell_{n-1}(t_n) & \ell_n(t_n) \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix}, F(y) = \begin{bmatrix} f\left(t_2, \sum_{i=1}^n y_i \ell_i(t_2), \sum_{i=1}^n y_i \ell_i'(t_2)\right) \\ \vdots \\ f\left(t_{n-1}, \sum_{i=1}^n y_i \ell_i(t_{n-1}), \sum_{i=1}^n y_i \ell_i'(t_{n-1})\right) \end{bmatrix}.$$

α β

Boundary Value Problems: Variational Methods

In particular, for the approximation space consisting of polynomials of degree $n - 1$ or less (i.e., \mathcal{P}_{n-1}), we have

$$u(t) \approx v(t, y) = \sum_{i=1}^n y_i \ell_i(t) \in \mathcal{P}_{n-1},$$

where the Lagrange basis, given by

$$\ell_i(t) = \frac{\prod_{k=1, k \neq i}^n (t - t_k)}{\prod_{k=1, k \neq i}^n (t_i - t_k)} \in \mathcal{P}_{n-1}$$

is used. Then the collocation method yields the following system of algebraic equations: $Ay = F(y)$ where

$$Ay = \begin{bmatrix} \ell_1(t_1) & \ell_2(t_1) & \cdots & \ell_n(t_1) \\ \ell_1''(t_2) & \ell_2''(t_2) & & \\ \vdots & \vdots & \ddots & \vdots \\ & & \ell_{n-1}''(t_{n-1}) & \ell_n''(t_{n-1}) \\ \ell_1(t_n) & \cdots & \ell_{n-1}(t_n) & \ell_n(t_n) \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{n-1} \\ y_n \end{bmatrix}, F(y) = \begin{bmatrix} f\left(t_2, \sum_{i=1}^n y_i \ell_i(t_2), \sum_{i=1}^n y_i \ell_i'(t_2)\right) \\ \vdots \\ f\left(t_{n-1}, \sum_{i=1}^n y_i \ell_i(t_{n-1}), \sum_{i=1}^n y_i \ell_i'(t_{n-1})\right) \end{bmatrix}.$$

α β

The Newton iterations take the form $y^{(m+1)} = y^{(m)} - \left(A - F'(y^{(m)})\right)^{-1} (Ay^{(m)} - F(y^{(m)}))$.



Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$



Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$



Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2\left(t - \frac{1}{2}\right)(t - 1)y_1 - 4t(t - 1)y_2 + 2t\left(t - \frac{1}{2}\right)y_3.$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2\left(t - \frac{1}{2}\right)(t - 1)y_1 - 4t(t - 1)y_2 + 2t\left(t - \frac{1}{2}\right)y_3.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\4y_1 - 8y_2 + 4y_3 &= 6\left(\frac{1}{2}\right), \\y_3 &= 1.\end{aligned}$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2\left(t - \frac{1}{2}\right)(t - 1)y_1 - 4t(t - 1)y_2 + 2t\left(t - \frac{1}{2}\right)y_3.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\4y_1 - 8y_2 + 4y_3 &= 6\left(\frac{1}{2}\right), \\y_3 &= 1.\end{aligned}$$

Thus the approximate solution is

$$u(t) \approx -4t(t - 1)\left(\frac{1}{8}\right) + 2t\left(t - \frac{1}{2}\right)$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

We seek the approximate solution in the space of quadratic polynomials of the form

$$v(t, y) = 2\left(t - \frac{1}{2}\right)(t - 1)y_1 - 4t(t - 1)y_2 + 2t\left(t - \frac{1}{2}\right)y_3.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\4y_1 - 8y_2 + 4y_3 &= 6\left(\frac{1}{2}\right), \\y_3 &= 1.\end{aligned}$$

Thus the approximate solution is

$$u(t) \approx -4t(t - 1)\left(\frac{1}{8}\right) + 2t\left(t - \frac{1}{2}\right) = -\frac{1}{2}t + \frac{3}{2}t^2.$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

Alternately, if we use the monomial basis $\{1, t, t^2\}$ for the space of quadratic polynomials, we then have approximate solution of the form

$$v(t, y) = y_1 + y_2 t + y_3 t^2.$$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned}u'' &= 6t, & 0 < t < 1, \\u(0) &= 0, & u(1) = 1.\end{aligned}$$

Let's take a mesh with single interior collocation point, that is, the mesh with three points:

$$t_1 = 0, \quad t_2 = \frac{1}{2}, \quad t_3 = 1.$$

Alternately, if we use the monomial basis $\{1, t, t^2\}$ for the space of quadratic polynomials, we then have approximate solution of the form

$$v(t, y) = y_1 + y_2 t + y_3 t^2.$$

The system of algebraic equation that we solve is

$$\begin{aligned}y_1 &= 0, \\2y_3 &= 6 \left(\frac{1}{2}\right), \\y_1 + y_2 + y_3 &= 1.\end{aligned}$$

Thus, the approximate solution is

$$u(t) \approx -\frac{1}{2}t + \frac{3}{2}t^2.$$

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods

- Least Squares Method



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Variational Methods



Instead of making $r(t, y) = 0$ at a finitely many points, there are other ways to enforce that the residual $r(t, y)$ is small so that the numerical solution $v(t, y)$ satisfies the ODE approximately.

Boundary Value Problems: Variational Methods



Instead of making $r(t, y) = 0$ at a finitely many points, there are other ways to enforce that the residual $r(t, y)$ is small so that the numerical solution $v(t, y)$ satisfies the ODE approximately.

*In one such approach, we minimize the square of L_2 norm of the residual. This approach is called the **least squares method**.*

Boundary Value Problems: Variational Methods

Instead of making $r(t, y) = 0$ at a finitely many points, there are other ways to enforce that the residual $r(t, y)$ is small so that the numerical solution $v(t, y)$ satisfies the ODE approximately.

In one such approach, we minimize the square of L_2 norm of the residual. This approach is called the **least squares method**.

Toward this, consider the functional $F: \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$F(y) = \frac{1}{2} \|r\|^2 = \frac{1}{2} \langle r, r \rangle,$$

where

$$\langle u, v \rangle = \int_a^b u(t)v(t)dt, \quad r(t, y) = \sum_{j=1}^n y_j \varphi_j''(t) - f\left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t)\right).$$

Boundary Value Problems: Variational Methods

Instead of making $r(t, y) = 0$ at a finitely many points, there are other ways to enforce that the residual $r(t, y)$ is small so that the numerical solution $v(t, y)$ satisfies the ODE approximately.

In one such approach, we minimize the square of L_2 norm of the residual. This approach is called the **least squares method**.

Toward this, consider the functional $F: \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$F(y) = \frac{1}{2} \|r\|^2 = \frac{1}{2} \langle r, r \rangle,$$

where

$$\langle u, v \rangle = \int_a^b u(t)v(t)dt, \quad r(t, y) = \sum_{j=1}^n y_j \varphi_j''(t) - f\left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t)\right).$$

We minimize F by setting its gradient to zero, that is,

$$0 = \frac{\partial F(y)}{\partial y_i}$$

Boundary Value Problems: Variational Methods

Instead of making $r(t, y) = 0$ at a finitely many points, there are other ways to enforce that the residual $r(t, y)$ is small so that the numerical solution $v(t, y)$ satisfies the ODE approximately.

In one such approach, we minimize the square of L_2 norm of the residual. This approach is called the **least squares method**.

Toward this, consider the functional $F: \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$F(y) = \frac{1}{2} \|r\|^2 = \frac{1}{2} \langle r, r \rangle,$$

where

$$\langle u, v \rangle = \int_a^b u(t)v(t)dt, \quad r(t, y) = \sum_{j=1}^n y_j \varphi_j''(t) - f\left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t)\right).$$

We minimize F by setting its gradient to zero, that is,

$$0 = \frac{\partial F(y)}{\partial y_i} = \int_a^b r(t, y) \frac{\partial r(t, y)}{\partial y_i} dt$$

Boundary Value Problems: Variational Methods

Instead of making $r(t, y) = 0$ at a finitely many points, there are other ways to enforce that the residual $r(t, y)$ is small so that the numerical solution $v(t, y)$ satisfies the ODE approximately.

In one such approach, we minimize the square of L_2 norm of the residual. This approach is called the **least squares method**.

Toward this, consider the functional $F: \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$F(y) = \frac{1}{2} \|r\|^2 = \frac{1}{2} \langle r, r \rangle,$$

where

$$\langle u, v \rangle = \int_a^b u(t)v(t)dt, \quad r(t, y) = \sum_{j=1}^n y_j \varphi_j''(t) - f \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right).$$

We minimize F by setting its gradient to zero, that is,

$$\begin{aligned} 0 &= \frac{\partial F(y)}{\partial y_i} = \int_a^b r(t, y) \frac{\partial r(t, y)}{\partial y_i} dt \\ &= \int_a^b r(t, y) \left(\varphi_i''(t) - f_2 \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right) \varphi_i(t) - f_3 \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right) \varphi_i'(t) \right) dt. \end{aligned}$$

Boundary Value Problems: Variational Methods

$$\int_a^b r(t, y) \left(\varphi_i''(t) - f_2 \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right) \varphi_i(t) - f_3 \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right) \varphi_i'(t) \right) dt = 0.$$

This yields a system of algebraic equations

$$\begin{aligned} \sum_{j=1}^n y_j \langle \varphi_j'', \varphi_i'' \rangle &= \langle f, \varphi_i'' \rangle + \\ \sum_{j=1}^n y_j \left\langle \varphi_j'', \varphi_i f_2 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle &- \left\langle f, \varphi_i f_2 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle + \\ \sum_{j=1}^n y_j \left\langle \varphi_j'', \varphi_i' f_3 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle &- \left\langle f, \varphi_i' f_3 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle, \quad i = 1, \dots, n. \end{aligned}$$

Boundary Value Problems: Variational Methods

$$\int_a^b r(t, y) \left(\varphi_i''(t) - f_2 \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right) \varphi_i(t) - f_3 \left(t, \sum_{j=1}^n y_j \varphi_j(t), \sum_{j=1}^n y_j \varphi_j'(t) \right) \varphi_i'(t) \right) dt = 0.$$

This yields a system of algebraic equations

$$\begin{aligned} \sum_{j=1}^n y_j \langle \varphi_j'', \varphi_i'' \rangle &= \langle f, \varphi_i'' \rangle + \\ \sum_{j=1}^n y_j \left\langle \varphi_j'', \varphi_i f_2 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle &- \left\langle f, \varphi_i f_2 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle + \\ \sum_{j=1}^n y_j \left\langle \varphi_j'', \varphi_i' f_3 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle &- \left\langle f, \varphi_i' f_3 \left(\cdot, \sum_{j=1}^n y_j \varphi_j, \sum_{j=1}^n y_j \varphi_j' \right) \right\rangle, \quad i = 1, \dots, n. \end{aligned}$$

In particular, for the linear problem with $f(t, u, v) = f(t)$, we have

$$\sum_{j=1}^n y_j \langle \varphi_j'', \varphi_i'' \rangle = \langle f, \varphi_i'' \rangle,$$

a symmetric system.

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods

- Galerkin Method



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Variational Methods



More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Boundary Value Problems: Variational Methods



More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Clearly, the least squares method uses $\psi_i = \varphi_i''$.

Boundary Value Problems: Variational Methods



More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Clearly, the least squares method uses $\psi_i = \varphi_i''$.

Also, the collocation method uses $\psi_i = \delta(t - t_i)$, where t_i is the i -th collocation point and δ is the Dirac delta function.

Boundary Value Problems: Variational Methods



More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Clearly, the least squares method uses $\psi_i = \varphi_i''$.

Also, the collocation method uses $\psi_i = \delta(t - t_i)$, where t_i is the i -th collocation point and δ is the Dirac delta function.

A popular alternative is known as the **Galerkin method** where $\psi_i = \varphi_i$ is used.

Boundary Value Problems: Variational Methods

More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Clearly, the least squares method uses $\psi_i = \varphi_i''$.

Also, the collocation method uses $\psi_i = \delta(t - t_i)$, where t_i is the i -th collocation point and δ is the Dirac delta function.

A popular alternative is known as the **Galerkin method** where $\psi_i = \varphi_i$ is used. In this case, for the linear problem with $f(t, u, v) = f(t)$, the method yields the linear system with n equations in n unknowns,

$$\sum_{j=1}^n y_j \langle \varphi_j'', \varphi_i \rangle = \langle f, \varphi_i \rangle, \quad i = 1, \dots, n.$$

Boundary Value Problems: Variational Methods



More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Clearly, the least squares method uses $\psi_i = \varphi_i''$.

Also, the collocation method uses $\psi_i = \delta(t - t_i)$, where t_i is the i -th collocation point and δ is the Dirac delta function.

A popular alternative is known as the **Galerkin method** where $\psi_i = \varphi_i$ is used. In this case, for the linear problem with $f(t, u, v) = f(t)$, the method yields the linear system with n equations in n unknowns,

$$\sum_{j=1}^n y_j \langle \varphi_j'', \varphi_i \rangle = \langle f, \varphi_i \rangle, \quad i = 1, \dots, n.$$

Typically, we assume that the basis functions satisfy the homogeneous boundary conditions $\varphi_i(a) = \varphi_i(b) = 0$ so that the approximation space $\text{span} \{\varphi_i: i = 1, \dots, n\}$ is a vector space.

Boundary Value Problems: Variational Methods

More generally, a **variational method** works by approximating the differential equation by the equation

$$\langle r, \psi_i \rangle = \int_a^b r(t, y) \psi_i(t) dt = 0$$

so that the residual is forced to be orthogonal to a given set of test functions $\{\psi_i: i = 1, \dots, n\}$.

Clearly, the least squares method uses $\psi_i = \varphi_i''$.

Also, the collocation method uses $\psi_i = \delta(t - t_i)$, where t_i is the i -th collocation point and δ is the Dirac delta function.

A popular alternative is known as the **Galerkin method** where $\psi_i = \varphi_i$ is used. In this case, for the linear problem with $f(t, u, v) = f(t)$, the method yields the linear system with n equations in n unknowns,

$$\sum_{j=1}^n y_j \langle \varphi_j'', \varphi_i \rangle = \langle f, \varphi_i \rangle, \quad i = 1, \dots, n.$$

Typically, we assume that the basis functions satisfy the homogeneous boundary conditions $\varphi_i(a) = \varphi_i(b) = 0$ so that the approximation space $\text{span} \{\varphi_i: i = 1, \dots, n\}$ is a vector space. In this case, the linear system takes the following form

$$-\sum_{j=1}^n y_j \langle \varphi_j', \varphi_i' \rangle = \langle f, \varphi_i \rangle, \quad i = 1, \dots, n.$$



Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

Boundary Value Problems: Variational Methods


Example

Consider the two-point BVP


$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by


$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases}$$



$$\varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1 - t), & \frac{1}{2} < t \leq 1, \end{cases}$$



$$\varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$



Boundary Value Problems: Variational Methods


Example

Consider the two-point BVP


$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by


$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases}$$



$$\varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1 - t), & \frac{1}{2} < t \leq 1, \end{cases}$$



$$\varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$



We seek solution of the form $u(t) = t + w(t)$ where w satisfies the same ODE $w'' = 6t$, $0 < t < 1$, but with homogeneous boundary conditions $w(0) = 0$, $w(1) = 0$.

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by

$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1 - t), & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$

We seek solution of the form $u(t) = t + w(t)$ where w satisfies the same ODE $w'' = 6t$, $0 < t < 1$, but with homogeneous boundary conditions $w(0) = 0$, $w(1) = 0$.

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by

$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1-t), & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$

We seek solution of the form $u(t) = t + w(t)$ where w satisfies the same ODE $w'' = 6t$, $0 < t < 1$, but with homogeneous boundary conditions $w(0) = 0$, $w(1) = 0$. We, therefore, approximate w as

$$w(t) \approx y_2 \varphi_2(t)$$

where y_2 can be found as solution to the equation

$$-y_2 \int_0^1 \varphi_2'(t) \varphi_2'(t) dt = \int_0^1 6t \varphi_2(t) dt$$

that is, $-4y_2 = 3/2$.

only φ_2 satisfies the boundary cond hence use only that for approx w in the finite dimensional functionalm space

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by

$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1 - t), & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$

We seek solution of the form $u(t) = t + w(t)$ where w satisfies the same ODE $w'' = 6t$, $0 < t < 1$, but with homogeneous boundary conditions $w(0) = 0$, $w(1) = 0$. We, therefore, approximate w as

$$w(t) \approx y_2 \varphi_2(t)$$

where y_2 can be found as solution to the equation

$$-y_2 \int_0^1 \varphi_2'(t) \varphi_2'(t) dt = \int_0^1 6t \varphi_2(t) dt$$

that is, $-4y_2 = 3/2$. We get $u(t) = t - \frac{3}{8} \varphi_2(t)$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by

$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1-t), & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$

We seek solution of the form $u(t) = t + w(t)$ where w satisfies the same ODE $w'' = 6t$, $0 < t < 1$, but with homogeneous boundary conditions $w(0) = 0$, $w(1) = 0$. We, therefore, approximate w as

$$w(t) \approx y_2 \varphi_2(t)$$

where y_2 can be found as solution to the equation

$$-y_2 \int_0^1 \varphi_2'(t) \varphi_2'(t) dt = \int_0^1 6t \varphi_2(t) dt$$

that is, $-4y_2 = 3/2$. We get $u(t) = t - \frac{3}{8} \varphi_2(t) = \left(\frac{1}{2} \varphi_2(t) + \varphi_3(t)\right) - \frac{3}{8} \varphi_2(t)$

Boundary Value Problems: Variational Methods

Example

Consider the two-point BVP

$$\begin{aligned} u'' &= 6t, & 0 < t < 1, \\ u(0) &= 0, & u(1) = 1. \end{aligned}$$

We seek the approximate solution in the space of piecewise linear functions with respect to the partition $\{[0, 1/2], [1/2, 1]\}$ spanned by

$$\varphi_1(t) = \begin{cases} 2\left(\frac{1}{2} - t\right), & 0 \leq t \leq \frac{1}{2} \\ 0, & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_2(t) = \begin{cases} 2t, & 0 \leq t \leq \frac{1}{2}, \\ 2(1 - t), & \frac{1}{2} < t \leq 1, \end{cases} \quad \varphi_3(t) = \begin{cases} 0, & 0 \leq t \leq \frac{1}{2}, \\ 2\left(t - \frac{1}{2}\right), & \frac{1}{2} < t \leq 1. \end{cases}$$

We seek solution of the form $u(t) = t + w(t)$ where w satisfies the same ODE $w'' = 6t$, $0 < t < 1$, but with homogeneous boundary conditions $w(0) = 0$, $w(1) = 0$. We, therefore, approximate w as

$$w(t) \approx y_2 \varphi_2(t)$$

where y_2 can be found as solution to the equation

$$-y_2 \int_0^1 \varphi_2'(t) \varphi_2'(t) dt = \int_0^1 6t \varphi_2(t) dt$$

that is, $-4y_2 = 3/2$. We get $u(t) = t - \frac{3}{8} \varphi_2(t) = \left(\frac{1}{2} \varphi_2(t) + \varphi_3(t)\right) - \frac{3}{8} \varphi_2(t) = \frac{1}{8} \varphi_2(t) + \varphi_3(t)$.

Boundary Value Problems: Variational Methods



Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).



Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*



Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

*Spectral methods **(a)** provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a a large amount of work and storage to solve.*



Boundary Value Problems: Variational Methods

Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

*Spectral methods **(a)** provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a large amount of work and storage to solve. **(b)** Moreover, for some choices of basis functions (e.g., monomials), the resulting system of equations becomes very ill-conditioned as the number of basis functions grows.*

Boundary Value Problems: Variational Methods

Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

*Spectral methods **(a)** provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a large amount of work and storage to solve. **(b)** Moreover, for some choices of basis functions (e.g., monomials), the resulting system of equations becomes very ill-conditioned as the number of basis functions grows. **(c)** To avoid this difficulty, orthogonal basis functions are used.*

Boundary Value Problems: Variational Methods

Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

Spectral methods (a) provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a large amount of work and storage to solve. (b) Moreover, for some choices of basis functions (e.g., monomials), the resulting system of equations becomes very ill-conditioned as the number of basis functions grows. (c) To avoid this difficulty, orthogonal basis functions are used.

For finite element methods, the highly localized support of basis functions automatically makes the “nearly” orthogonal resulting in (a) a relatively well-conditioned equations to solve and (b) it also makes the system sparse, so that a much less work and storage is required to solve it.



Boundary Value Problems: Variational Methods

Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

Spectral methods (a) provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a large amount of work and storage to solve. (b) Moreover, for some choices of basis functions (e.g., monomials), the resulting system of equations becomes very ill-conditioned as the number of basis functions grows. (c) To avoid this difficulty, orthogonal basis functions are used.

For finite element methods, the highly localized support of basis functions automatically makes the “nearly” orthogonal resulting in (a) a relatively well-conditioned equations to solve and (b) it also makes the system sparse, so that a much less work and storage is required to solve it. However, (c) these methods typically produce lower accuracy, and therefore, a relatively large system may be required to achieve acceptable accuracy.

Boundary Value Problems: Variational Methods

Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

Spectral methods (a) provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a large amount of work and storage to solve. (b) Moreover, for some choices of basis functions (e.g., monomials), the resulting system of equations becomes very ill-conditioned as the number of basis functions grows. (c) To avoid this difficulty, orthogonal basis functions are used.

For finite element methods, the highly localized support of basis functions automatically makes the “nearly” orthogonal resulting in (a) a relatively well-conditioned equations to solve and (b) it also makes the system sparse, so that a much less work and storage is required to solve it. However, (c) these methods typically produce lower accuracy, and therefore, a relatively large system may be required to achieve acceptable accuracy. In principle, (d) the accuracy is increased by reducing the mesh spacing (h-refinement) or increasing the degree of the B-splines (p-refinement) or both.

Boundary Value Problems: Variational Methods

Remark

In our examples, we have used two kinds of basis functions, one with global support (where the basis function is non-zero over the entire interval, e.g., polynomials, trigonometric polynomials, etc.) and another with local support (e.g., B-splines, etc.).

*A method using basis functions with global support is typically referred to a **spectral** or a **pseudo-spectral method**. A method that used locally supported basis functions is an example of **finite element method**.*

Spectral methods (a) provide high accuracy for the number of points used but the global support of the basis functions yield a dense system of equations to solve that potentially requires a large amount of work and storage to solve. (b) Moreover, for some choices of basis functions (e.g., monomials), the resulting system of equations becomes very ill-conditioned as the number of basis functions grows. (c) To avoid this difficulty, orthogonal basis functions are used.

For finite element methods, the highly localized support of basis functions automatically makes the “nearly” orthogonal resulting in (a) a relatively well-conditioned equations to solve and (b) it also makes the system sparse, so that a much less work and storage is required to solve it. However, (c) these methods typically produce lower accuracy, and therefore, a relatively large system may be required to achieve acceptable accuracy. In principle, (d) the accuracy is increased by reducing the mesh spacing (h-refinement) or increasing the degree of the B-splines (p-refinement) or both. (e) The local nature also makes them more flexible than spectral methods in dealing with complicated domains and boundary conditions.

Numerical Analysis & Scientific Computing II

Lesson 3

Boundary Value Problems for ODEs

3.2 Shooting Method

3.3 Finite Difference Method

3.4 Variational Methods

- Convergence Analysis for Galerkin Method



Akash Anand
MATH, IIT KANPUR

Boundary Value Problems: Variational Methods

For analyzing the computational error in the Galerkin method, note that the solution u to the boundary value problem

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

and any function of the form

$$v(t, x) = \sum_{i=1}^n x_i \varphi_i(t),$$

with $\varphi_i(a) = \varphi_i(b) = 0$, satisfies

$$\int_a^b u''(t) v(t, x) dt = \int_a^b f(t) v(t, x) dt.$$

Boundary Value Problems: Variational Methods

For analyzing the computational error in the Galerkin method, note that the solution u to the boundary value problem

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

and any function of the form

$$v(t, x) = \sum_{i=1}^n x_i \varphi_i(t),$$

with $\varphi_i(a) = \varphi_i(b) = 0$, satisfies

$$-\int_a^b u'(t) v'(t, x) dt = \int_a^b u''(t) v(t, x) dt = \int_a^b f(t) v(t, x) dt.$$

Boundary Value Problems: Variational Methods

For analyzing the computational error in the Galerkin method, note that the solution u to the boundary value problem

$$\begin{aligned}u'' &= f(t), & a < t < b, \\u(a) &= 0, & u(b) = 0,\end{aligned}$$

and any function of the form

$$v(t, x) = \sum_{i=1}^n x_i \varphi_i(t),$$

with $\varphi_i(a) = \varphi_i(b) = 0$, satisfies

$$-\int_a^b u'(t) v'(t, x) dt = \int_a^b u''(t) v(t, x) dt = \int_a^b f(t) v(t, x) dt.$$

Similarly, for the approximate solution $v(t, y)$, we have

$$-\int_a^b v'(t, y) v'(t, x) dt = \int_a^b f(t) v(t, x) dt.$$

Boundary Value Problems: Variational Methods

For analyzing the computational error in the Galerkin method, note that the solution u to the boundary value problem

$$\begin{aligned} u'' &= f(t), & a < t < b, \\ u(a) &= 0, & u(b) &= 0, \end{aligned}$$

and any function of the form

$$v(t, x) = \sum_{i=1}^n x_i \varphi_i(t),$$

with $\varphi_i(a) = \varphi_i(b) = 0$, satisfies

$$-\int_a^b u'(t) v'(t, x) dt = \int_a^b u''(t) v(t, x) dt = \int_a^b f(t) v(t, x) dt.$$

Similarly, for the approximate solution $v(t, y)$, we have

$$-\int_a^b v'(t, y) v'(t, x) dt = \int_a^b f(t) v(t, x) dt.$$

Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z)) v'(t, x) dt = \int_a^b (u'(t) - v'(t, z)) v'(t, x) dt.$$



Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Boundary Value Problems: Variational Methods



Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*)



Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*), and check that (*exercise*), for $w \in C_0^1([a, b])$, we have

$$\|w\|_{L^2} \leq c(b-a)\|w'\|_{L^2}$$

for some $c > 0$.

Boundary Value Problems: Variational Methods

Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*), and check that (*exercise*), for $w \in C_0^1([a, b])$, we have

$$\|w\|_{L^2} \leq c(b-a)\|w'\|_{L^2}$$

for some $c > 0$. In particular, we have

$$\|v(\cdot, y) - v(\cdot, z)\|_{H^1}^2 \leq (1 + c^2(b-a)^2) \int_a^b (v'(t, y) - v'(t, z))^2 dt$$

Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*), and check that (*exercise*), for $w \in C_0^1([a, b])$, we have

$$\|w\|_{L^2} \leq c(b-a)\|w'\|_{L^2}$$

for some $c > 0$. In particular, we have

$$\begin{aligned} \|v(\cdot, y) - v(\cdot, z)\|_{H^1}^2 &\leq (1 + c^2(b-a)^2) \int_a^b (v'(t, y) - v'(t, z))^2 dt \\ &= (1 + c^2(b-a)^2) \int_a^b (u'(t) - v'(t, z))(v'(t, y) - v'(t, z)) dt \end{aligned}$$

Boundary Value Problems: Variational Methods

Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*), and check that (*exercise*), for $w \in C_0^1([a, b])$, we have

$$\|w\|_{L^2} \leq c(b-a)\|w'\|_{L^2}$$

for some $c > 0$. In particular, we have

$$\begin{aligned} \|v(\cdot, y) - v(\cdot, z)\|_{H^1}^2 &\leq (1 + c^2(b-a)^2) \int_a^b (v'(t, y) - v'(t, z))^2 dt \\ &= (1 + c^2(b-a)^2) \int_a^b (u'(t) - v'(t, z))(v'(t, y) - v'(t, z)) dt \\ &\leq (1 + c^2(b-a)^2) \left(\int_a^b |u'(t) - v'(t, z)|^2 dt \right)^{\frac{1}{2}} \left(\int_a^b |v'(t, y) - v'(t, z)|^2 dt \right)^{\frac{1}{2}} \end{aligned}$$

Boundary Value Problems: Variational Methods

Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*), and check that (*exercise*), for $w \in C_0^1([a, b])$, we have

$$\|w\|_{L^2} \leq c(b-a)\|w'\|_{L^2}$$

for some $c > 0$. In particular, we have

$$\begin{aligned} \|v(\cdot, y) - v(\cdot, z)\|_{H^1}^2 &\leq (1 + c^2(b-a)^2) \int_a^b (v'(t, y) - v'(t, z))^2 dt \\ &= (1 + c^2(b-a)^2) \int_a^b (u'(t) - v'(t, z))(v'(t, y) - v'(t, z)) dt \\ &\leq (1 + c^2(b-a)^2) \left(\int_a^b |u'(t) - v'(t, z)|^2 dt \right)^{\frac{1}{2}} \left(\int_a^b |v'(t, y) - v'(t, z)|^2 dt \right)^{\frac{1}{2}} \\ &\leq (1 + c^2(b-a)^2) \|u - v(\cdot, z)\|_{H^1} \|v(\cdot, y) - v(\cdot, z)\|_{H^1} \end{aligned}$$

Boundary Value Problems: Variational Methods

Thus, for any $z \in \mathbb{R}^n$, we have

$$\int_a^b (v'(t, y) - v'(t, z))v'(t, x)dt = \int_a^b (u'(t) - v'(t, z))v'(t, x)dt.$$

Define the H^1 norm $\|\cdot\|_{H^1}$ on $C_0^1([a, b]) = \{w \in C^1([a, b]): w(a) = w(b) = 0\}$ as

$$\|w\|_{H^1}^2 := \|w\|_{L^2}^2 + \|w'\|_{L^2}^2$$

(*exercise: check that it is a norm*), and check that (*exercise*), for $w \in C_0^1([a, b])$, we have

$$\|w\|_{L^2} \leq c(b-a)\|w'\|_{L^2}$$

for some $c > 0$. In particular, we have

$$\begin{aligned} \|v(\cdot, y) - v(\cdot, z)\|_{H^1}^2 &\leq (1 + c^2(b-a)^2) \int_a^b (v'(t, y) - v'(t, z))^2 dt \\ &= (1 + c^2(b-a)^2) \int_a^b (u'(t) - v'(t, z))(v'(t, y) - v'(t, z)) dt \\ &\leq (1 + c^2(b-a)^2) \left(\int_a^b |u'(t) - v'(t, z)|^2 dt \right)^{\frac{1}{2}} \left(\int_a^b |v'(t, y) - v'(t, z)|^2 dt \right)^{\frac{1}{2}} \\ &\leq (1 + c^2(b-a)^2) \|u - v(\cdot, z)\|_{H^1} \|v(\cdot, y) - v(\cdot, z)\|_{H^1} \end{aligned}$$

Therefore,

$$\|v(\cdot, y) - v(\cdot, z)\|_{H^1} \leq (1 + c^2(b-a)^2) \|u - v(\cdot, z)\|_{H^1}.$$



Therefore,

$$\|v(\cdot, y) - v(\cdot, z)\|_{H^1} \leq (1 + c^2(b - a)^2)\|u - v(\cdot, z)\|_{H^1}.$$

Now,

$$\|u - v(\cdot, y)\|_{H^1} \leq \|u - v(\cdot, z)\|_{H^1} + \|v(\cdot, z) - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2)\|u - v(\cdot, z)\|_{H^1}.$$

As $z \in \mathbb{R}^n$ is arbitrary, we conclude that

$$\|u - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1}$$

Boundary Value Problems: Variational Methods

Therefore,

$$\|v(\cdot, y) - v(\cdot, z)\|_{H^1} \leq (1 + c^2(b - a)^2) \|u - v(\cdot, z)\|_{H^1}.$$


Now,

$$\|u - v(\cdot, y)\|_{H^1} \leq \|u - v(\cdot, z)\|_{H^1} + \|v(\cdot, z) - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \|u - v(\cdot, z)\|_{H^1}.$$

As $z \in \mathbb{R}^n$ is arbitrary, we conclude that

$$\|u - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1}$$

Best approximation
of u by functions in
the subspace
 $\text{span}\{\varphi_1, \dots, \varphi_n\}$



Boundary Value Problems: Variational Methods

Therefore,

$$\|v(\cdot, y) - v(\cdot, z)\|_{H^1} \leq (1 + c^2(b - a)^2) \|u - v(\cdot, z)\|_{H^1}.$$

Now,

$$\|u - v(\cdot, y)\|_{H^1} \leq \|u - v(\cdot, z)\|_{H^1} + \|v(\cdot, z) - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \|u - v(\cdot, z)\|_{H^1}.$$


As $z \in \mathbb{R}^n$ is arbitrary, we conclude that

$$\|u - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1}$$

For piecewise linear elements, we have

$$\inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1} \leq Ch \|u''\|_{L^2}$$

Best approximation
of u by functions in
the subspace
 $\text{span}\{\varphi_1, \dots, \varphi_n\}$



Boundary Value Problems: Variational Methods

Therefore,

$$\|v(\cdot, y) - v(\cdot, z)\|_{H^1} \leq (1 + c^2(b - a)^2) \|u - v(\cdot, z)\|_{H^1}.$$

Now,

$$\|u - v(\cdot, y)\|_{H^1} \leq \|u - v(\cdot, z)\|_{H^1} + \|v(\cdot, z) - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \|u - v(\cdot, z)\|_{H^1}.$$

As $z \in \mathbb{R}^n$ is arbitrary, we conclude that

$$\|u - v(\cdot, y)\|_{H^1} \leq (2 + c^2(b - a)^2) \inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1}$$

For piecewise linear elements, we have

$$\inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1} \leq Ch \|u''\|_{L^2}$$

and, more generally, for elements of degree r , we have an estimate

$$\inf_{z \in \mathbb{R}^n} \|u - v(\cdot, z)\|_{H^1} \leq Ch^r \|u\|_{H^{r+1}}$$

Best approximation
of u by functions in
the subspace
 $\text{span}\{\varphi_1, \dots, \varphi_n\}$

