Coherence and Entanglement Monogamy in the Discrete Analogue of Analog Grover Search

Namit Anand*
Department of Physics
National Institute of Science Education and Research,
Bhubaneswar, 752050, India

Arun Kumar Pati†
Quantum Information and Computation Group
Harish-Chandra Research Institute, Chhatnag Road,
Jhunsi, Allahabad 211 019, India
(Dated: November 4, 2016)

Grover's search algorithm is the optimal quantum algorithm that can search an unstructured database quadratically faster than any known classical algorithm. The role of entanglement and correlations in the search algorithm has been studied in great detail and it is known that entanglement between the qubits is necessary to gain a quadratic speedup, for pure state implementation of the Grover search algorithm. Here, we systematically investigate the behavior of quantum coherence, concurrence and monogamy of entanglement in the discrete analogue of the *analog analogue of Grover search algorithm*. The analog analogue of Grover search is a continuous time quantum algorithm based on adiabatic Hamiltonian evolution that gives a quadratic speedup, similar to the original Grover search algorithm. We analytically show that the decrease of quantum coherence, quantified using various coherence monotones, is a clear signature of attaining the maximum success probability in the analog Grover search. We also show that for any two qubit reduced density matrix of the system, the concurrence evolves in close vicinity to the increasing rate of success probability, hence implying that entanglement is indeed necessary for the discrete analogue of the analog Grover search algorithm. Furthermore, we show that the system satisfies a N-party monogamy inequality for arbitrary times, hence bounding the amount of pairwise entanglement during the quantum search.

I. INTRODUCTION

In the early 1980s, Feynman [1] first suggested that in order to simulate physical systems efficiently, one would be ideally suited to constructing a computer built from quantum mechanical systems since the physical world is ultimately quantum mechanical. This idea, that quantum systems can be visualized as information processing devices, is at the heart of the theory of quantum information and computation. Deutsch [2] constructed the first example of an algorithm which would require two queries to solve on a classical computer but that can be solved with only one quantum query. Subsequently, Deutsch and Jozsa [3], Bernstein and Vazirani [4] and Simon [5] demonstrated the striking difference between the classical and quantum query complexity. This culminated with Grover's algorithm [6] achieving a quadratic speedup over any classical algorithm for the unstructured search problem and Shor's algorithm [7] that could factor integers efficiently and also calculate discrete logarithms.

Grover's algorithm gives an optimal solution to the unstructured search problem, the problem of deciding whether a black-box Boolean function has any input that evaluates to 1. It provides a quadratic temporal speedup over the best classical search algorithms, even when they both require the same spatial resources to perform the search task. Zalka [8] proved the optimality of the Grover search algorithm. Grover's algorithm was subsequently generalized to the framework of amplitude amplification and to counting the number of solutions by Brassard *et al.* [9]. Grover's search algorithm has since been applied to a wide variety of similar problems[10][11][12].

The usual paradigm of computation (quantum or classical) is defined in a discrete setting. However, adiabatic quantum computation [13] provides a continuous time model for quantum computing by using the quantum adiabatic theorem. Here, one begins by finding an appropriate Hamiltonian whose ground state describes the solution to the problem of interest. Next, a system with a simple Hamiltonian is prepared and initialized to the ground state. Finally, the simple Hamiltonian is adiabatically evolved to the desired Hamiltonian. By the adiabatic theorem, the system remains in the ground state at all times during this adibatic evolution, depending on spectral properties of the Hamiltonian [14][15]. At the end, the final state of the system describes the solution to the problem. The time complexity for an adiabatic algorithm is simply defined as the time taken to complete the adiabatic evolution, which is known to depend on the spectral gap of the Hamiltonian. The adiabatic model of quantum computing is universal and equivalent to standard quantum computation [16]. Farhi and Gutmann constructed the first analog algorithm for solving

* namitniser11@gmail.com
† akpati@hri.res.in

the Grover search problem [17] referred to hereafter as the *analog analogue of Grover search algorithm* or simply the *analog Grover search algorithm*.

It is understood that the speedup in quantum algorithms comes from quantum mechanical features like quantum coherence and quantum correlations like quantum entanglement, that exists amongst the qubits. Quantum correlations, especially quantum entanglement is one of the crucial resources in quantum information theory and has been extensively studied [18]. The indispensible role of entanglement in quantum information tasks such as quantum teleportation [19], superdense coding [20] and remote state preparation [21] [22] , etc. has been quantified and understood deeply. However, despite showing that quantum entanglement is *necessary* for the pure state implementation of Grover search [23] and Shor's factorizing algorithm [24], the innate role of entanglement is not clear for general quantum computing tasks.

One of the fundamental attributes of quantum systems is their ability to exist in linear superpositions of different physical states. This physical phenomenon is called quantum superposition. Quantum coherence arises from superposition and is at the heart of several quantum features like quantum interference, multiparticle entanglement [25], quantum biology [26] [27], quantum thermodynamics [28] [29] [30] [31], quantum game theory [32] [33] [34] etc., which in turn are some of the most important applications of quantum physics and quantum information science. Inspired from the resource theory of entanglement [18] [36], there has been a lot of effort recently to quantify coherence as a resource theory [37]. In a standard resource-theoretic treatment of coherence, the "incoherent" states are those that are diagonal in some fixed reference basis. The amount of coherence is then defined as the distance from these reference states. A 'coherence measure' is a real-valued function over the quantum state-space, such that it vanishes for all the states that are deemed to be incoherent and cannot increase under some class of operations that preserve incoherence. In this regard, several coherence measures like the l_1-norm of coherence and the relative entropy of coherence have also been introduced.

In this paper, we quantitatively analyze the role of quantum coherence and monogamy of entanglement [38] [39] in the analog Grover search algorithm. We give a geometric representation to the analog Grover search where we show that the time evolving state vector follows a geodesic in the two-dimensional subspace of the N-dimensional Hilbert space. To quantify the dynamics of coherence in this analog Grover search, we use two coherence monotones namely the l_1-norm and the relative entropy of coherence. We then discretize the analog Grover search algorithm to study how quantum entanglement evolves parallel to the system's adiabatic evolution. We use an entanglement monotone called the concurrence [40] to study the same.

We find that the amount of coherence (as quantified by the coherence measures) is non-zero at all times during the search. The only time this coherence goes to zero is when the search algorithm attains the maximum success probability equal to one. Since we begin with a maximally coherent state, in a resource theoretic sense, this amount of coherence is actually consumed during the quantum search. Using coherence monotones, we relate the success probability of the search algorithm with quantum the amount of coherence. Further, during the analog search, the final state is a product state and so all entanglement monotones must go to zero at the end of the searching process. The bipartite entanglement entropy, concurrence and the monogamy score, all behave in accord with this, since for both qubit partitions, these measures of quantum correlation go to zero and mark the completion of the search algorithm. Furthermore, the two qubit concurrence and the monogamy score peak simultaneously with the increasing rate of success probability, hence implying that entanglement is indeed necessary for the discrete analogue of the analog analogue of Grover search algorithm.

The paper is organized as follows. In Sec. II, we briefly review the analog Grover search algorithm. In Sec. **??**, we geometrically analyze the analog Grover search algorithm and show that the trajectory traced by the time evolving state vector in the N-dimensional Hilbert space is a geodesic. In Sec. III, we use the two coherence monotones and calculate the coherence dynamics and elucidate their behavior. In Sec. IV, we explore the entanglement entropy and the concurrence and demonstrate the synonymous behavior of the rate of change of concurrence in the continuous and the discrete Grover search. In Sec. **??** we calculate the two-qubit concurrence and show its connection to the success probability of the search algorithm. In Sec. IV C we use the monogamy inequality to bound the bipartite entanglement at all times. We conclude our paper with the discussions in Sec. V.

II. GEOMETRY OF ANALOG SEARCH

Adiabatic quantum computation can be described as a controlled Hamiltonian time evolution of a system, obeying the Schrodinger equation

$$i\frac{d}{dt}|\psi\rangle = H(t)|\psi\rangle. \tag{1}$$

We briefly recapitulate the analog analogue of Grover search algorithm [17], where the problem is to use quantum evolution to find a marked state among N orthonormal states.

Imagine that we are given a Hamiltonian in an N dimensional Hilbert space such that it has only one non-zero eigenvalue, $E \neq 0$. The task then is to find the corresponding eigenvector $|w\rangle$ which has eigenvalue E. The Hamiltonian can be represented as

$$H_w = E|w\rangle\langle w| \tag{2}$$

with $|w\rangle$ unknown and $\langle w|w\rangle = 1$. Now we choose some normalized vector $|s\rangle$, which is independent of $|w\rangle$ (since we do not know what $|w\rangle$ is yet). Now we add to H_w, the "driving" Hamiltonian [17]

$$H_D = E|s\rangle\langle s| \tag{3}$$

so that the full Hamiltonian is given by

$$H = H_w + H_D. \tag{4}$$

Now, starting from the initial state $|s\rangle$ at $t = 0$, we calculate the time evolution of the state $|\psi(t)\rangle$. Since the total Hamiltonian is time-independent, we can simply write the time evolved state as:

$$|\psi(t)\rangle = e^{-iHt}\ |s\rangle. \tag{5}$$

It suffices to confine our attention to the two dimensional subspace spanned by $|s\rangle$ and $|w\rangle$. The vectors $|s\rangle$ and $|w\rangle$ are not orthogonal (in general) and let us denote their inner product as x, i.e., $\langle s|w\rangle = x$, where x can be taken to be real and positive since any phase in $\langle s|w\rangle$ can ultimately be absorbed in $|s\rangle$. Now the vectors

$$|r\rangle = \frac{1}{\sqrt{1-x^2}}\left(|s\rangle - x|w\rangle\right) \tag{6}$$

and $|w\rangle$ are orthonormal. In the $|w\rangle$, $|r\rangle$ basis the Hamiltonian is given by

$$H = E\begin{bmatrix} 1+x^2 & x\sqrt{1-x^2} \\ x\sqrt{1-x^2} & 1-x^2 \end{bmatrix} \tag{7}$$

and

$$|s\rangle = \begin{bmatrix} x \\ \sqrt{1-x^2} \end{bmatrix}. \tag{8}$$

Now the state of the system at time t is given by

$$|\psi(t)\rangle = e^{-iEt}\begin{bmatrix} x\ \cos(Ext) - i\ \sin(Ext) \\ \sqrt{1-x^2}\cos(Ext) \end{bmatrix}. \tag{9}$$

Thus, we see that at time t the probability of finding the state $|w\rangle$ is given by

$$P(t) = \sin^2(Ext) + x^2\ \cos^2(Ext) \tag{10}$$

and the probability is one at time t_m given by

$$t_m = \frac{\pi}{2Ex}. \tag{11}$$

The inner product between the vectors $|w\rangle$ and $|s\rangle$ (defined as x above), is assumed to be non-zero, else it will take infinite time for the quantum search to complete.

The analog model has been generalized further. For example, the same algorithm was recast in the form of a spatial search problem [41], where there is a N-dimensional lattice and the basis state $|i\rangle$ is localized at the ith lattice site. The on-site potential energy, E is zero everywhere except at $|w\rangle$, where it takes value 1. The objective is same as before, to reach the marked state starting from an equal superposition of all the $|i\rangle$'s. The kinetic term is formulated through the Laplacian of the lattice, which effectively introduces uniform hopping to all nearest neighbors from any given lattice sites, and is kept constant.

III. QUANTUM COHERENCE IN GROVER SEARCH

Quantum coherence is one of the salient features of the quantum world. As this crucial feature drives several quantum technologies, it is very desirable to quantify the usefulness of coherence as a resource. This is done using the mathematical framework called a 'resource theory'. To characterize something as a resource, we must first impose certain constraints on what specific physical operations are allowed (e.g. the local operations and classical communication(LOCC) framework for quantum entanglement [44] restricts one from performing joint quantum operations between distant laboratories), which define the freely accessible operations. To be able to execute general quantum operations under such a constraint then requires some "special" quantum states that contain a relevant resource (e.g. entangled states) and can be consumed in the process. In fact, a quantitative resource theory for entanglement is already in place [18] [36] and has later expanded to encompass a wider range of operational phenomenon [45] [46].

However, for a long time there did not exist a resource theoretic framework to quantify the aspects of coherence. Coherence was often defined in terms of functions of the off-diagonal entries of a density matrix, and the definition justified on the basis of physical intuition, which in turn lead to the idea of decoherence developing along similar lines [47]. Eventually, in 2014, Baumgratz *et al.* [37] defined a resource theory of coherence and rigourously quantified the role of coherence in close analogy to the resource theory of entanglement.

Let us formally introduce the measures of coherence in the framework of resource theory, that is based on the set of incoherent operations and incoherent states. Coherence is a basis dependent property and hence first, we must fix a *reference basis*. The choice of the reference basis may be dictated by the underlying physics of the problem(say the energy eigenbasis) or the task to be performed where we wish to use coherence. Let $\{|i\rangle_{i=0,1,2,\ldots,d-1}\}$ be a basis for $\mathcal{H}^d$. The set of all density matrices that are diaogonal in this chosen basis form the so called "incoherent states". The set of all the states of the form $\rho_I = \sum_i p_i|i\rangle\langle i|$, where $p_i \geq 0$ and $\sum_i p_i = 1$, forms the set $\mathcal{I}$ of all incoherent states. Any quantum state that does not belong to the set $\mathcal{I}$ will be called a coherent state and will act as a resource. A maximally coherent state in this basis is then given

by $|\psi_d\rangle = \frac{1}{\sqrt{d}}\sum_{i=0}^{d-1}|i\rangle$ as any other state can be created from $|\psi_d\rangle$ using only the set of 'incoherent operations'. Baumgratz *et al.* [37] define incoherent operations as the set of trace preserving completely positive maps $\Lambda : B(H) \longmapsto B(H)$ admitting a set of Kraus operators K_n [42] such that $\sum_n {K_n}^\dagger K_n = \mathbb{1}$ (trace preservation) and, for all n and $\rho \in \mathcal{I}$,

$$\frac{K_n \rho {K_n}^\dagger}{Tr[K_n \rho {K_n}^\dagger]} \in \mathcal{I}. \tag{12}$$

The definition of the set of incoherent operations is not unique and so we redirect the reader to a recent review for further discussion [48]. We shall use two coherence monotones introduced in [37], namely, the l_1-norm of coherence and the relative entropy of coherence.

Given a density matrix ρ for the system, the l_1-norm of coherence is defined as

$$\mathcal{C}_{l_1}(\rho) = \sum_{i,j;i\neq j} \left|\rho_{i,j}\right|. \tag{13}$$

For the analog Grover search, the l_1-norm of coherence for the state given in eqn. 9 is

$$\mathcal{C}_{l_1}(\rho) = 2\left|\cos(Ext)\right| \sqrt{(1-x^2)\left(x^2\cos^2(Ext)+\sin^2(Ext)\right)}.$$

Similarly, the relative entropy of coherence is given as:

$$\mathcal{C}_r(\rho) = S(\rho_{diag}) - S(\rho). \tag{14}$$

The relative entropy of coherence is calculated to be

$$\mathcal{C}_r(\rho) = -\left(1-x^2\right)\cos^2(Ext)\log_2\left(\left(1-x^2\right)\cos^2(Ext)\right) - \left(x^2\cos^2(Ext)+\sin^2(Ext)\right)\log_2\left(x^2\cos^2(Ext)+\sin^2(Ext)\right).$$

The l_1-norm of coherence has a closed form expression in terms of the probability of success given by,

$$\mathcal{C}_{l_1}(P) = 2\sqrt{P(1-P)}. \tag{15}$$

Similarly, the relative entropy of coherence in terms of the probability of success takes the form of a binary entropy function, i.e.,

$$\mathcal{C}_r(P) = -\left(P\log_2 P + (1-P)\log_2(1-P)\right). \tag{16}$$

It can be observed analytically from equations 15, 16 and numerically from the figure 1 that both coherence measures go to zero iff the probability of success peaks to one, hence acting as a clear signature for the success of the search algorithm. Even for other general initial states during the analog Grover search, the coherence attains a minima as the success probability peaks to 1.



FIG. 1. Variations of the l_1-norm, the relative entropy of coherence and the probability of success as a function of time at parameter values $E = 1$ and $x = 0.707$.

A. Discrete analogue of the analog analogue of Grover search algorithm

It is easier to quantify the role of entanglement in the original Grover search due to its discrete nature, however it is not so in the analog version. To overcome this obstacle, we consider a mapping through which (i) the N eigenstates of the system are mapped to the logical states of n qubits (where $N = 2^n$). (ii) The Hamiltonian initially acting on the state $|\psi\rangle$ of the system now acts on the n qubit system. (iii) In this setting entanglement can arise between the n qubits.

Now, consider the initial state to be an equal superposition of all eigenstates, i.e.,

$$|s\rangle = \frac{1}{\sqrt{N}}\sum_{i=0}^{n-1}|i\rangle \tag{17}$$

and the final state $|w\rangle$ to be of the form $|000...000\rangle \equiv |0\rangle$. The $|\psi_w, t\rangle$ can be expressed in the $|w\rangle, |r\rangle$ basis as:

$$|\psi_w, t\rangle = \alpha(t)|w\rangle + \beta(t)|r\rangle, \tag{18}$$

where $x = \frac{1}{\sqrt{N}}$, $\alpha = x\cos(Ext) - i\sin(Ext)$ and $\beta = \sqrt{1-x^2}\cos(Ext)$.

Therefore the density matrix in the $|w\rangle, |r\rangle$ basis is:

$$\begin{aligned}\rho = \alpha^2(t)|w\rangle\langle w| + \alpha(t)\beta(t)|w\rangle\langle r| \\ +\alpha^*(t)\beta(t)|r\rangle\langle w| + \beta^2(t)|r\rangle\langle r|.\end{aligned} \tag{19}$$

After discretizing the analog Grover search, the l_1-norm of coherence for the density matrix in eqn. 19 is

analytically calculated to be

$$\mathcal{C}_{l_1}(\rho) = 2\sqrt{\left(1-\frac{1}{N}\right)\left(\frac{1}{N}\cos^2(\frac{Et}{\sqrt{N}})+\sin^2(\frac{Et}{\sqrt{N}})\right)} \text{ x } \left|\cos(\frac{Et}{\sqrt{N}})\right|.$$

Similarly the relative entropy of coherence is calculated to be

$$\mathcal{C}_r(\rho) = -\left(1-\frac{1}{N}\right)\cos^2(\frac{Et}{\sqrt{N}})\log_2\left(\left(1-\frac{1}{N}\right)\cos^2(\frac{Et}{\sqrt{N}})\right) - \left(\frac{1}{N}\cos^2(\frac{Et}{\sqrt{N}})+\sin^2(\frac{Et}{\sqrt{N}})\right)\log_2\left(\frac{1}{N}\cos^2(\frac{Et}{\sqrt{N}})+\sin^2(\frac{Et}{\sqrt{N}})\right)$$

The l_1-norm and the relative entropy of coherence display a similar behavior as before, and it is important to note that the initial state for the analog Grover search algorithm is an equal superposition of all eigenstates of the Hamiltonian, i.e.,

$$|s\rangle = \frac{1}{\sqrt{N}}\sum_{i=0}^{n-1}|i\rangle, \tag{20}$$

which corresponds to a maximally coherent state in this eigenbasis. However, at time $t = \frac{\pi\sqrt{N}}{2E}$ (which is the time at which the search algorithm succeeds), the coherence is reduced to zero. This implies that a maximally coherent state is actually consumed during the search algorithm.

IV. ENTANGLEMENT MONOGAMY IN GROVER SEARCH

A. Entanglement entropy

Entanglement entropy is a measure of entanglement for many body quantum states. Bipartite entanglement entropy is defined with respect to a bipartition of a state into two partitions say A and B. For this, consider a quantum system whose Hilbert space $\mathcal{H}$ is constructed from two Hilbert spaces $\mathcal{H}_A$ and $\mathcal{H}_B$ such that $\mathcal{H} = \mathcal{H}_A \otimes \mathcal{H}_B$. The bipartite entanglement entropy S of a state $|\Psi\rangle \in \mathcal{H}$ is defined as the von Neumann entropy of either of its reduced states; the result is independent of which one we pick (this can be easily proved using the Schmidt deomposition of the original pure state). That is, for a pure state $\rho_{AB} = |\Psi\rangle\langle\Psi|_{AB}$, it is given by

$$\mathcal{S}(\rho_A) = -\operatorname{Tr}(\rho_A \log \rho_A) = -\operatorname{Tr}(\rho_B \log \rho_B) = \mathcal{S}(\rho_B),$$

where $\rho_A = \operatorname{Tr}_B(\rho_{AB})$ and $\rho_B = \operatorname{Tr}_A(\rho_{AB})$ are the reduced density matrices for each partition. The von Neumann entanglement entropy in the eigenbasis is defined as

$$S(\rho_l) = -\sum_i \lambda_i \log \lambda_i. \tag{21}$$

Now we calculate the entanglement entropy of the system by partitioning it as one qubit (say l) vs the other $n-1$ qubits.

The reduced density matrix for the lth qubit, ρ_l is obtained by tracing out the other $(n-1)$ qubits and is given as

$$\begin{bmatrix} |\alpha|^2 + \frac{(N-2)\beta^2}{2N-2} & \frac{(N-2)\beta^2}{2N-2} + \frac{\alpha\beta}{\sqrt{N-1}} \\ \frac{(N-2)\beta^2}{2N-2} + \frac{\alpha^*\beta}{\sqrt{N-1}} & \frac{\beta^2 N}{2N-2} \end{bmatrix}. \tag{22}$$

The eigenvalues of the matrix λ_i are given by

$$\frac{2N \pm \sqrt{N\left((N-2)\cos\left(\frac{4Et}{\sqrt{N}}\right)+3N+2\right)}}{4N}. \tag{23}$$

B. Concurrence

Concurrence is an entanglement measure, which for the special case of a pair of qubits is closely related to the entanglement of formation [40]. For an arbitrary two qubit density matrix ρ, concurrence is defined as follows - we first define a "spin-flipped" density matrix, γ as $(\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$ where σ_y is the Pauli matrix $\begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$. Then we calculate the square root of the eigenvalues of the matrix $(\rho\gamma)$ and arrange them in decreasing order (say λ_1,λ_2,λ_3,λ_4). The concurrence is then defined as

$$\max\,(\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4, 0).$$

For pure states, the concurrence is also defined via another quantity known as the "tangle", where

$$\tau(\psi) = 2(1 - Tr(\rho^2)) \tag{24}$$

and the concurrence $= \sqrt{\tau}$.

The above concurrence can be extended to the situation of higher dimension pure bipartite state$|\psi\rangle$. We will use this form to study the entanglement of Grover search. The concurrence of $|\psi\rangle$ is defined as

$$C(|\psi\rangle) = \sqrt{\frac{d}{d-1}(1 - Tr\rho_r^2)}, \tag{25}$$

where ρ_r is the reduced density matrix obtained by tracing out one of the two subsystems, and d is the dimension of ρ_r.

Concurrence for the pure state across the bipartition $A_1|A_2A_3...A_N$ is calculated using the density matrix in eqn. 22 to be

$$C(\psi) = \sqrt{\frac{(N-2)}{2N}}\left|\sin\left(\frac{2Et}{\sqrt{N}}\right)\right|. \tag{26}$$



FIG. 2. Variations of the entanglement entropy, concurrence (across system partition of $A_1|A_2A_3...A_N$) and the probability of success as a function of time at parameter values $E = 1$ and $N = 4$.

The entanglement entropy and concurrence across the system partition of $A_1|A_2A_3...A_N$ is calculated analytically and its variation with time along with the probability of success is shown in Fig. 2. Note that for both entanglement measures, the entanglement goes to zero as the success probability peaks to 1.

Rate of change of the concurrence with time is calculated to be

$$\frac{dC(\psi)}{dt} = \frac{E\sqrt{2(N-2)}}{N}\cos\left(\frac{2Et}{\sqrt{N}}\right). \tag{27}$$

Figure 3 shows the analogous rate of change between the discrete and the continuous Grover search and suggests that the chosen mapping in Sec. III A preserves the properties of the original Grover search algorithm.

The reduced density matrix for two qubits is:

$$\rho_{AB} = \begin{bmatrix} \frac{(\frac{N}{4}-1)\beta^2}{N-1} + |\alpha|^2 & \frac{(\frac{N}{4}-1)\beta^2}{N-1} + \frac{\alpha\beta}{\sqrt{N-1}} & \frac{(\frac{N}{4}-1)\beta^2}{N-1} + \frac{\alpha\beta}{\sqrt{N-1}} & \frac{(\frac{N}{4}-1)\beta^2}{N-1} + \frac{\alpha\beta}{\sqrt{N-1}} \\ \frac{(\frac{N}{4}-1)\beta^2}{N-1} + \frac{\alpha^*\beta}{\sqrt{N-1}} & \frac{\beta^2 N}{4(N-1)} & \frac{\beta^2 N}{4(N-1)} & \frac{\beta^2 N}{4(N-1)} \\ \frac{(\frac{N}{4}-1)\beta^2}{N-1} + \frac{\alpha^*\beta}{\sqrt{N-1}} & \frac{\beta^2 N}{4(N-1)} & \frac{\beta^2 N}{4(N-1)} & \frac{\beta^2 N}{4(N-1)} \\ \frac{(\frac{N}{4}-1)\beta^2}{N-1} + \frac{\alpha^*\beta}{\sqrt{N-1}} & \frac{\beta^2 N}{4(N-1)} & \frac{\beta^2 N}{4(N-1)} & \frac{\beta^2 N}{4(N-1)} \end{bmatrix} \tag{28}$$

The concurrence of an arbitrary two-qubit state is calculated according to the formula in section IV B, using the spin flipped qubit. The eigenvalues for the density matrix $\rho\gamma$ are:

$\lambda_1 = 0$, $\lambda_2 = 0$, $\lambda_3 = \frac{1}{4N}\left(\frac{N+4}{4} - \sqrt{N}\right)\sin^2\left(\frac{2Et}{\sqrt{N}}\right)$ and $\lambda_4 = \frac{1}{4N}\left(\frac{N+4}{4} + \sqrt{N}\right)\sin^2\left(\frac{2Et}{\sqrt{N}}\right)$. Therefore the two qubit concurrence is analytically calculated to be

$$\mathcal{C}(\rho_{\mathcal{AB}}) = \left(\frac{1}{\sqrt{N}}\right)\left|\sin\left(\frac{2Et}{\sqrt{N}}\right)\right|. \tag{29}$$

This is the analytic pairwise entanglement in the analog Grover search. The evolution of the pairwise entanglement is calculated numerically and the result is shown in figure 4 along with the rate of success probability in the search algorithm. The two peak simultaneously suggesting that entanglement is indeed necessary for the discrete analogue of analog Grover search algorithm.

As can be seen from the form of the $\mathcal{C}(\rho_{\mathcal{AB}})$, the concurrence is independent of which pair of qubits we choose to keep in such a partitioning of the system. For the multipartite system, in particular, the pairwise entanglement sharing and other pairwise correlations are monogamous (see Sec. IV C); when n tends to infinity all of the pairwise correlations should vanish. The above equation clearly satisfies the same.

C. Monogamy score

The amount of quantum correlations that can be shared amongst the subsystems of a multipartite quantum state is captured by the idea of monogamy. The central idea of monogamy is that entanglement cannot be freely shared. Precisely put, *if two qubits A and B are maximally entangled then they cannot be entangled at all with a third qubit C.*

For the tripartite case, if $\mathcal{C}$ is a bipartite quantum correlation measure, then this measure is said to be monogamous (or satisfy monogamy) for a tripartite quantum state ρ_{ABC}, and with A as the "nodal observer", if

$$\mathcal{C}(\rho_{\mathcal{A}:\mathcal{BC}}) \geq \mathcal{C}(\rho_{\mathcal{AB}}) + \mathcal{C}(\rho_{\mathcal{AC}}). \tag{30}$$

Here $\mathcal{C}(\rho_{\mathcal{AB}})$ is the quantum correlation (with respect to the measure $\mathcal{C}$) between subsystems A and B, $\mathcal{C}(\rho_{\mathcal{AC}})$ is the quantum correlation between subsystems A and C, and $\mathcal{C}(\rho_{\mathcal{A}:\mathcal{BC}})$ is quantum correlation between subsystem A and subsystems B and C taken together. For example, if we have a three-qubit pure state $\rho_{ABC} = (|\Psi\rangle\langle\Psi|)_{ABC}$



(a)Figure a



(b)Figure b

FIG. 3. Rate of change of concurrence with time in (a) the analog analogue of Grover search and (b) the original Grover search



FIG. 4. Variations of the concurrence(across system partition of $A_1A_2|A_3...A_N$) and the probability of success as a function of time at parameter values $E = 1$ and $N = 4$.

and a quantum correlation measure say concurrence then it can be shown that

$$\mathcal{C}^2_{AB} + \mathcal{C}^2_{AC} \leq 4 \text{ det } \rho_A. \tag{31}$$

Even though BC is a two qubit system with a four dimensional subspace, it can be shown that the support of ρ_{BC} is spanned by the eigenvectors corresponding to at most two non-zero eigenvalues of the reduced density matrix ρ_{BC}, and hence effectively becomes a two-dimensional space. This allows to treat the bipartite split of A and BC as an effective two-qubit system whose concurrence, $\mathcal{C}_{A:BC}$, is simply given by $2\sqrt{\det \rho_A}$. Substituting the value of $\mathcal{C}_{A:BC}$ above, we get the Coffman, Kundu, Wootters (CKW) inequality [49]

$$\mathcal{C}^2_{AB} + \mathcal{C}^2_{AC} \leq \mathcal{C}^2_{A:BC}. \tag{32}$$

This leads to the concept of quantum monogamy score, which, for a given bipartite quantum correlation measure, is defined as

$$\delta\mathcal{C} \equiv \mathcal{C}(\rho_{\mathcal{A}:\mathcal{BC}}) - \mathcal{C}(\rho_{\mathcal{AB}}) - \mathcal{C}(\rho_{\mathcal{AC}}) \geq 0. \tag{33}$$

If a tripartite state satisfies entanglement monogamy then its monogamy score is positive else it is negative.

Squared concurrence satisfies a monogamy inequality for arbitrary pure and mixed states, where for mixed states, the $\mathcal{C}^2_{A:BC}$ is calculated by the minimum average concurrence squared over all possible pure state decompositions $\{p_i, |\psi_i\rangle\}$ of ρ_{BC}. Another measure of bipartite entanglement is the entanglement of formation (EoF)[40], which is closely related to two-qubit concurrence. Consider a bipartite quantum state ρ_{AB}, and the ensemble $\{p_i, |\psi_i\rangle\}$ denoting a possible pure state decomposition of ρ_{AB}, satisfying $\rho_{AB} = \sum_i p_i|\psi_i\rangle\langle\psi_i|$. The EoF is defined as

$$E_f(\rho_{AB}) = \min_{\{p_i,|\psi_i\rangle\}} \sum_i p_i S(\text{Tr}_B[|\psi_i\rangle\langle\psi_i|]), \tag{34}$$

where $S(\text{Tr}_B[|\psi_i\rangle\langle\psi_i|])$ is the von Neumann entropy of the reduced density matrix corresponding to the A part of ρ_{AB}. For a two-qubit mixed state ρ_{AB}, $E_f(\rho_{AB}) = h\left((1+\sqrt{1-\mathcal{C}^2_{AB}})/2\right)$, where $h(x) = -x\log_2 x - (1-x)\log_2(1-x)$ is the binary entropy function. The EoF, being a concave function of squared concurrence, does not obey the CKW inequality. However, the the square of the EoF does obey the same relation as the squared concurrence for tripartite systems [50].

D. Monogamy in analog Grover search

The discrete analogue of analog Grover search satisfies the squared concurrence monogamy calculated as

$$\begin{aligned} C^2(\rho_{A_1|A_2A_3...A_N}) - C^2(\rho_{A_1A_2}) - C^2(\rho_{A_1A_3}) \\ -... - C^2(\rho_{A_1A_N}) \geq 0. \end{aligned} \tag{35}$$

Since in the analog Grover search case, the concurrence between any two pair of qubits is the same, i.e., $\mathcal{C}(\rho_{ij}) = \mathcal{C}(\rho_{kl})$ $\forall$ pairs of qubits ij, kl, as a result, the monogamy score between the n qubits reduces to the following

$$\delta\mathcal{C} \equiv \mathcal{C}\left(\rho_{A:BC...}\right) - (\text{n-1})\ \mathcal{C}(\rho_{AB}). \tag{36}$$

The monogamy score is thus, given by

$$\delta\mathcal{C} = \left(\frac{N-2}{2N} - \frac{1}{N}\log_2\frac{N}{2}\right)\sin^2\left(\frac{2Et}{\sqrt{N}}\right) \tag{37}$$



FIG. 5. Variations of the monogamy score for the concurrence, the monogamy score for the squared entanglement of formation and the probability of success as a function of time at parameter values $E = 1$ and $N = 32$

$$\delta\mathcal{C} = \left(\frac{\left(-\sqrt{1-\frac{(n-2)\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{2n}}-1\right)\log\left(\frac{1}{2}\left(\sqrt{1-\frac{(n-2)\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{2n}}+1\right)\right)}{2\log(2)} \right)$$

$$-\left(-\frac{\left(\frac{1}{2}\left(-\sqrt{1-\frac{(n-2)\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{2n}}-1\right)+1\right)\log\left(\frac{1}{2}\left(-\sqrt{1-\frac{(n-2)\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{2n}}-1\right)+1\right)}{\log(2)} \right)\text{^}2$$

$$-\log_2\left(\frac{n}{2}\right)\left(-\frac{\log\left(1-\sqrt{1-\frac{4\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{\left(\sqrt{\left(\sqrt{n}-2\right)^2+\sqrt{n}+2}\right)^2}}\right)+\log\left(\sqrt{1-\frac{4\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{\left(\sqrt{\left(\sqrt{n}-2\right)^2+\sqrt{n}+2}\right)^2}}+1\right)+2\sqrt{1-\frac{4\sin^2\left(\frac{2t}{\sqrt{n}}\right)}{\left(\sqrt{\left(\sqrt{n}-2\right)^2+\sqrt{n}+2}\right)^2}}\tanh^{-1}\left(\right.}{\log(4)} \right.$$

Monogamy inequality bounds the amount of pairwise entanglement that can be shared between multiple qubits, and we have seen that the discrete analog of Grover search algorithm satisfies two monogamy inequalities for arbitrary times.

Similarly, the squared entanglement of formation satisfies a monogamy inequality, where the monogamy score is given as

V. CONCLUSION

To summarize, in this paper we have explored the role of quantum coherence and monogamy of entanglement in the discrete analogue of the analog analogue of Grover search algorithm. By using the l_1-norm and the relative entropy of coherence, it was shown that coherence acts

as a signature for the success of the analog Grover search algorithm. It was also shown that a maximally coherent state $|s\rangle = \frac{1}{\sqrt{N}} \sum_{i=1}^{i=N-1} |i\rangle$ ends up into an incoherent state $|w\rangle$ as the search algorithm evolves and hence this amount of coherence is actually consumed during the searching process. The path traversed by the time evolving vector under the analog Grover Hamiltonian follows a geodesic in the two dimensional subspace of the N dimensional Hilbert space. This in turn provides an alternate geometric proof of the optimality of the analog Grover search algorithm.

The variation of entanglement and correlations was also quantified and the analogous rate of change of concurrence between the discrete and analog Grover search algorithms suggests that our mapping preserves the original behavior of the algorithm. The pairwise entanglement was shown to peak simultaneously with the rate of success probability as evidence that entanglement is indeed necessary, for pure state implementation of analog Grover search algorithm. The n going to infinity limit of pairwise entanglement suggested a monogamous behavior of quantum correlations in the analog Grover search and it was then shown that the analog Grover search also satisfies the entanglement monogamy inequality for both entanglement measures namely the concurrence and the entanglement of formation, for all times during the search algorithm.

VI. ACKNOWLEDGMENTS

Namit Anand would like to acknowledge the hospitality of the Harish-Chandra Research Institute for allowing him to use their facilities during several visits made throughout the preparation of this manuscript.

[1] R. Feynman, International Journal Of Theoretical Physics 21, (1982).
[2] D. Deutsch, Proceedings Of The Royal Society A: Mathematical, Physical And Engineering Sciences 400, (1985).
[3] D. Deutsch and R. Jozsa, Proceedings Of The Royal Society A: Mathematical, Physical And Engineering Sciences 439, (1992).
[4] E. Bernstein and U. Vazirani, SIAM Journal On Computing 26, (1997).
[5] D. Simon, SIAM Journal On Computing 26, (1997).
[6] L. Grover, Physical Review Letters 79, (1997).
[7] P. Shor, SIAM Journal On Computing 26, (1997).
[8] C. Zalka, Physical Review A 60, (1999).
[9] Brassard, G., P. Hyer, M. Mosca, and A. Tapp, Quantum Computation and Information, AMS Contemporary Mathematics Series Vol. 305 pp. 5374 (2002); e-print arXiv:quant-ph/0005055.
[10] G. Brassard, P. Hyer and A. Tapp, SIGACT News 28, (1997).
[11] C. Drr, M. Heiligman, P. HOyer and M. Mhalla, SIAM Journal On Computing 35, (2006).
[12] Ambainis, A., and R. palek, Symposium on Theoretical Aspects of Computer Science, Lecture Notes in Computer Science Vol. 3884 (Springer, Berlin), pp. 172183 (2006); e-print arXiv:quant-ph/0508205.
[13] E. Farhi, J. Goldstone, S. Gutmann, M. Sipserhttps arXiv:quant-ph/0001106
[14] M. Born and V. Fock, Zeitschrift für Physik 51, (1928).
[15] S. Jansen, M. Ruskai and R. Seiler, J. Math. Phys. 48, (2007).
[16] D. Aharonov, W. Dam, J. Kempe, Z. Landau, S. Lloyd, O. Regev ,SIAM Journal of Computing, Vol. 37, Issue 1, p. 166-194 (2007).
[17] E. Farhi and S. Gutmann, Physical Review A 57, (1998).
[18] R. Horodecki, P. Horodecki, M. Horodecki and K. Horodecki, Reviews Of Modern Physics 81, (2009).
[19] C. Bennett, G. Brassard, C. Crpeau, R. Jozsa, A. Peres and W. Wootters, Phys. Rev. Lett. 70, (1993).
[20] C. Bennett and S. Wiesner, Phys. Rev. Lett. 69, (1992).
[21] A. K. Pati, Phys. Rev. A 63, 014303, (2000).
[22] C. Bennett, D. DiVincenzo, P. Shor, J. Smolin, B. Terhal and W. Wootters, Physical Review Letters 87, (2001).
[23] S. L. Braunstein and A. K. Pati, Quant. Info. Comput. 2, 399 (2002); A. K. Pati and S. L. Braunstein, J. of Indian Inst. of Sci., (2009).
[24] A. Ekert and R. Jozsa, Reviews Of Modern Physics 68, (1996).
[25] D. Meyer and N. Wallach, J. Math. Phys. 43, (2002).
[26] N. Lambert, Y. Chen, Y. Cheng, C. Li, G. Chen and F. Nori, Nature Physics 9, (2012).
[27] S. Huelga and M. Plenio, Contemporary Physics 54, (2013).
[28] J. Goold, M. Huber, A. Riera, L. Rio and P. Skrzypczyk, Journal Of Physics A: Mathematical And Theoretical 49, (2016).
[29] G. Gour, M. Mller, V. Narasimhachar, R. Spekkens and N. Yunger Halpern, Physics Reports 583, (2015).
[30] D. Janzing, International Journal Of Theoretical Physics 39, (2000).
[31] E. Ruch, Theoretica Chimica Acta 38, (1975).
[32] D. Meyer, Physical Review Letters 82, (1999).
[33] J. Eisert, M. Wilkens and M. Lewenstein, Physical Review Letters 83, (1999).
[34] N. Anand and C. Benjamin, Quantum Inf Process 14, (2015).
[35] M. Horodecki, P. Horodecki and J. Oppenheim, Physical Review A 67, (2003).
[36] M.B. Plenio and S. Virmani, Quantum Inf. Comput. 7, 1, (2007); e-print arXiv:quant-ph/0504163.
[37] T. Baumgratz, M. Cramer and M. Plenio, Physical Review Letters 113, (2014).
[38] B. Terhal, IBM Journal Of Research And Development 48, (2004).
[39] H. S. Dhar, A. K. Pal, D. Rakshit, A. Sen De, U. Sen e-print arXiv:1610.01069.
[40] W. Wootters, Physical Review Letters 80, 2245, (1998).
[41] A. Childs and J. Goldstone, Physical Review A 70, (2004).
[42] M. Nielsen and I. Chuang, Quantum Computation And Quantum Information (Cambridge University Press,

Cambridge, 2004).
[43] J. Thorpe, Elementary Topics In Differential Geometry (Springer-Verlag, New York, 1979).
[44] E. Chitambar, D. Leung, L. Maninska, M. Ozols and A. Winter, Commun. Math. Phys. 328, (2014).
[45] B. Coecke, T. Fritz and R. Spekkens, Information And Computation 250, (2016).
[46] M. HORODECKI and J. OPPENHEIM, International Journal Of Modern Physics B 27, (2013).
[47] M. Schlosshauer, Reviews Of Modern Physics 76, (2005).
[48] A., G. Adesso, M. B. Plenio, e-print arXiv:1609.02439.
[49] V. Coffman, J. Kundu, and W. K. Wootters, Phys. Rev. A 61, 052306 (2000).
[50] Y.-K. Bai, Y.-F. Xu, and Z. D.Wang, Phys. Rev. Lett. 113, 100503 (2014).