

Name: Namrata Bhorade

Class: BE COMPS

UID: 2018130004

Data Science, 2022

## Tut 4: Independent Component Analysis

---

Ex. 1

Exercise: Mixing statistically independent sources

Given some scalar and statistically independent random variables (signals)  $s_i$  with zero mean, unit variance, and a value  $a_i$  for the kurtosis that lies between  $-a$  and  $+a$ , with arbitrary but fixed value of  $0 < a$ . The  $s_i$  shall be mixed like

$$x := \sum_i w_i s_i$$

with constant weights  $w_i$ .

- Which constraints do you have to impose on the weights  $w_i$  to guarantee that the mixture has unit variance as well?

Hint

$$\begin{aligned} \text{var}(x) &= \langle (x - \langle x \rangle)^2 \rangle \\ &= \langle x^2 \rangle - \langle x \rangle^2 \end{aligned}$$

Data Science, 2022

## Tut 4: Independent Component Analysis

Ex 1 Mixing statistically independent sources

Answer: Variance of the mixture is given as

$$\text{Var}(x) = \langle (x - \langle x \rangle)^2 \rangle$$

$$= \langle x^2 \rangle - \langle x \rangle^2$$

$$= \left\langle \left( \sum_i w_i s_i \right)^2 \right\rangle - \left\langle \sum_i w_i s_i \right\rangle^2$$

$$= \left\langle \left( \sum_i w_i s_i \right)^2 \right\rangle - \left( \sum_i w_i \langle s_i \rangle \right)^2$$

$$= \left\langle \left( \sum_i w_i s_i \right) \left( \sum_j w_j s_j \right) \right\rangle - \left( \sum_i w_i \langle s_i \rangle \right) \left( \sum_j w_j \langle s_j \rangle \right)$$

$$= \left\langle \sum_{i,j} w_i w_j s_i s_j \right\rangle - \sum_{i,j} w_i w_j \langle s_i \rangle \langle s_j \rangle$$

$$= \sum_{i,j} w_i w_j \langle s_i s_j \rangle - \sum_{i,j} w_i w_j \langle s_i \rangle \langle s_j \rangle$$

$$= \sum_i w_i w_i (\langle s_i s_i \rangle - \langle s_i \rangle \langle s_i \rangle) +$$

$$\sum_{i,j: i \neq j} w_i w_j (\langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle)$$

$$= \sum_i w_i^2 (\langle s_i^2 \rangle - \langle s_i \rangle^2) + \sum_{i,j: i \neq j} w_i w_j (\langle s_i s_j \rangle - \langle s_i \rangle \langle s_j \rangle)$$

$s_i$  &  $s_j$  are statistically independent for  $i \neq j \Rightarrow \langle s_i \rangle \langle s_j \rangle - \langle s_i \rangle \langle s_j \rangle = 0$

Also  $\text{Var}(s_i) = 1$

$$\therefore \text{Var}(x) = \sum_i w_i^2$$

To guarantee that the mixture has unit variance,

$$\text{var}(x) = 1$$


$$\therefore \sum_i w_i^2 = 1$$

$\therefore$  The following constraint has to be imposed on the weights  $w_i$  for the mixture to have unit variance

$$\boxed{\sum_i w_i^2 = 1}$$

## Ex.2


Two examples of joint probability densities are shown in following figure. One is a mixture of arbitrary non-Gaussian densities, and the other one a mixture of Gaussians. The dashed curves around the densities plot the projected variance measured in all directions. The dashed line marks the direction of maximum variance, that is, the first principal component. Similarly, the values of kurtosis are shown using solid curves and the direction of maximum kurtosis with a solid line.



Example joint probability densities. (a) For non-Gaussian densities the principal (dashed line) and independent (solid line) directions can be identified, whereas (b) for Gaussian ones the directions are all equal. The corresponding dashed and solid curves show the values of variance and kurtosis in all directions respectively.

Referring to above provide the guess independent components and distributions from data


- Decide whether the following distributions can be linearly separated into independent components. If yes,
- sketch the (not necessarily orthogonal) axes onto which the data must be projected to extract the independent components. Draw these axes also the marginal distributions of the corresponding components.






Ex 2 Guess independent components and distribution from data


a)



b)



c)



Cannot be separated into independent components.