

A method for recursively generating sequential rational approximations to $\sqrt[n]{k}$

Joe Nance
nance2uiuc@gmail.com
Department of Mathematics
University of Illinois at Urbana-Champaign

November 23, 2010

Abstract

The goal of this paper is to derive a simple recursion that generates a sequence of fractions approximating $\sqrt[n]{k}$ ($n \in \mathbb{N}$) with increasing accuracy. The recursion is ~~then~~ defined in terms of a series of first-order non-linear difference equations and then analyzed as a discrete dynamical system. Convergence behavior is then discussed in the language of initial trajectories and eigenvectors, effectively proving convergence without notions from standard analysis of infinitesimals.

1 Introduction and motivation

Consider for a moment the simple recursion

$$\frac{x_{t+1}}{y_{t+1}} = \frac{x_t + 2y_t}{x_t + y_t}.$$

should the reader immediately recognize that this heads off (1) to $\sqrt{2}$?

(Choose initial values to set it marching on its way towards $\sqrt{2}$.) If we choose initial values $x_0 = y_0 = 1$, the recursion (1) gives a sequence of fractions approximating $\sqrt{2}$ whose whose behavior is summarized in the table below.

| t | $\frac{x_t}{y_t}$ | \approx |
|---|-------------------|------------|
| 0 | $\frac{1}{1}$ | 1 |
| 1 | $\frac{3}{2}$ | 1.5 |
| 2 | $\frac{7}{5}$ | 1.4 |
| 3 | $\frac{17}{12}$ | 1.4167... |
| 4 | $\frac{41}{29}$ | 1.41379... |
| 5 | $\frac{99}{70}$ | 1.41429... |

You might hold off on revealing $\sqrt{2}$ as the limit until after this table, then remark that the terms seem to head towards $\sqrt{2}$. Then you give a justification.

For reassurance that the recursion generates a sequence that does in fact converge to $\sqrt{2}$, employ the following analysis: For some sequence a_t , $a_t \rightarrow L$ then $a_{t+1} \rightarrow L$ as well. So suppose that the recursion (1) has limit L . Multiply the top and bottom of the right hand side of (1) by $\frac{1}{y_t}$ and we get

$$\frac{x_{t+1}}{y_{t+1}} = \frac{x_t + 2y_t}{x_t + y_t} \frac{\frac{1}{y_t}}{\frac{1}{y_t}} = \frac{\frac{x_t}{y_t} + 2}{\frac{x_t}{y_t} + 1}$$

and using the previous fact about limits, we have

$$L = \frac{L + 2}{L + 1}.$$

you should make sure to tell the reader that you're using $\frac{x_t}{y_t}$ as your sequence. This will be resolved if you make this more clear

by which you mean $\lim \frac{x_{t+1}}{y_{t+1}}$.

use words or include \lim

Incomplete sentence

why is this unsettling?

Which gives $L = \pm\sqrt{2}$. In this analysis, we have picked up the unsettling possibility of this recursion generating a sequence of fractions converging to $-\sqrt{2}$. Discussion of which initial values generate such a sequence is withheld momentarily.

2 \sqrt{k} Recursion

It is convenient to consider recursion (1) as an action on a system of first order linear difference equations given by

$$\begin{cases} x_{t+1} = x_t + 2y_t \\ y_{t+1} = x_t + y_t \end{cases} \quad (2)$$

which is a discrete dynamical system, or "DDS". Clearly, if we replace 2 by a positive integer $k \in \mathbb{Z}^+$ we obtain a recursion similar in structure to recursion (1) converging to $\pm\sqrt{k}$ for any initial values x_0, y_0 . This recursion is

$$\frac{x_{t+1}}{y_{t+1}} = \frac{x_t + ky_t}{x_t + y_t} \quad (3)$$

and gives a corresponding DDS

$$\begin{cases} x_{t+1} = x_t + ky_t \\ y_{t+1} = x_t + y_t \end{cases} \quad (4)$$

which can be represented in matrix form as

$$\begin{pmatrix} x_{t+1} \\ y_{t+1} \end{pmatrix} = \begin{pmatrix} 1 & k \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x_t \\ y_t \end{pmatrix} \quad (5)$$

and any term in a sequence generated by (5) is generalized as

$$\begin{pmatrix} x_t \\ y_t \end{pmatrix} = \begin{pmatrix} 1 & k \\ 1 & 1 \end{pmatrix}^t \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}. \quad (6)$$

Recursion (3) is recovered by taking ratios of terms with equal indices.

long sentence. lots of displayed equations. Do you need them all?

3 $\sqrt[n]{k}$ Recursion

A natural question at this point would be "Does there exist a structurally simple recursion similar to (3) generating a sequence of fractions approximating $\sqrt[n]{k}$?" To which the answer is "kind of". In the spirit of the previous analysis, start out with

$$L = \sqrt[n]{k} \Rightarrow L^n = k$$

and add L to both sides to get

$$L^n + L = L + k$$

then factor out an L

$$L(L^{n-1} + 1) = L + k$$

now divide both sides by $L^{n-1} + 1$ to get

$$L = \frac{L + k}{L^{n-1} + 1}$$

now use the fact about sequences to obtain

$$\frac{x_{t+1}}{y_{t+1}} = \frac{x_t + ky_t}{\frac{x_t^{n-1}}{y_t^{n-2}} + y_t} \quad (7)$$

which has the corresponding DDS

$$\begin{cases} x_{t+1} &= x_t + ky_t \\ y_{t+1} &= \frac{x_t^{n-1}}{y_t^{n-2}} + y_t \end{cases} \quad (8)$$

which is not representable as a simple 2×2 matrix with real entries for arbitrary n . This is because there are now *three* different terms in this system: x_t , y_t , and $\frac{x_t^{n-1}}{y_t^{n-2}}$ which is non-linear.

4 Convergence behavior of the DDS given by Π_n

We are now in a position to analyze (8) using matrix methods by force! Suppose there *did exist* an $n \times n$ matrix, call it Π_n such that (8) could be

Incomplete surface.

represented as

$$\begin{pmatrix} \vec{x}_{t,1} \\ \vec{x}_{t,2} \\ \vdots \\ \vec{x}_{t,n} \end{pmatrix} = \Pi_n^t \begin{pmatrix} \vec{x}_{0,1} \\ \vec{x}_{0,2} \\ \vdots \\ \vec{x}_{0,n} \end{pmatrix} \quad (9)$$

where $\vec{x}_{a,b}$ denotes the b^{th} entry of the vector corresponding to a . Or equivalently, $\vec{R}_t = \Pi_n^t \vec{R}_0$. But what would the matrix Π_n look like? Consider the properties of (8) that Π_n must capture in order to faithfully represent convergence behavior of (7). First, recognize that the action on (8) needed to arrive at (7) is taking ratios of terms with equal indices which is equivalent to taking the ratio of successive entries of a vector \vec{R}_t . The entries in an arbitrary vector \vec{R}_t must tend toward those of the dominant eigenvector of Π_n , $\vec{\lambda}_d$. So the ratio of any two successive entries in $\vec{\lambda}_d$ should be equal to $\sqrt[n]{k}$. That is, $\frac{\vec{\lambda}_{d,i}}{\vec{\lambda}_{d,i+1}} = \sqrt[n]{k}$ where $1 \leq i \leq n-1$ and $\vec{\lambda}_{d,i}$ is the i^{th} entry of the dominant eigenvector of Π_n . Such an eigenvector would look like $\vec{\lambda}_d = \begin{pmatrix} \sqrt[n]{k}^{n-1} & \sqrt[n]{k}^{n-2} & \dots & 1 \end{pmatrix}^T$. Recovery of the dominant eigenvalue from the dominant eigenvector is had by solving for λ_d in $(\Pi_n - \lambda_d I_n) \vec{\lambda}_d = \vec{0}$ where $\vec{0}$ is the column vector consisting entirely of zeros. This calculation yields $\lambda_d = 1 + \sqrt[n]{k}$.

does it? how when we don't know what Π_n is yet?

Now suppose Π_n is diagonalizable. Then Π_n has n distinct eigenvectors corresponding to n distinct eigenvalues. This means that Π_n admits a basis for \mathbb{R}^n consisting entirely of eigenvectors of Π_n . So any initial vector \vec{R}_0 in \mathbb{R}^n can be written as a linear combination of eigenvectors $\vec{R}_0 = \sum_{i=1}^n c_i \vec{\lambda}_i$.

Applying Π_n to our initial vector \vec{R}_0 ,

$$\begin{aligned} \vec{R}_1 &= \Pi_n \vec{R}_0 = \Pi_n \sum_{i=1}^n c_i \vec{\lambda}_i \\ &= c_1 \Pi_n \vec{\lambda}_1 + \dots + c_n \Pi_n \vec{\lambda}_n \\ &= c_1 \lambda_1 \vec{\lambda}_1 + \dots + c_n \lambda_n \vec{\lambda}_n \end{aligned}$$

diagonalizable \Rightarrow

distinct eigenvalues

diagonalizable \Rightarrow basis of eigenvectors \checkmark

this is confusing as it stands, but it's a critical statement in your paper. work hard to make this 100% crystal clear.

Applying Π_n again,

$$\begin{aligned}\vec{R}_2 &= \Pi_n^2 \vec{R}_0 = \Pi_n(\Pi_n \vec{R}_0) = \Pi_n(c_1 \lambda_1 \vec{\lambda}_1 + \cdots + c_n \lambda_n \vec{\lambda}_n) \\ &= c_1 \lambda_1 \Pi_n \vec{\lambda}_1 + \cdots + c_n \lambda_n \Pi_n \vec{\lambda}_n \\ &= c_1 \lambda_1^2 \vec{\lambda}_1 + \cdots + c_n \lambda_n^2 \vec{\lambda}_n.\end{aligned}$$

We can see the pattern now,

$$\vec{R}_t = \Pi_n^t \vec{R}_0 = c_1 \lambda_1^t \vec{\lambda}_1 + \cdots + c_n \lambda_n^t \vec{\lambda}_n = \sum_{i=1}^n c_i \lambda_i^t \vec{\lambda}_i. \quad (10)$$

Verify that indeed,

funny wording.

$$\vec{R}_t = \Pi_n^t \vec{R}_0 = \begin{pmatrix} c_1 \lambda_1^t \vec{\lambda}_{1,1} + \cdots + c_n \lambda_n^t \vec{\lambda}_{n,1} \\ c_1 \lambda_1^t \vec{\lambda}_{1,2} + \cdots + c_n \lambda_n^t \vec{\lambda}_{n,2} \\ \vdots \\ c_1 \lambda_1^t \vec{\lambda}_{1,n} + \cdots + c_n \lambda_n^t \vec{\lambda}_{n,n} \end{pmatrix}$$

where $\vec{\lambda}_{a,b}$ denotes the b^{th} entry of the eigenvector corresponding to λ_a .

Note that with increasing t , only one of $c_i \lambda_i^t \vec{\lambda}_{i,l}$ terms becomes dominant term. The dominant term is the one involving the dominant eigenvalue, λ_d and the entries of its corresponding eigenvector, $\vec{\lambda}_{d,i}$. Since contributions of the other terms become negligible in the quotient, we can make the statement

$$\vec{R}_t = \Pi_n^t \vec{R}_0 \approx c \lambda_d^t \vec{\lambda}_d = c(1 + \sqrt[n]{k})^t \begin{pmatrix} (\sqrt[n]{k})^{n-1} \\ (\sqrt[n]{k})^{n-2} \\ \vdots \\ 1 \end{pmatrix} \text{ for } t \gg 1.$$

I'm worried about this

It follows that

$$\frac{\vec{R}_{t,i}}{\vec{R}_{t,i+1}} = \frac{c(1 + \sqrt[n]{k})^t \vec{\lambda}_{d,i} + \cdots + c_n \lambda_n^t \vec{\lambda}_{n,i}}{c(1 + \sqrt[n]{k})^t \vec{\lambda}_{d,i+1} + \cdots + c_n \lambda_n^t \vec{\lambda}_{n,i+1}} \approx \frac{c(1 + \sqrt[n]{k})^t \vec{\lambda}_{d,i}}{c(1 + \sqrt[n]{k})^t \vec{\lambda}_{d,i+1}} = \frac{\vec{\lambda}_{d,i}}{\vec{\lambda}_{d,i+1}} = \sqrt[n]{k} \quad (11)$$

for $(1 \leq i \leq n-1)$ and $t \gg 1$.

When $n = 2$ (finding square roots), let $c = 0$ and notice that (11) is approximately $-\sqrt{k}$. To settle our curiosity from Section 1, if we are to

awkward wording

how do you know λ_d is dominant term?

I'm happy w/ this, provided $c \neq 0$.

have a sequence generated by (2) converging to $-\sqrt{k}$ then the appropriate initial values are ones which are components of some multiple of the second eigenvector, not a linear combination of the dominant eigenvector and the other. This way, ratios of successive entries of an evolving vector equal the slope of the second eigenvector, $-\sqrt{k}$. If we restrict our choices of initial values to \mathbb{Q}^n then we do not run into this problem of multiple limits.

Q2

why? what is second eigenvector, and why isn't it in Q2?

5 Derivation of the matrix Π_n

In the preceding section we showed that if (8) can be represented as (9), then $\lim_{t \rightarrow +\infty} \frac{x_{t,i}}{x_{t,i+1}} = \sqrt[n]{k}$, as desired. It only remains to find the exact form for Π_n . In the analysis above, we showed that long-term time evolution of an initial vector depends heavily on "hitting" $\vec{\lambda}_d$ with its corresponding eigenvalue λ_d , which in turn depends on left-multiplying \vec{R}_0 by Π_n . Carrying this calculation out gives

$$\lambda_d \vec{\lambda}_d = \left(1 + \sqrt[n]{k}\right) \begin{pmatrix} \left(\sqrt[n]{k}\right)^{n-1} \\ \left(\sqrt[n]{k}\right)^{n-2} \\ \left(\sqrt[n]{k}\right)^{n-3} \\ \left(\sqrt[n]{k}\right)^{n-4} \\ \vdots \\ 1 \end{pmatrix} = \begin{pmatrix} \left(\sqrt[n]{k}\right)^{n-1} \\ \left(\sqrt[n]{k}\right)^{n-2} \\ \left(\sqrt[n]{k}\right)^{n-3} \\ \left(\sqrt[n]{k}\right)^{n-4} \\ \vdots \\ 1 \end{pmatrix} + \begin{pmatrix} k \\ \left(\sqrt[n]{k}\right)^{n-1} \\ \left(\sqrt[n]{k}\right)^{n-2} \\ \left(\sqrt[n]{k}\right)^{n-3} \\ \vdots \\ \sqrt[n]{k} \end{pmatrix}$$

Evidently, Π_n is such that when an initial vector \vec{R}_0 is left multiplied by it, returned is the dominant eigenvector plus another vector, call it \vec{v} , along with other insignificant terms in the limit that $t \gg 1$. This means that Π_n must be the sum of two matrices acting on the linear combination of eigenvectors that comprises \vec{R}_0 . $\Pi_n \vec{R}_0 = (I_n + \pi_n) (\lambda_d \vec{\lambda}_d + \dots + \lambda_n \vec{\lambda}_n)$ where I_n is the n -dimensional identity matrix. But what is π_n ? Consider what action is taken by π_n to return \vec{v} from λ_d . Apparently, π_n is an $n \times n$ matrix such that when it left multiplies a column vector, it has the effect of scalar multiplication by $\sqrt[n]{k}$. By inspection, we see that $\pi_n =$

awkward wording

What?

incomplete sentence. should explain why you picked I_n as "first" matrix.

if Π_n is scalar mult by $\sqrt[n]{k}$,
 then $\Pi_n = \begin{pmatrix} \sqrt[n]{k} & & \\ & \sqrt[n]{k} & \\ & & \dots \\ & & & \sqrt[n]{k} \end{pmatrix}$
 but your Π_n isn't this!

$$\begin{pmatrix} 0 & 0 & 0 & \dots & 0 & k \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \end{pmatrix}. \text{ The exact form of } \Pi_n \text{ is given by}$$

$$\Pi_n = I_n + \pi_n = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & k \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & k \\ 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 1 \end{pmatrix}.$$

This is critical, so
 deserves some kind
 of explanation.

6 Algebraic properties of the matrix Π_n

We must still confirm diagonalizability of Π_n since most of our case depends upon this property of Π_n . The characteristic polynomial, $P(\lambda)$, of Π_n can be found by computing $\det(\Pi_n - \lambda I_n)$ by expanding in minors along the top row, giving $P(\lambda) = (1 - \lambda)^n + (-1)^{n+1}k$. The eigenvalues are had by solving $P(\lambda) = 0$ giving $\lambda_j = 1 - \sqrt[n]{k} e^{i\frac{J\pi}{n}}$ where $1 \leq j \leq n$ and $J = 2j$ when n is even and $2j+1$ when n is odd. The largest of these eigenvalues is $\lambda_d = \lambda_{\frac{n}{2}} = \lambda_{\frac{n-1}{2}} = 1 + \sqrt[n]{k}$ as desired. These n distinct eigenvalues give n

distinct eigenvectors given by $\vec{\lambda}_j = \left(\frac{k}{\lambda_j - 1}, \frac{k}{(\lambda_j - 1)^2}, \frac{k}{(\lambda_j - 1)^3}, \frac{k}{(\lambda_j - 1)^4}, \dots, 1 \right)^T$,

the largest of which corresponds to the eigenvalue λ_d , this vector is $\vec{\lambda}_d = \left((\sqrt[n]{k})^{n-1}, (\sqrt[n]{k})^{n-2}, (\sqrt[n]{k})^{n-3}, (\sqrt[n]{k})^{n-4}, \dots, 1 \right)^T$. Since Π gives n distinct eigenvalues and eigenvectors, Π_n is diagonalizable. We are now in the position to make the generalization of (7) as follows

$$\Pi_n^t \vec{R}_0 = \vec{R}_t \Rightarrow \lim_{t \rightarrow +\infty} \frac{\vec{R}_{t,i}}{\vec{R}_{t,i+1}} = \sqrt[n]{k} \text{ for } (1 \leq i \leq n-1; t \in \mathbb{Z}) \quad (12)$$


careful here.
 it makes sense
 to talk about

awkward wording.




or equivalently

$$\begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & k \\ 1 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 1 \end{pmatrix}^t \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \\ x'_4 \\ \vdots \\ x'_n \end{pmatrix} \Rightarrow \lim_{t \rightarrow +\infty} \frac{x'_i}{x'_{i+1}} = \sqrt[n]{k}.$$

This result effectively fulfils the goal of the paper which was to derive a simple recursion that generates a sequence of fractions approximating $\sqrt[n]{k}$ with increasing accuracy. Below are two graphical representations of the recursive process for $n = 2, k = 2$ and $n = 3, k = 2$.



plot 2D.jpg



plot 3D.jpg

are there better
caption names?

7 Computation

The result of the previous section satisfies the technical goal of this paper, but it is left to the reader to judge the practicality of this result. Accuracy of an approximation depends on taking powers of an $n \times n$ matrix. This tedious task can be tiresome for even relatively small powers of n and t . So where do we look to find aid in this computation? One could certainly start with the Cayley-Hamilton Theorem, which states that every $n \times n$ matrix over a commutative ring satisfies its own characteristic equation, $P(\lambda) = \det(\lambda I_n - A)$. Applying this theorem to Π_n gives

$$P(\Pi_n) = (I_n - \Pi_n)^n + (-1)^{n+1}k I_n = \mathbf{0}$$

where $\mathbf{0}$ is the $n \times n$ matrix consisting entirely of zeros. The binomial theorem then gives

$$\sum_{i=0}^n \frac{n!(-1)^i}{i!(n-i)!} \Pi_n^i I_n = (-1)^n k I_n$$

and solving for Π_n^n gives

$$\Pi_n^n = \sum_{i=1}^{n-1} \frac{n!(-1)^{n-1-i}}{i!(n-i)!} \Pi_n^i + [(-1)^{n-1} + k] I_n \quad (13)$$

which is an explicit equation expressing Π_n^n in terms of lower powers of Π_n and I_n . This is useful because if one is able to calculate powers of Π_n up to and including Π_n^{n-1} , then one is able to generate arbitrarily large powers of Π_n iteratively which then can be used to generate arbitrarily close approximations to $\sqrt[n]{k}$.

Let's take a look at the $n = 2$ case. Equation (13) gives

$$\Pi_2^2 = 2\Pi_2 + (k - 1)I_2 \quad (14)$$

which is an explicit expression of Π_2 in first powers of Π_2 and I_2 . Because no higher powers of Π_2 need to be calculated to arrive at (14), arbitrary

integer powers of Π_2 are gotten with ease from iterative multiplication and substitution of powers of Π_2 . This gives rise to a Fibonacci-like sequence in the exponents of Π_2 :

$$\begin{aligned}
 \Pi_2^2 &= 2\Pi_n + (k-1)I_2 \\
 \Pi_2^3 &= \Pi_2\Pi_2^2 \\
 &= 2\Pi_2^2 + (k-1)\Pi_2 \\
 &= (k+3)\Pi_2 + 2(k-1)I_2 \\
 \Pi_2^5 &= \Pi_2^3\Pi_2^2 \\
 &= (k^2 + 10k + 5)\Pi_2 + 4(k^2 - 1)I_2 \\
 &\vdots \\
 \Pi_2^{F_i} &= \Pi_2^{F_{i-1}}\Pi_2^{F_{i-2}}
 \end{aligned}$$

The reader is encouraged to try this for the $n = 3, 4, 5, \dots$ cases to see that once harrowing computations are done to make Π_n^{n-1} known, precise approximate computation soon follows.

I'm not sure what this means.