

INFORMATION LEAFLET:

VITAMIN B₁₂ solution for injections 500 γ

5 ml

VITAMIN B₁₂ solutio pro injectionibus 500 γ

5 ml

For large and small ruminants, pigs, swines, dogs and hens

NAME AND PERMANENT ADDRESS OF MARKETING AUTHORIZATION HOLDER AND MANUFACTURER, IF DIFFERENT

Marketing authorization holder and manufacturer:

VETPROM AD, Bulgaria, Radomir 2400, 26 Otetz Paisiy Street

NAME OF THE VETERINARY MEDICINAL PRODUCT

VITAMIN B₁₂ solution for injections 500 g / VITAMIN B₁₂ solutio pro injectionibus 500 γ

CONTENTS OF THE ACTIVE SUBSTANCE(S) AND EXCIPIENT(S)

5 ml of the medicinal product contain 0.0005 g Cyanocobalamin as an active substance and sodium chloride and water for injections as excipients.

THERAPEUTIC INDICATIONS

In cases of hyperchromatic, macrocytic and hypoplastic anemias; secondary anemias due to pregnancy, helminthoses, haemosporidiosis; in heavy metal intoxications, benzene and organochlorine insecticides products; for therapy and prevention of liver degeneration, neuritis, myelitis, neuralgia; radiation sickness; at feed poisoning; for improvement of digestion; as a stimulating growth factor in adolescent animals; for increased food consumption; at runts.

CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substance or to any of the excipients!

UNDESIRABLE ADVERSE REACTIONS

Not known.

ANIMAL SPECIES INDICATED FOR TREATMENT WITH THE VMP

For large and small ruminants, pigs, swines, dogs and hens

DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

Dosage: Average dose 2 γ/kg:

Cattle and buffaloes - 500 γ-0.001 /5-10 ml of 0.01 % solution/;

Calves - 150-200 γ /1.5-2 ml of 0.01 % solution /;

Sheep, goats, swine - 100-300 γ /1-3 ml of 0.01 % solution /;

Pigs - 10-50 γ /0.5-2.5 ml of 0.002 % solution /;

Dogs - 15-30 γ /0.75-1.5 ml of 0.002 % solution /;

Hens - 2-5 γ /0.1-0.25 ml of 0.002 % solution /.

In cases of severe inflammatory diseases of large animals 0.1-0.2, occasionally 0.3-0.5, for pigs up to 500 γ can be injected for antiallergic and antiinflammatory effect.

Route of administration:

Intramuscularly and subcutaneously.

ADVICE ON PROPER ADMINISTRATION

Always use a vitamin B12 injection solution just as you have been told your veterinarian. If you are unsure, ask your veterinarian.

WITHDRAWAL PERIOD

Not required

11. SPECIAL STORAGE PRECAUTIONS

Store at temperature below 25°C. Keep away from light.

Store in a dry place.

Do not use this product after the expiry date stated on the label

Keep out of children

Shelf life after first opening: 28 days. Store in refrigerator (2°C – 8 °C).

SPECIAL PRECAUTIONS

None

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUTILIZED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste should be disposed in accordance with local regulations and guidelines

DATE OF LATEST REVISION OF THE TEXT

August 2010.

ADDITIONAL INFORMATION

For any information about this veterinary product, please contact the local representative of the Marketing Authorisation Holder

VETPROM AD,

Bulgaria, Radomir 2400, 26 Otetz Paisiy Street

Tel.: +359777/8-24-93, 8-02-68; Fax +359777/8-23-91

E-mail: office@vetprom.bg