

Published in final edited form as:

Stem Cells. 2014 January ; 32(1): 85–92. doi:10.1002/stem.1525.

Partial Proteasome Inhibitors Induce Hair Follicle Growth by Stabilizing β -catenin

Gozde Yucel^a, John Van Arnem^a, Paula Casey Means^a, Erik Huntzicker^a, Banu Altindag^a,
Maria Fernanda Lara^a, Jenny Yuan^b, Calvin Kuo^b, and Anthony E. Oro^a

^aProgram in Epithelial Biology, Stanford University, School of Medicine, Stanford, California, USA

^bDepartment of Medicine, Stanford University, School of Medicine, Stanford, California, USA

Abstract

The activation of tissue stem cells from their quiescent state represents the initial step in the complex process of organ regeneration and tissue repair. While the identity and location of tissue stem cells are becoming known, how key regulators control the balance of activation and quiescence remains mysterious. The vertebrate hair is an ideal model system where hair cycling between growth and resting phases is precisely regulated by morphogen signaling pathways, but how these events are coordinated to promote orderly signaling in a spatial and temporal manner remains unclear. Here, we show that hair cycle timing depends on regulated stability of signaling substrates by the ubiquitin-proteasome system. Topical application of partial proteasomal inhibitors (PaPIs) inhibits epidermal and dermal proteasome activity throughout the hair cycle. PaPIs prevent the destruction of the key anagen signal β -catenin, resulting in more rapid hair growth and dramatically shortened telogen. We show that PaPIs induce excess β -catenin, act similarly to the GSK3 β antagonist LiCl, and antagonize Dickkopf-related protein-mediated inhibition of anagen. PaPIs thus represent a novel class of hair growth agents that act through transiently modifying the balance of stem cell activation and quiescence pathways.

Keywords

Hair growth; Stem cells; Wnt signaling; Proteasome; Tissue regeneration

Introduction

Despite extensive interest in the identification of stem cells and their location within vertebrate tissues, a key remaining issue is the nature of signals that regulate the activation

© AlphaMed Press 2013

Correspondence: Anthony Oro, M.D., Ph.D., Stanford University, Program in Epithelial Biology, 269 Campus Drive, CCSR 2145, Stanford, California 94305, USA. Telephone: 650-723-7843; Fax: 650-723-8762; Oro@stanford.edu.

Author Contributions: G.Y., J.V.A., and P.C.M.: conception, collection of data, data analysis, and manuscript writing; E.H.: conception, collection of data, and data analysis; B.A. and M.F.L.: collection and assembly of data; J.Y. and C.K.: provision of study materials; A.E.O.: conception, data analysis, manuscript writing, and financial support. G.Y., J.V.A., and P.C.M. share equal first authorship.

Disclosure of Potential Conflicts of Interest: The authors indicate no potential conflicts of interest.

See www.StemCells.com for supporting information available online.