

SCIENTIFIC REPORTS

OPEN

Isolation and characterization of myogenic precursor cells from human cremaster muscle

Neia Naldaiz-Gastesi^{1,2,3}, María Goicoechea^{2,15}, Isabel M.-a Aragón^{4,5}, Virginia Pérez-López¹, Sandra Fuertes-Alvarez¹, Bernardo Herrera-Imbroda^{4,5}, Adolfo López de Munain^{2,3,6,7}, Resi de Luna-Díaz⁸, Pedro M. Baptista^{9,10,11,12}, M. Alejandro Fernández¹³, **María Fernanda Lara^{4,5} & Ander Izeta^{1,14}**

Human myogenic precursor cells have been isolated and expanded from a number of skeletal muscles, but alternative donor biopsy sites must be sought after in diseases where muscle damage is widespread. Biopsy sites must be relatively accessible, and the biopsied muscle dispensable. Here, we aimed to histologically characterize the cremaster muscle with regard number of satellite cells and regenerative fibres, and to isolate and characterize human cremaster muscle-derived stem/precursor cells in adult male donors with the objective of characterizing this muscle as a novel source of myogenic precursor cells. Cremaster muscle biopsies (or adjacent non-muscle tissue for negative controls; N = 19) were taken from male patients undergoing routine surgery for urogenital pathology. Myosphere cultures were derived and tested for their *in vitro* and *in vivo* myogenic differentiation and muscle regeneration capacities. Cremaster-derived myogenic precursor cells were maintained by myosphere culture and efficiently differentiated to myotubes in adhesion culture. Upon transplantation to an immunocompromised mouse model of cardiotoxin-induced acute muscle damage, human cremaster-derived myogenic precursor cells survived to the transplants and contributed to muscle regeneration. These precursors are a good candidate for cell therapy approaches of skeletal muscle. Due to their location and developmental origin, we propose that they might be best suited for regeneration of the rhabdosphincter in patients undergoing stress urinary incontinence after radical prostatectomy.

In striated muscle, adult myogenic stem cells are known as satellite cells, due to their superficial position on muscle fibres¹. The myogenic process is a multifaceted transition between precursor states (quiescence, activation, proliferation and differentiation) that precede fusion of the myoblasts to regenerative muscle fibres². Besides, satellite cells reside in a complex niche, which includes other precursors such as fibro-adipogenic precursor cells (FAPs) that modulate the regenerative response³, along with signals arising from nerve and capillary terminals and other interstitial cells. For cell-based therapeutic purposes, it would thus be desirable to obtain and characterize the diverse types of human muscle precursor cells from an accessible source.

¹Tissue Engineering group, Instituto Biodonostia, San Sebastian, Spain. ²Neuromuscular diseases group, Instituto Biodonostia, San Sebastian, Spain. ³Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED), Instituto de Salud Carlos III, Madrid, Spain. ⁴CNIO-IBIMA Genitourinary Cancer Research Unit, Institute of Biomedical Research in Málaga (IBIMA), Málaga, Spain. ⁵Urology Department, Hospital Universitario Virgen de la Victoria, Málaga, Spain. ⁶Department of Neurosciences, Faculty of Medicine and Dentistry, UPV-EHU, San Sebastian, Spain. ⁷Department of Neurology, Hospital Universitario Donostia, San Sebastian, Spain. ⁸Department of Surgery, Hospital Universitario Virgen de la Victoria, Málaga, Spain. ⁹Health Research Institute of Aragón (IIS Aragón), Zaragoza, Spain. ¹⁰Centro de Investigación Biomédica en Red de Enfermedades Hepáticas y Digestivas (CIBEREHD), Instituto de Salud Carlos III, Madrid, Spain. ¹¹Health Research Institute of Jiménez Díaz Foundation (IIS FJD), Madrid, Spain. ¹²Biomedical and Aerospace Engineering Department, University Carlos III of Madrid, Madrid, Spain. ¹³Digital industry, Product design, IDONIAL Technology Center, Gijón, Spain. ¹⁴Department of Biomedical Engineering and Science, School of Engineering, **Tecnun-University of Navarra, San Sebastian, Spain.**

¹⁵Present address: Viralgen Vector Core, San Sebastian, Spain. Correspondence and requests for materials should be addressed to M.F.L. (email: mf.lara@fimabis.org) or A.I. (email: ander.izeta@biodonostia.org)