

Published in final edited form as:

*J Control Release*. 2013 March 28; 166(3): 211–219. doi:10.1016/j.jconrel.2012.12.030.

## Gene silencing following siRNA delivery to skin via coated steel microneedles: *in vitro* and *in vivo* proof-of-concept

Rosalind H.E. Chong<sup>a,b</sup>, Emilio Gonzalez-Gonzalez<sup>c,1</sup>, **Maria F. Lara<sup>d</sup>**, Tycho J. Speaker<sup>d</sup>, Christopher H. Contag<sup>c,e</sup>, Roger L. Kaspar<sup>c,d</sup>, Sion A. Coulman<sup>a</sup>, Rachel Hargest<sup>b</sup>, and James C. Birchall<sup>a,\*</sup>

<sup>a</sup>School of Pharmacy and Pharmaceutical Sciences, Cardiff University, Cardiff, CF10 3NB, UK.

<sup>b</sup>Academic Department of Surgery, University Hospital of Wales, Cardiff University, Cardiff, CF14 4XN, United Kingdom

<sup>c</sup>Department of Pediatrics, Stanford School of Medicine, and Molecular Imaging Program at Stanford, California, USA

<sup>d</sup>Transderm Inc., Santa Cruz, California, USA.

<sup>e</sup>Departments of Radiology and Microbiology & Immunology, Stanford School of Medicine, Stanford, California, USA

### Abstract

The development of siRNA-based gene silencing therapies has significant potential for effectively treating debilitating genetic, hyper-proliferative or malignant skin conditions caused by aberrant gene expression. To be efficacious and widely accepted by physicians and patients, therapeutic siRNAs must access the viable skin layers in a stable and functional form, preferably without painful administration. In this study we explore the use of minimally-invasive steel microneedle devices to effectively deliver siRNA into skin. A simple, yet precise microneedle coating method permitted reproducible loading of siRNA onto individual microneedles. Following recovery from the microneedle surface, lamin A/C siRNA retained full activity, as demonstrated by significant reduction in lamin A/C mRNA levels and reduced lamin A/C protein in HaCaT keratinocyte cells. However, lamin A/C siRNA pre-complexed with a commercial lipid-based transfection reagent (siRNA lipoplex) was less functional following microneedle coating. As Accell-modified “self-delivery” siRNA targeted against CD44 also retained functionality after microneedle coating, this form of siRNA was used in subsequent *in vivo* studies, where gene silencing was determined in a transgenic reporter mouse skin model. Self-delivery siRNA targeting the reporter (luciferase/GFP) gene was coated onto microneedles and delivered to mouse footpad. Quantification of reporter mRNA and intravital imaging of reporter expression in the outer skin layers confirmed functional *in vivo* gene silencing following microneedle delivery of siRNA. The use of coated metal microneedles represents a new, simple, minimally-invasive, patient-friendly and potentially self-administrable method for the delivery of therapeutic nucleic acids to the skin.

© 2012 Elsevier B.V. All rights reserved.

\*Corresponding author: birchalljc@cf.ac.uk Phone: +44 2920875815.

<sup>1</sup>Current address: The Andalusian Centre for Nanomedicine and Biotechnology, BIONAND, Malaga, Spain.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.