

Published in final edited form as:

Oncogene. 2014 September 11; 33(37): 4599–4612. doi:10.1038/onc.2013.417.

p21 suppresses inflammation and tumorigenesis on pRB-deficient stratified epithelia

Cristina Saiz-Ladera^{#1}, María Fernanda Lara^{#1}, Marina Garín³, Sergio Ruiz⁴, Mirentxu Santos¹, Corina Lorz¹, Ramón García-Escudero¹, Mónica Martínez-Fernández¹, Ana Bravo⁵, Oscar Fernández-Capetillo⁴, Carmen Segrelles^{1,6}, and Jesús M Paramio^{1,6}

¹Molecular Oncology Unit. Division of Biomedicine, CIEMAT (ed70A). Ave. Complutense 40, E-28040 Madrid, Spain

³Division of Hematopoietic Innovative Therapies (HIT). Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT)/Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBER-ER). Madrid, Spain

⁴Genomic Instability Group, Spanish National Cancer Research Center, Madrid, Spain

⁵Department of Veterinary Clinical Sciences Veterinary Faculty, University of Santiago de Compostela, E-27002 Lugo, Spain

[#] These authors contributed equally to this work.

Abstract

The retinoblastoma gene product (pRb) controls proliferation and differentiation processes in stratified epithelia. Importantly, in contrast to other tissues, *Rb* deficiency does not lead to spontaneous skin tumor formation. As the cyclin dependent kinase inhibitor p21 regulates proliferation and differentiation in the absence of pRb, we analyzed the consequences of deleting p21 in pRb-ablated stratified epithelia (hereafter pRb^{Epi};p21^{-/-}). These mice display an enhancement of the phenotypic abnormalities observed in pRb^{Epi} animals, indicating that p21 partially compensates pRb absence. Remarkably, pRb^{Epi};p21^{-/-} mice show an acute skin inflammatory phenotype and develop spontaneous epithelial tumors, particularly affecting tongue and oral tissues. Biochemical analyses and transcriptome studies reveal changes affecting multiple pathways, including DNA damage and p53-dependent signaling responses. Comparative metagenomic analyses, together with the histopathological profiles, indicate that these mice constitute a faithful model for human head and neck squamous cell carcinomas. Collectively, our findings demonstrate that p21, in conjunction with pRb, plays a central role in regulating multiple epithelial processes and orchestrating specific tumor suppressor functions.

Introduction

The pocket protein family (pRb, p107 and p130) plays unique and overlapping roles in differentiation and cell cycle control (1). Of them, the *Rb* gene is the predominant family member mutated in human tumors. This reflects the essential role of pRb in the control of

⁶To whom correspondence should be addressed. jesusm.paramio@ciemat.es and carmen.segrelles@ciemat.es.