
AMLAN


Topic: 7

AMLAN: towards vision of Anaemia Mukta Odisha

Goal: Accelerated Reduction in the prevalence of anaemia in the State across all age groups.

Objectives:

- To reduce anaemia by 10% among each target beneficiary every year taking NFHS-5 as the base.
- To provide a package of interventions to reduce anaemia by addressing nutritional and non-nutritional causes.
- To increase the awareness on affordable micro-nutrient rich food.



Target Beneficiaries



Children
6 – 59
months



Children
5 – 9
years (In
School)



Adolesce
nts 10-
19 years



Pregnant
Women



Lactatin
g
Mothers



Women
of
Reprodu
ctive
Age
Group
20-24
years

Test, Treat ,Talk & Track

- In order to implement the programme effectively and to ensure maximum coverage of the target beneficiaries, 3 major platforms have been identified for ensuring the service deliveries categorized under Test, Treat and Talk. They are:
 - Schools
 - VHSND
 - T3 camps
- The testing of the beneficiaries categorised under different groups will be done by CHO / MPHW(F) / MPHW(M) and so also the treatment.
- The talk and track activities will be carried out by Teacher, MPHW(F), MPHW(M), CHO, AWW and ASHA



T3 at VHSND

- Service provider – MPHW(F)/MPHW(M)/AWW/ASHA
- Target Group
 - Children aged 6-59 months
 - Out of school Adolescents girls 10-19 years
 - Pregnant women
 - Lactating mother
 - Women in Reproductive Age (20-24 years)
- Day designated for T3(Test, Treat & Talk) – VHSND
- Test & Treat - MPHW(F)/MPHW(M)
- Counselling , IFA Tablet and facilitate filling the follow-up card – MPHW(F) and AWW
- Record keeping – MPHW(F) and AWW
- Facilitate, monitoring and home visit - ASHA

T3 in Schools

- Service provider – MPHW(F)/MPHW(M)/CHO/Teacher/ASHA
- Target Group – Students of class 1 to 12
- Day designated for T3(Test, Treat & Talk) – Monday
- Test & Treat by - MPHW(F)/MPHW(M)/CHO
- Counselling , IFA Tablet and facilitate filling the follow-up card – Teacher
- Record keeping – Teacher and MPHW(F)
- Facilitate, monitoring and home visit - ASHA

Key Activities Under AMLAN

- The beneficiary category of Children 6 to 59 months. Out of school Adolescent girls 10-19 years, Women in Reproductive Age (WRA) 20-24 year, Pregnant Women and Lactating Mother will be tested for haemoglobin estimation at their respective VHSND/UHSND.
- All students studying in class 1st to 12th in Govt and Govt. aided schools will be tested for haemoglobin estimation at their respective schools.
- Those beneficiaries who are left out or in out reach areas will be tested in T3 camps, which will be organized yearly twice at every CHC.
- After testing for the Hb. level, it should be recorded in the prescribed register by MPHW(F)/CHO/MPHW(M)/ School nodal teacher/AWW for AMLAN
- Based on the test results, necessary therapeutic dose is to be prescribed/ referral is to be done to higher centre.
- Counselling & follow-up (tracking) to be carried out.

Frequency of Screening

Every year there will be **four time screening**. First time-Universal screening of all beneficiaries

Catagory	1 st Screening	Rescreening	Action	Action
6-59 Month	Screening of all beneficiaries in first Quarter	Rescreening of only identified anaemic cases after two month consumption of therapeutic doses	After 2 months if Hb<11 gm/dl but improvement in Hb. Level >1gm then continue therapeutic dose for next two months	If no improvement or decrease of Hb then refer for investigation.
5-9yrs Sishubatika to class-5	Screening of all beneficiaries by 2 nd Quarter	Rescreening of only identified anaemic cases after two month consumption of therapeutic doses	After 2 months if Hb<11.5 gm/dl but improvement in Hb. Level >1gm then continue therapeutic dose for next two months	If no improvement or decrease of Hb then refer for investigation.
10-19yrs(In schoolclass 6-12 and out of school adolescent girls	Screening of all beneficiaries by 2 nd Quarter	Rescreening of only identified anaemic cases after three month consumption of therapeutic doses	After 3 months if Hb<12 gm/dl but improvement in Hb. Level >1gm then continue therapeutic dose for next three months	If no improvement or decrease of Hb then refer for investigation.

Frequency of Screening

Catagor y	1 st Screening	Rescreening	Action	Action
WRA (20-24months)	Screening of all beneficiaries in first Quarter	Rescreening of only identified anaemic cases after three month consumption of therapeutic doses	After 3 months if Hb <12 gm/dl but improvement in Hb. Level >1gm then continue therapeutic dose for next three months	If no improvement or decrease of Hb then refer for investigation.
Pregnant Women	Screening of all beneficiaries in every visit	Rescreening after one month consumption of therapeutic doses	In next month if Hb<11.0 gm/dl but improvement in Hb. Level >1gm then continue therapeutic dose	If no improvement or decrease of Hb then refer CHC/SDH/DHH for further investigations and will be managed with IV Iron Sucrose/ FCM
Lactating Mother	Screening of all beneficiaries in every visit	Rescreening after one month consumption of therapeutic doses	In next month months if Hb<12.0 gm/dl but improvement in Hb. Level >1gm continue therapeutic dose	If no improvement or decrease of Hb then refer CHC/SDH/DHH for further investigations and will be managed with IV Iron Sucrose/ FCM

Hemoglobin levels to diagnose Anaemia

Population	No anemia (g/dl)	Anemia		
		Mild (g/dl)	Moderate (g/dl)	Severe (g/dl)
Children 6-59 months	11.0	10 - 10.9	7 - 9.9	< 7.0
Children 5-11 years	11.5	11.0 - 11.4	8.0 - 10.9	< 8.0
Children 12 -14 years	12.0	11.0 - 11.9	8.0 - 10.9	<8.0
Non pregnant women (≥ 15 years)	12.0	11.0 - 11.9	8.0 - 10.9	< 8.0
Pregnant women	11.0	10.0 - 10.9	7.0 - 9.9	< 7.0

Source: WHO, 2011

Modalities for Oral IFA consumptions for all age groups

SN	Beneficiaries	Prophylactic	Therapeutic	Time for Consumption	Monitored by
1	6 – 59 month	Biweekly (Tuesday & Friday) 1ml of IFA syrup containing 20mg of elemental iron & 100mcg of folic acid	- Children 6–12 months- 1 ml IFA syrup, once a day - Children 1–3 years - 1.5 ml IFA syrup, once a day - Children 3–5 years - 2 ml IFA syrup, once a day	One hour after Complementary food/meal	ASHA through home visit
2	Sishubatika to Class-V	Weekly once(Monday) 1 Iron and Folic Acid Pink colour tablet	One Iron and Folic Acid Pink colour tablet daily for two months	One IFA pink colour tablet one hour after Mid-day meal at school	Teacher

<i>SN</i>	<i>Beneficiaries</i>	<i>Prophylactic</i>	<i>Therapeutic</i>	<i>Time for Consumption</i>	<i>Monitored by</i>
3	Class-VI to Class- XII	Weekly once(Monday) 1 Iron and Folic Acid Blue colour tablet	Two Iron and Folic Acid Blue colour tablet daily for three months	•Two Iron and Folic Acid Blue colour tablet one hour after Mid-day meal for Class- VI to X •Two Iron and Folic Acid Blue colour tablet after prayer class for Students of Class-XI & XII	Teacher
4	Out of School adolescent Girls 10-19years	Weekly once(Monday) 1 Iron and Folic Acid Blue colour tablet	Two Iron and Folic Acid Blue colour tablet daily for three months	One Iron and Folic Acid Blue colour tablet one hour after Breakfast and one Iron and Folic Acid Blue colour tablet one hour after evening snacks	AWW

SN	Beneficiaries	Prophylactic	Therapeutic	Time for Consumption	Monitored by
5	WRA(20-24years)	Weekly once(Saturday) 1 Iron and Folic Acid Red colour tablet	Two Iron and Folic Acid Red colour tablet daily for three months	One Iron and Folic Acid Red colour tablet one hour after Breakfast and one Iron and Folic Acid Red colour tablet one hour after evening snacks	ASHA through home visit
6	Pregnant Women	One Iron and Folic Acid tablet Red colour daily for 180 days	Two Iron and Folic Acid tablet Red colour daily for 180days	•For non anaemic PW, One Iron and Folic Acid Red colour tablet one hour after Breakfast for prophylactic cases •For mild and moderate anaemic PW, one Iron and Folic Acid Red colour tablet one hour after Breakfast and One Iron and Folic Acid Red colour tablet one hour after evening snacks for therapeutic cases	ASHA & Poshan Sakhi through home visit
		2 Calcium tablet daily for 180 days		One calcium Tablet after lunch and one calcium Tablet after Dinner	ASHA & Poshan Sakhi through home visit
		From the beginning of 2 nd trimester Multivitamin tablet once daily for 90days	From the beginning of 2 nd trimester Multivitamin tablet once daily for 180days	One multivitamin Tablet after breakfast with tablet IFA	ASHA & Poshan Sakhi through home visit

<i>S N</i>	<i>Beneficiaries</i>	<i>Prophylactic</i>	<i>Therapeutic</i>	<i>Time for Consumption</i>	<i>Monitored by</i>
7	Lactating Mother	One Iron and Folic Acid tablet Red colour daily for 180 days	Two Iron and Folic Acid tablet Red colour daily for 180days	•One Iron and Folic Acid Red colour tablet one hour after Breakfast for prophylactic cases •One Iron and Folic Acid Red colour tablet one hour after Breakfast and One Iron and Folic Acid Red colour tablet one hour after evening snacks for therapeutic cases	ASHA & Poshan Sakhi through home visit
		2 Calcium tablet daily for 180days		One calcium Tablet after lunch and one calcium Tablet after Dinner	ASHA & Poshan Sakhi through home visit

Way Forward

- Ensure 100% Hb testing of all beneficiaries in the first quarter and timely re-testing of anaemic cases after 3 months of therapeutic IFA.
- Strengthen AMLAN micro-planning and stakeholder coordination for universal screening, treatment, and follow-up.
- Conduct regular reviews and supportive supervision to address implementation gaps and improve programme performance.
- Build capacity of frontline workers on anaemia detection, therapeutic IFA administration, follow-up, and reporting.
- Ensure timely reporting, regular updating of high-priority beneficiary lists, and proper maintenance of AMLAN registers.
- Improve therapeutic IFA compliance through effective counselling, MCP card tracking, and use of IEC materials like AMLAN flipbook.