

Contents

1	Rings and Ideals	2
2	Modules	4
3	Rings and Modules of Fractions	7
4	Primary Decomposition	9
5	Integral Dependence and Valuations	11
6	Chain Conditions	13
7	Noetherian Rings	14
8	Artinian Rings	15
9	Discrete Valuation Rings and Dedekind Domain	16
10	Completion	18
11	Dimension Theory	21
EXERCISES		23
1	Rings and Ideals	23
2	Modules	24
3	Rings and Modules of Fractions	26
4	Primary Decomposition	28
5	Integral Dependence and Valuations	29
6	Chain Conditions	32
7	Noetherian Rings	32
8	Artinian Rings	34
9	Discrete Valuation Rings and Dedekind Domain	35
10	Completion	36
11	Dimension Theory	37
SUMMARY		39

1 Rings and Ideals

IDEALS. PRIME AND MAXIMAL IDEALS

Proposition 1.1. *There is a one-to-one order-preserving correspondence between the ideals $\mathfrak{b}$ of A which contain $\mathfrak{a}$, and the ideals $\bar{\mathfrak{b}}$ of $A/\mathfrak{a}$, given by $\mathfrak{b} = \phi^{-1}(\bar{\mathfrak{b}})$.*

Corollary 1.5. *Every non-unit of A is contained in a maximal ideal.*

Corollary 1.6. (i) *Let A be a ring and $\mathfrak{m} \neq (1)$ an ideal of A such that every $x \in A - \mathfrak{m}$ is a unit in A , Then A is a local ring and $\mathfrak{m}$ its maximal ideal.*

(ii) *Let A be a ring and $\mathfrak{m}$ a maximal ideal of A , such that every element of $1 + \mathfrak{m}$ is a unit in A , then A is a local ring.*

NILRADICAL AND JACOBSON RADICAL

We denote nilradical to be $\mathfrak{N}$ and Jacobson radical to be $\mathfrak{J}$.

Proposition 1.8. *The nilradical of A is the intersection of all the prime ideals of A .*

Sketch proof. Assume that $f \notin \mathfrak{N}$. Let Σ be a set of ideals $\mathfrak{a}$ with property $f^n \notin \mathfrak{a}$. Then Σ has maximal element which is prime. $\square$

Proposition 1.9. $x \in \mathfrak{J} \Leftrightarrow 1 - xy$ is a unit in A for all $y \in A$.

OPERATIONS ON IDEALS

Two ideals $\mathfrak{a}, \mathfrak{b}$ are said to be **coprime** if $\mathfrak{a} + \mathfrak{b} = 1$. Thus we have $\mathfrak{a} \cap \mathfrak{b} = \mathfrak{a}\mathfrak{b}$.

Proposition 1.10 (Chinese Remainder Theorem). *Define ϕ to be*

$$\phi : A \rightarrow \prod_{i=1}^n (A/\mathfrak{a}_i), \quad \phi(x) = (x + \mathfrak{a}_1, \dots, x + \mathfrak{a}_n).$$

(i) *If $\mathfrak{a}_i$ and $\mathfrak{a}_j$ are coprime whenever $i \neq j$, then $\mathfrak{a}_1 \cdots \mathfrak{a}_n = \bigcap_{i=1}^n \mathfrak{a}_i$.*

(ii) *ϕ is surjective $\Leftrightarrow \mathfrak{a}_i$ and $\mathfrak{a}_j$ are coprime whenever $i \neq j$.*

(iii) *ϕ is injective $\Leftrightarrow \bigcap_{i=1}^n \mathfrak{a}_i = (0)$.*

Proposition 1.11. *Denote $\mathfrak{a}$ to be ideal and $\mathfrak{p}$ to be prime ideal.*

(i) *If $\mathfrak{a} \subseteq \bigcup_{i=1}^n \mathfrak{p}_i$, then $\mathfrak{a} \subseteq \mathfrak{p}_i$ for some i .*

(ii) *If $\mathfrak{p} \supseteq \bigcap_{i=1}^n \mathfrak{a}_i$, then $\mathfrak{p} \supseteq \mathfrak{a}_i$ for some i . If $\mathfrak{p} = \bigcap_{i=1}^n \mathfrak{a}_i$, then $\mathfrak{p} = \mathfrak{a}_i$ for some i .*

The **ideal quotient** of $\mathfrak{a}$ and $\mathfrak{b}$ is

$$(\mathfrak{a} : \mathfrak{b}) = \{x \in A : x\mathfrak{b} \subseteq \mathfrak{a}\}.$$

The annihilator of $\mathfrak{b}$ is $\text{Ann}(\mathfrak{b}) = (0 : \mathfrak{b})$.

The **radical** of $\mathfrak{a}$ is

$$\sqrt{\mathfrak{a}} = \{x \in A : x^n \in \mathfrak{a} \text{ for some } n > 0\}.$$

EXTENSION AND CONTRACTION

Let $f : A \rightarrow B$ be a ring homomorphism. Let $\mathfrak{a} \in A$ and $\mathfrak{b} \in B$. Define

$$\mathfrak{a}^e = Bf(\mathfrak{a}), \quad \mathfrak{b}^c = f^{-1}(\mathfrak{b})$$

to be the **extension** of $\mathfrak{a}$ and the **contraction** of $\mathfrak{b}$, respectively. If $\mathfrak{b}$ is prime, $\mathfrak{b}^c$ is prime; if $\mathfrak{a}$ is prime, $\mathfrak{a}^e$ need not be prime.

Proposition 1.17. $\mathfrak{a} \subseteq \mathfrak{a}^{ec}$, $\mathfrak{b} \supseteq \mathfrak{b}^{ce}$.

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

Idempotent is an element e which satisfies $e^2 = e$.

A ring in which every element is an idempotent is a **Boolean ring**.

A topological space X is **reducible** if it can be written as a union $X = X_1 \cup X_2$ of two closed proper subsets X_1, X_2 of X . Equivalently, X is **irreducible** if every pair of non-empty open sets in X intersect, or if every non-empty open set is dense in X .

Let A be a ring and let $\text{Spec}(A)$ be the set of all prime ideals of A . For each subset E of A , let $V(E)$ denote the set of all prime ideals of A which contain E . By [Exercise 1.15](#), the sets $V(E)$ satisfy the axioms for closed sets in a topological space. The resulting topology on $\text{Spec}(A)$ is called **Zariski topology**, and $\text{Spec}(A)$ is called the **prime spectrum** of A .

For each $f \in A$, the complement of $V(f)$ in $\text{Spec}(A)$, denote to be X_f , is called **basic open set** of $\text{Spec}(A)$.

The subspace of $\text{Spec}(A)$ consisting of the maximal ideals of A with the induced topology, is called the **maximal spectrum** of A and is denoted by $\text{Max}(A)$.

Proposition ([Exercise 1.27](#)). *Let k be algebraically closed. Then the affine space k^n with its classical Zariski topology is homeomorphic to $\text{Max}(k[x_1, \dots, x_n])$.*

2 Modules

Proposition 2.1. (i) If $L \supseteq M \supseteq N$ are A -modules, then $(L/N)/(M/N) \cong L/M$.

(ii) If M_1, M_2 are submodules of M , then $(M_1 + M_2)/M_1 \cong M_2/(M_1 \cap M_2)$.

An A -module is **faithful** if $\text{Ann}(M) = 0$.

FINITELY GENERATED MODULES

Proposition 2.3. M is a finitely generated A -module $\Leftrightarrow M$ is isomorphic to a quotient of A^n for some integer $n > 0$. More precisely, n is the number of generators of M and may not be unique.

Proposition 2.4 (**Cayley–Hamilton theorem**). Let M be a finitely generated A -module, let $\mathfrak{a}$ be an ideal of A , and let ϕ be an A -module endomorphism of M such that $\phi(M) \subseteq \mathfrak{a}M$. Then ϕ satisfies an equation of the form

$$\phi^n + a_1\phi^{n-1} + \cdots + a_n = 0$$

where the a_i are in $\mathfrak{a}$.

Proposition 2.6 (**Nakayama's lemma**). Let M be a finitely generated A -module and $\mathfrak{a}$ an ideal of A contained in the Jacobson radical of A . Then $\mathfrak{a}M = M$ implies $M = 0$.

EXACT SEQUENCES

Proposition 2.10 (Snake lemma). Let

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \xrightarrow{u} & M & \xrightarrow{v} & M'' \longrightarrow 0 \\ & & \downarrow f' & & \downarrow f & & \downarrow f'' \\ 0 & \longrightarrow & N' & \xrightarrow{u'} & N & \xrightarrow{v'} & N'' \longrightarrow 0 \end{array}$$

be a commutative diagram of A -modules and homomorphisms, with the rows exact. Then there exists an exact sequence

$$0 \longrightarrow \ker(f') \xrightarrow{\bar{u}} \ker(f) \xrightarrow{\bar{v}} \ker(f'') \xrightarrow{\delta} \text{coker}(f') \xrightarrow{\bar{u}'} \text{coker}(f) \xrightarrow{\bar{v}'} \text{coker}(f'') \longrightarrow 0.$$

Let C be a class of A -modules and let λ be a function on C with values in $\mathbb{Z}$. The function λ is **additive** if for each short exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

in which all terms belong to C , we have $\lambda(M') - \lambda(M) + \lambda(M'') = 0$.

TENSOR PRODUCT

Proposition 2.12 (Universal property). Let M, N be A -modules. The tensor product of M and N over A is an A -module $M \otimes_A N$ together with an A -bilinear map

$$f : M \times N \rightarrow M \otimes_A N, \quad (m, n) \mapsto m \otimes n,$$

such that, for every A -module P and every A -bilinear map $\tau : M \times N \rightarrow P$, there exists a unique A -linear map $\tilde{\tau} : M \otimes_A N \rightarrow P$ such that $\tau = \tilde{\tau} \circ f$. Equivalently, the following diagram commutes:

$$\begin{array}{ccc} M \times N & \xrightarrow{f} & M \otimes_A N \\ & \searrow \tau & \downarrow \tilde{\tau} \\ & & P \end{array}$$

Proposition 2.15. Let A, B be rings, let M be an A -module, P a B -module and N an (A, B) -bimodule. then

$$(M \otimes_A N) \otimes_B P \cong M \otimes_A (N \otimes_B P).$$

Let M be an A -module and $f : A \rightarrow B$ be a homomorphism of rings. Define $M_B = B \otimes_A M$.

Proposition 2.19. For an A -module N , the following are equivalent:

- (i) The functor $T_N : M \mapsto M \otimes_A N$ on the category of A -modules is exact.
- (ii) If $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is any exact sequence of A -modules, then $0 \rightarrow M' \otimes N \rightarrow M \otimes N \rightarrow M'' \otimes N \rightarrow 0$ is exact.
- (iii) If $f : M' \rightarrow M$ is injective, then $f \otimes 1 : M' \otimes N \rightarrow M \otimes N$ is injective.
- (iv) If $f : M' \rightarrow M$ is injective and M, M' are finitely generated, then $f \otimes 1 : M' \otimes N \rightarrow M \otimes N$ is injective.

An A -module N satisfying these properties is said to be a **flat** A -module.

ALGEBRA

Let $f : A \rightarrow B$ be a ring homomorphism. If $a \in A$ and $b \in B$, define a product

$$ab = f(a)b.$$

Thus, B has an A -module structure. The ring B is said to be an **A -algebra**.

A ring homomorphism $f : A \rightarrow B$ is **finite**, if B is finitely generated A -module. We can also say B is a **finite** A -algebra, or B is **finite** over A .

The ring homomorphism f is **of finite type**, if B is a finitely generated A -algebra.

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A partially ordered set I is said to be a **directed** set if for each pair $i, j \in I$ there exists $k \in I$ such that $i \leq k$ and $j \leq k$.

Let A be a ring, let I be a directed set and let $(M_i)_{i \in I}$ be a family of A -modules indexed by I . For each pair $i, j \in I$ such that $i \leq j$, let $\mu_{ij} : M_i \rightarrow M_j$ be an A -homomorphism, and suppose:

- (1) μ_{ii} is the identity mapping of M_i , for all $i \in I$;
- (2) $\mu_{ik} = \mu_{jk} \circ \mu_{ij}$ whenever $i \leq j \leq k$.

Then the modules M_i and homomorphisms μ_{ij} are said to form a **direct system** $\mathbf{M} = (M_i, \mu_{ij})$ over the directed set I .

Let D be the submodule of $\bigoplus M_i$ generated by all elements of the form $x_i - \mu_{ij}(x_i)$ whenever $i \leq j$ and $x_i \in M_i$. The **direct limit** of the direct system $\mathbf{M}$ is defined to be

$$\varinjlim M_i = \bigoplus_{i \in I} M_i / D.$$

Let $\mu : \bigoplus M_i \rightarrow \varinjlim M_i$ be the projection of quotient module. Define μ_i to be the restriction of μ to M_i .

A ring A (not necessarily commutative) is **von Neumann regular** if for every $a \in A$, there is an $x \in A$ with $a = axa$. A commutative ring A is **absolutely flat** if every A -module is flat. By [Exercise 2.27](#), absolute flat and von Neumann regular are identical properties.

3 Rings and Modules of Fractions

MULTIPLICATIVELY CLOSED SETS

Proposition 3.3. *The operation S^{-1} is exact.*

Proposition 3.5. *Let M be an A -module. Then there exists a unique $S^{-1}A$ module isomorphism $f : S^{-1}A \otimes_A M \rightarrow S^{-1}M$.*

LOCAL PROPERTIES

Let P be a property of a ring A (or of an A -module M). Then the property P is said to be a **local property**, if A (or M) has $P \Leftrightarrow A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has P for each prime ideal $\mathfrak{p}$ of A .

Proposition 3.8. *Let M, N be A -modules. Then the following are equivalent:*

- (i) $M = 0$;
- (ii) $M_{\mathfrak{p}} = 0$ for all prime ideals $\mathfrak{p}$ of A ;
- (iii) $M_{\mathfrak{m}} = 0$ for all maximal ideals $\mathfrak{m}$ of A .

Proposition 3.9. *Let $\phi : M \rightarrow N$ be an A -module homomorphism. Then the following are equivalent:*

- (i) ϕ is injective;
- (ii) $\phi_{\mathfrak{p}} : M_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}$ is injective for each prime ideal $\mathfrak{p}$;
- (iii) $\phi_{\mathfrak{m}} : M_{\mathfrak{m}} \rightarrow N_{\mathfrak{m}}$ is injective for each prime ideal $\mathfrak{m}$.

Similarly with “injective” replaced by “surjective” throughout.

Proposition 3.10. *Let M be A -modules. Then the following are equivalent:*

- (i) M is a flat A -module;
- (ii) $M_{\mathfrak{p}}$ is a flat $A_{\mathfrak{p}}$ -module for each prime ideal $\mathfrak{p}$;
- (iii) $M_{\mathfrak{m}}$ is a flat $A_{\mathfrak{m}}$ -module for each maximal ideal $\mathfrak{m}$.

EXTENSIONS AND CONTRACTIONS OF FRACTIONS

If $\mathfrak{a}$ is an ideal in A , its extension $\mathfrak{a}^e$ in $S^{-1}A$ is $S^{-1}\mathfrak{a}$. Denote the contraction of $S^{-1}\mathfrak{a}$ in A to be $S(\mathfrak{a})$.

Proposition 3.11. (i) *Every ideal in $S^{-1}A$ is an extended ideal.*

(ii) *If $\mathfrak{a}$ is an ideal in A , then $S(\mathfrak{a}) = \mathfrak{a}^{ec} = \bigcup_{s \in S} (\mathfrak{a} : s)$.*

(iii) *$\mathfrak{a}$ is a contracted ideal $\Leftrightarrow$ no element of S is a zero-divisor in $A/\mathfrak{a}$.*

(iv) *Them prime ideal of $S^{-1}A$ are in one-to-one correspondence ($\mathfrak{p} \leftrightarrow S^{-1}\mathfrak{p}$) with the prime ideal of A which don't meet S .*

(v) *The operation S^{-1} commutes with formation of finite sums, products, intersections and radicals.*

Corollary 3.15. *If N, P are submodules of an A -module M and if P is finitely generated, then $S^{-1}(N : P) = (S^{-1}N : S^{-1}P)$.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A multiplicatively closed subset S of a ring A is said to be **saturated** if

$$xy \in S \Leftrightarrow x \in S \text{ and } y \in S.$$

Let M be an A -module. An element $x \in M$ is a **torsion** element of M if there exist a non-zero-divisor $a \in A$ such that $ax = 0$. All torsion elements of M form a submodule of M , which is called the **torsion submodule** of M and denote by $T(M)$. If $T(M) = 0$, the module M is said to be **torsion free**.

An A -module M is called **faithfully flat** if the tensor product functor $- \otimes_A M$ is exact and faithful. See [Exercise 3.16](#).

Let M be an A -module. The **support** of M is defined to be the set $\text{Supp}(M)$ of prime ideals $\mathfrak{p}$ such that $M_{\mathfrak{p}} \neq 0$.

$\text{Spec}(A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \otimes_A B)$ is called the **fiber** of $f^* : \text{Spec}(B) \rightarrow \text{Spec}(A)$ over $\mathfrak{p}$.

4 Primary Decomposition

PRIMARY IDEALS

An ideal $\mathfrak{q} \in A$ is **primary** if $\mathfrak{q} \neq A$, and if $x \notin \mathfrak{q} \Rightarrow y^n \in \mathfrak{q}$ for some $n > 0$. This also means $x \notin \sqrt{\mathfrak{q}} \Rightarrow y \in \mathfrak{q}$.

Proposition 4.2. *If $\sqrt{\mathfrak{a}}$ is maximal, then $\mathfrak{a}$ is primary.*

Sketch proof. Let $xy \in \mathfrak{a}$ but $x \notin \sqrt{\mathfrak{a}}$. We have $\sqrt{\mathfrak{a}} + (x) = A$, so $m + ax = 1$ with $m^n \in \mathfrak{a}$. Therefore $m^n + bx = 1$ and multiply y to both sides. $\square$

Proposition 4.4. *Let $\mathfrak{q}$ be a $\mathfrak{p}$ -primary ideal, $x \in A$ an element. Then*

- (i) *if $x \in \mathfrak{q}$ then $(\mathfrak{q} : x) = (1)$;*
- (ii) *if $x \notin \mathfrak{q}$ then $(\mathfrak{q} : x)$ is $\mathfrak{p}$ -primary;*
- (iii) *if $x \notin \mathfrak{p}$ then $(\mathfrak{q} : x) = \mathfrak{q}$;*

Proposition 4.8. *Let S be a multiplicatively closed subset of A , and let $\mathfrak{q}$ be a $\mathfrak{p}$ -primary ideal.*

- (i) *If $S \cap \mathfrak{p} \neq \emptyset$, then $S^{-1}\mathfrak{q} = S^{-1}A$.*
- (ii) *If $S \cap \mathfrak{p} = \emptyset$, then $S^{-1}\mathfrak{q}$ is $S^{-1}\mathfrak{p}$ -primary and its contraction in A is $\mathfrak{q}$.*

A **primary decomposition** of an ideal $\mathfrak{a}$ in A is an expression $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$.

If $\sqrt{\mathfrak{q}_i}$ are all distinct and $\mathfrak{q}_i \not\supseteq \bigcap_{j \neq i} \mathfrak{q}_j$, the primary decomposition is said to be **irredundant**.

UNIQUENESS

Proposition 4.5 (1st uniqueness theorem). *Let $\mathfrak{a}$ be a decomposable ideal and let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ be a minimal primary decomposition of $\mathfrak{a}$. Let $\mathfrak{p}_i = \sqrt{\mathfrak{q}_i}$. Then $\mathfrak{p}_i$ are precisely the prime ideals which occur in the set of ideals $\sqrt{(\mathfrak{a} : x)}$, and hence are independent of the particular decomposition of $\mathfrak{a}$.*

The prime ideals $\mathfrak{p}_i$ are said to belong to $\mathfrak{a}$, or to be **associated** with $\mathfrak{a}$. The minimal elements of the set $\{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$ are called the minimal or **isolated** prime ideals belonging to $\mathfrak{a}$. The others are called **embedded** prime ideals.

Proposition 4.9. *Let S be a multiplicatively closed subset of A and let $\mathfrak{a}$ be a decomposable ideal. Let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ be a minimal primary decomposition of $\mathfrak{a}$. Let $\mathfrak{p}_i = \sqrt{\mathfrak{q}_i}$ and suppose the $\mathfrak{q}_i$ numbered so that S meets $\mathfrak{p}_{m+1}, \dots, \mathfrak{p}_n$ but not $\mathfrak{p}_1, \dots, \mathfrak{p}_m$. Then we have minimal primary decompositions*

$$S^{-1}\mathfrak{a} = \bigcap_{i=1}^m S^{-1}\mathfrak{q}_i, \quad S(\mathfrak{a}) = \bigcap_{i=1}^m \mathfrak{q}_i.$$

A set Σ of prime ideals belonging to $\mathfrak{a}$ is said to be **isolated** if it satisfies the following condition: if $\mathfrak{p}'$ is a prime ideal belonging to $\mathfrak{a}$ and $\mathfrak{p}' \subseteq \mathfrak{p}$ for some $\mathfrak{p} \in \Sigma$, then $\mathfrak{p}' \in \Sigma$. Thus, every isolated set contains an isolated prime ideal.

Proposition 4.10 (2nd uniqueness theorem). *Let $\mathfrak{a}$ be a decomposable ideal, let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ be a minimal primary decomposition of $\mathfrak{a}$, and let $\{\mathfrak{p}_{i_1}, \dots, \mathfrak{p}_{i_m}\}$ be an isolated set of prime ideals of $\mathfrak{a}$. Then the ideal $\mathfrak{q}_{i_1} \cap \dots \cap \mathfrak{q}_{i_m}$ is independent of the decomposition.*

Corollary 4.11. *The isolated primary components are uniquely determined by $\mathfrak{a}$.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

Let A be a ring and $\mathfrak{p}$ a prime ideal of A . The n th **symbolic power** of $\mathfrak{p}$ is defined to be the ideal $\mathfrak{p}^{(n)} = S_{\mathfrak{p}}(\mathfrak{p}^n)$.

Let M be an A -module. A submodule $P \subseteq M$ is called a **prime submodule** if for any $a \in A$ and $x \in M$, $ax \in P$ implies $x \in P$ or $aM \subseteq P$.

A submodule $Q \subseteq M$ is called a **primary submodule** if $ax \in Q$ implies $x \in Q$ or $a^n M \subseteq Q$ for some $n > 0$.

Let M be an A -module, N a submodule of M . The **ideal radical** $\text{rad}_M(N)$ of N in M is defined to be $\text{rad}_M(N) = \sqrt{(N : M)}$. The **prime radical** $\text{Rad}_M(N)$ of N in M is defined to be the intersection of all prime submodules containing N .

Proposition. *The ideal radical and the prime radical satisfy*

$$(\text{Rad}_M(N) : M) = \text{rad}_M(N).$$

Sketch proof. $(\text{Rad}_M(N) : M) = \bigcap_{P \supseteq N} (P : M)$, so we need to prove $\bigcap_{P \supseteq N} (P : M) = \sqrt{(N : M)}$.

If $N \subseteq P$, $(N : M) \subseteq (P : M)$. Since $(P : M)$ prime, $\sqrt{(N : M)} \subseteq (P : M)$ and therefore $\sqrt{(N : M)} \subseteq \bigcap_{P \supseteq N} (P : M)$.

Let $a \in (P : M)$ for all $P \supseteq N$. For any $\mathfrak{p} \supseteq (N : M)$, let

$$K_{\mathfrak{p}}(N) = \{m \in M : sm \in \mathfrak{p}M + N \text{ for some } s \in A - \mathfrak{p}\},$$

we have $N \subseteq K_{\mathfrak{p}}(N)$. If $(K_{\mathfrak{p}}(N) : M) = \mathfrak{p}$, then $a \in \mathfrak{p}$ and therefore $a \in \sqrt{(N : M)}$.

Let $x \in (K_{\mathfrak{p}}(N) : M)$, then $xsm \in \mathfrak{p}M + N$ for all $m \in M$ and some $s \in A - \mathfrak{p}$. If $x \notin \mathfrak{p}$, $xs \in A - \mathfrak{p}$, and therefore $K_{\mathfrak{p}}(N) = M$, contradiction. Hence $(K_{\mathfrak{p}}(N) : M) \subseteq \mathfrak{p}$.

For any $x \in \mathfrak{p}$, $xM \subseteq \mathfrak{p}M \subseteq \mathfrak{p}M + N$. Therefore, $x \in (K_{\mathfrak{p}}(N) : M)$ and so $\mathfrak{p} \subseteq (K_{\mathfrak{p}}(N) : M)$. $\square$

5 Integral Dependence and Valuations

INTEGRAL DEPENDENCE

Proposition 5.1. *The following are equivalent:*

- (i) $x \in B$ is integral over A ;
- (ii) $A[x]$ is a finitely generated A -module;
- (iii) $A[x]$ is contained in a subring C of B such that C is a finitely generated A -module;
- (iv) There exist a faithful $A[x]$ -module M which is finitely generated as an A -module.

The set $\bar{A}$ of elements of B which are integral over A is called the **integral closure** of A in B . If $\bar{A} = A$, then A is said to be **integrally closed in B** . If $\bar{A} = B$, the ring B is said to be **integral over A** .

THE GOING-UP THEOREM

Proposition 5.7. *Let $A \subseteq B$ be integral domains, B integral over A . Then B is a field iff A is a field.*

Corollary 5.9. *Let $A \subseteq B$ be rings, B integral over A . Let $\mathfrak{q}_1, \mathfrak{q}_2$ be prime ideals of B such that $\mathfrak{q}_1 \subseteq \mathfrak{q}_2$ and $\mathfrak{q}_1^e = \mathfrak{q}_2^e = \mathfrak{p}$. Then $\mathfrak{q}_1 = \mathfrak{q}_2$.*

Theorem 5.10 (Lying over property). *Let $A \subseteq B$ be rings, B integral over A , and let $\mathfrak{p}$ be a prime ideal of A . Then there exists a prime ideal $\mathfrak{q}$ of B such that $\mathfrak{q} \cap A = \mathfrak{p}$.*

Theorem 5.11 (Going-up theorem). *Let $A \subseteq B$ be rings, B integral over A . Let $\mathfrak{p}_1 \subseteq \cdots \subseteq \mathfrak{p}_n$ be a chain of prime ideals of A and $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_m$ ($m < n$) a chain of prime ideals of B such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq m$. Then the chain $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_m$ can be extended to a chain $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_n$ such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq n$.*

THE GOING-DOWN THEOREM

Proposition 5.12. *Let $A \subseteq B$ be rings, C the integral closure of A in B . Let S be a multiplicatively closed subset of A . Then $S^{-1}C$ is the integral closure of $S^{-1}A$ in $S^{-1}B$.*

An integral domain is said to be **integrally closed** (without qualification) if it is integrally closed in its field of fractions.

Proposition 5.13. *Let A be an integral domain. Then the following are equivalent:*

- (i) A is integrally closed;
- (ii) $A_{\mathfrak{p}}$ is integrally closed, for each prime ideal $\mathfrak{p}$;
- (iii) $A_{\mathfrak{m}}$ is integrally closed, for each maximal ideal $\mathfrak{m}$.

Theorem 5.16 (Going-down theorem). *Let $A \subseteq B$ be integral domains, A integrally closed, B integral over A . Let $\mathfrak{p}_1 \supseteq \cdots \supseteq \mathfrak{p}_n$ be a chain of prime ideals of A and $\mathfrak{q}_1 \supseteq \cdots \supseteq \mathfrak{q}_m$ ($m < n$) a chain of prime ideals of B such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq m$. Then the chain $\mathfrak{q}_1 \supseteq \cdots \supseteq \mathfrak{q}_m$ can be extended to a chain $\mathfrak{q}_1 \supseteq \cdots \supseteq \mathfrak{q}_n$ such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq n$.*

VALUATION RINGS

Let B be an integral domain. B is a **valuation ring** of $\text{Frac}(B)$ if, for each $x \neq 0$, either $x \in B$ or $x^{-1} \in B$ (or both).

Proposition 5.18. *Let B be a valuation ring, $K = \text{Frac}(B)$ its field of fractions. Then*

- (i) B is a local ring;
- (ii) If B' is a ring such that $B \subseteq B' \subseteq K$, then B' is a valuation ring of K ;
- (iii) B is integrally closed.

Theorem 5.21. *Let K be a field, Ω an algebraically closed field. Let Σ be the set of all pairs (A, f) , where A is a subring of K and $f : A \rightarrow \Omega$ a homomorphism. Define a partial order of Σ to be:*

$$(A, f) \leq (B, g) \Leftrightarrow A \subseteq B \text{ and } g|_A = f.$$

Then the maximal element of Σ is a valuation ring of K .

Corollary 5.22. *Let A be a subring of a field K . Then the integral closure $\bar{A}$ of A in K is the intersection of all the valuation rings of K which contain A .*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A ring homomorphism $f : A \rightarrow B$ is said to have the **going-up property** if for $\mathfrak{p}_1 \subseteq \mathfrak{p}_2$ and $\mathfrak{p}_1 = f^{-1}(\mathfrak{q}_1)$, there exist a $\mathfrak{q}_2$ such that $\mathfrak{q}_1 \subseteq \mathfrak{q}_2$ and $f^{-1}(\mathfrak{q}_2) = \mathfrak{p}_2$. Reversing the inclusions of prime ideals we have the **going-down property**.

Proposition (Noether normalization lemma). *Let k be a field and let $A \neq 0$ be a finitely generated k -algebra. Then there exist elements $y_1, \dots, y_r \in A$ which are algebraically independent over k and such that A is integral, and therefore finite over $k[y_1, \dots, y_r]$.*

A ring A satisfied the following equivalent property is a **Jacobson ring**.

- (i) Every prime ideal in A is an intersection of maximal ideals.
- (ii) In every homomorphic image of A the nilradical is equal to the Jacobson radical.
- (iii) Every prime ideal in A which is not maximal is equal to the intersection of the prime ideals which contain it strictly.

A subset of a topological space X is **locally closed** if it is the intersection of an open set and a closed set.

A subset X_0 satisfying the following equivalent conditions is said to be **very dense** in X :

- (1) Every non-empty locally closed subset of X meets X_0 .
- (2) For every closed set E in X we have $\overline{E \cap X_0} = E$.
- (3) The mapping $U \mapsto U \cap X_0$ of the collection of open sets of X on to the collection of open sets of X_0 is bijective.

Let A be a valuation ring of a field K . The group U of units of A is a subgroup of the multiplicative group K^* of K . Let $\Gamma = K^*/U$. If $\xi, \iota \in \Gamma$ are represented by $x, y \in K$, define $\xi \geq \iota$ to mean $xy^{-1} \in A$. Then Γ is a totally ordered abelian group. It is called the **value group** of A .

Conversely, let Γ be a totally ordered abelian group. A **valuation** of K with values in Γ is a mapping $v : K^* \rightarrow \Gamma$ such that

- (1) $v(xy) = v(x) + v(y)$,
- (2) $v(x + y) \geq \min(v(x), v(y))$,

for all $x, y \in K^*$. Then the set of elements $x \in K^*$ such that $v(x) \geq 0$ is a valuation ring of K . This ring is called the **valuation ring** of v , and the subgroup $v(K^*)$ of Γ is the **value group** of v . Thus the concepts of valuation ring and valuation are equivalent.

6 Chain Conditions

Let Σ be a partially ordered set. Σ said to be satisfying the **ascending chain condition** (a.c.c) if, for every increasing sequence $x_1 \leq x_2 \leq \dots$ in Σ there exists n such that $x_n = x_{n+1} = \dots$.

Similarly, Σ said to be satisfying the **descending chain condition** (d.c.c) if, for every decreasing sequence $x_1 \geq x_2 \geq \dots$ in Σ there exists n such that $x_n = x_{n+1} = \dots$.

A module satisfying the ascending chain condition is said to be **Noetherian**. A module satisfying the descending chain condition is said to be **Artinian**.

Proposition 6.2. *M is a Noetherian A -module $\Leftrightarrow$ every submodule of M is finitely generated.*

Proposition 6.3. *Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence of A -modules. Then*

(i) *M is Noetherian $\Leftrightarrow M'$ and M'' are Noetherian.*

(ii) *M is Artinian $\Leftrightarrow M'$ and M'' are Artinian.*

Proposition 6.5. *Let A be a Noetherian (resp. Artinian) ring, M a finitely generated A -module. Then M is Noetherian (resp. Artinian).*

A **chain** of submodules of a module M is a sequence (M_i) of submodules of M such that

$$M = M_0 \supsetneq M_1 \supsetneq \dots \supsetneq M_n = 0.$$

The **length** of the chain is n . A **composition series** of M is a maximal chain, that is one which no extra submodules can be inserted, or equivalently M_{i-1}/M_i is simple.

Proposition 6.7. *Suppose that M has a composition series of length n . Then every composition series of M has length n , and every chain can be extended to a composition series.*

The length of a composition series of M is called the **length** of M , denoted as $l(M)$.

Proposition 6.10. *For a vector space,*

$$\text{finite dimension} \iff \text{finite length} \iff \text{a.c.c.} \iff \text{d.c.c.}$$

If the conditions are satisfied, length equals dimension.

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A topological space X is said to be **Noetherian** if the open subsets of X satisfy the ascending chain condition. Or equivalently, the closed subsets satisfy descending chain condition.

7 Noetherian Rings

Proposition 7.1. *Any homomorphic image of Noetherian ring is Noetherian.*

Proposition 7.2. *Let A be a subring of B ; suppose that A is Noetherian and that B is finitely generated as an A -module. Then B is Noetherian.*

Proposition 7.3. *If A is Noetherian and S is any multiplicatively closed subset of A , then $S^{-1}A$ is Noetherian.*

Theorem 7.5 (Hilbert's basis theorem). *If A is Noetherian, then the polynomial ring $A[x]$ is Noetherian.*

Proposition 7.8. *Let $A \subseteq B \subseteq C$ be rings. Suppose that A is Noetherian, that C is finitely generated as an A -algebra and that C is either (i) finitely generated as a B -module or (ii) integral over B . Then B is finitely generated as an A -algebra.*

Proposition 7.9 (Zariski's lemma). *Let k be a field, E a finitely generated k -algebra. If E is a field then it is a finite algebraic extension of k .*

PRIMARY DECOMPOSITION IN NOETHERIAN RINGS

Proposition 7.11. *In a Noetherian ring every ideal is a finite intersection of irreducible ideals.*

Proposition 7.12. *In a Noetherian ring every irreducible ideal is primary.*

Proposition 7.13 (Lasker Noether theorem). *In a Noetherian ring every ideal has a primary decomposition.*

Proposition 7.14. *In a Noetherian ring every ideal contains a power of its radical.*

Corollary 7.15. *In a Noetherian ring the nilradical is nilpotent.*

Corollary 7.16. *Let A be a Noetherian ring, $\mathfrak{m}$ a maximal ideal of A , $\mathfrak{q}$ any ideal of A . Then the following are equivalent:*

- (i) $\mathfrak{q}$ is $\mathfrak{m}$ -primary;
- (ii) $\sqrt{\mathfrak{q}} = \mathfrak{m}$;
- (iii) $\mathfrak{m}^n \subseteq \mathfrak{q} \subseteq \mathfrak{m}$ for some $n > 0$.

Proposition 7.17. *Let $\mathfrak{a} \neq (1)$ be an ideal in a Noetherian ring. Then the prime ideals which belong to $\mathfrak{a}$ are precisely the prime ideals which occur in the set of ideals $(\mathfrak{a} : x)$.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A subset of a topological space X is **constructible set** if it is a finite union of locally closed sets.

8 Artinian Rings

Proposition 8.1. *In an Artinian ring every prime ideal is maximal.*

Proposition 8.3. *An Artinian ring has only a finite number of maximal ideals.*

Proposition 8.4. *In an Artinian ring the nilradical is nilpotent.*

The **dimension** of A is the supremum of the length of all chains of prime ideals in A .

Proposition 8.5. *A ring A is Artinian $\Leftrightarrow A$ is Noetherian and $\dim(A) = 0$.*

Proposition 8.6. *Let A be a Noetherian local ring, $\mathfrak{m}$ its maximal ideal. Then exactly one of the following statements is true:*

- (i) $\mathfrak{m}^n \neq \mathfrak{m}^{n+1}$ for all n ;
- (ii) $\mathfrak{m}^n = 0$ for some n . In this case A is an Artinian local ring.

Proposition 8.7 (Structure theorem of Artinian rings). *An Artinaian ring A is uniquely (up to isomorphism) a finite direct product of Artinian local rings.*

Proposition 8.8. *Let A be an Artinian local ring. Then the following are equivalent:*

- (i) *every ideal in A is principal;*
- (ii) *the maximal ideal $\mathfrak{m}$ is principal;*
- (iii) $\dim_{A/\mathfrak{m}}(\mathfrak{m}/\mathfrak{m}^2) \leq 1$.

9 Discrete Valuation Rings and Dedekind Domain

Proposition 9.1. *Let A be a Noetherian domain of dimension 1. Then every non-zero ideal $\mathfrak{a}$ in A can be uniquely expressed as a product of primary ideals whose radicals are all distinct.*

DISCRETE VALUATION RINGS

Let K be a field. A **discrete valuation** on K is a mapping v of K^* onto $\mathbb{Z}$ such that

- (1) $v(xy) = v(x) + v(y)$;
- (2) $v(x + y) \geq \min(v(x), v(y))$.

The set consisting of all $x \in K^*$ such that $v(x) \geq 0$ called the **valuation ring** of v .

An integral domain A is a **discrete valuation ring** if there is a discrete valuation v of its field of fractions K such that A is the valuation ring of v .

Proposition 9.2. *Let A be a Noetherian local domain of dimension 1, $\mathfrak{m}$ its maximal ideal. Then the following are equivalent:*

- (i) A is a discrete valuation ring;
- (ii) A is integrally closed;
- (iii) $\mathfrak{m}$ is a principal ideal;
- (iv) $\dim_k(\mathfrak{m}/\mathfrak{m}^2) = 1$;
- (v) Every non-zero ideal is a power of $\mathfrak{m}$;
- (vi) There exists $x \in A$ such that every non-zero ideal is of the form (x^k) for $k \geq 0$.

DEDEKIND DOMAINS

Theorem 9.3. *Let A be a Noetherian local domain of dimension 1. Then the following are equivalent:*

- (i) A is integrally closed;
- (ii) Every primary ideal in A is a prime power;
- (iii) Every local ring $A_{\mathfrak{p}}$ with $\mathfrak{p} \neq 0$ is a discrete valuation ring.

A ring satisfying the conditions is called a **Dedekind domain**.

FRACTIONAL IDEALS

Let A be an integral domain, K its field of fractions. An A -submodule M of K is a **fractional ideal** of A if $xM \subseteq A$ for some $x \neq 0$.

An A -submodule M of K is an **invertible ideal** if there exists a submodule N of K such that $MN = A$. Then $N = (A : M)$, and M is finitely generated fractional ideal.

Proposition 9.6. *For a fractional ideal M , the following are equivalent:*

- (i) M is invertible;

- (ii) M is finitely generated and , for each prime ideal $\mathfrak{p}$, $M_{\mathfrak{p}}$ is invertible;
- (iii) M is finitely generated and , for each maximal ideal $\mathfrak{m}$, $M_{\mathfrak{m}}$ is invertible;

Theorem 9.8. *Let A be an integral domain. Then A is a Dedekind domain $\Leftrightarrow$ every non-zero fractional ideal of A is invertible.*

10 Completion

TOPOLOGIES AND COMPLETIONS

A **topological abelian group** G is both a topological space and an abelian group, with two structural mappings

$$\begin{aligned} G \times G &\rightarrow G : (x, y) \mapsto x + y \\ G &\rightarrow G : x \mapsto -x \end{aligned}$$

being continuous.

If $\{0\}$ is closed in G , then G is Hausdorff.

Since $T_a(x) = x + a$ is a homeomorphism, the topology of G is uniquely determined by the neighborhoods of 0 in G .

Proposition 10.1. *Let H be the intersection of all neighborhoods of 0 in G . Then*

- (i) H is a subgroup.
- (ii) H is the closure of $\{0\}$.
- (iii) G/H is Hausdorff.
- (iv) G is Hausdorff $\Leftrightarrow H = 0$.

Assume that $0 \in G$ has a countable fundamental system of neighborhoods. A **Cauchy sequence** (x_ν) of G is that, for any neighborhood U of 0, there exists an integer $s(U)$ such that $x_m - x_n \in U$ for all $m, n \geq s(U)$.

Let $C(G)$ be the set of all Cauchy sequences in G . Define an equivalence relation $\sim$ on $C(G)$ where $(x_\nu) \sim (y_\nu)$ means $x_n - y_n \rightarrow 0$ in G . The **completion** $\hat{G}$ of G is the set of all equivalence classes in $C(G)/\sim$.

For each $x \in G$, define a homomorphism $\phi : G \rightarrow \hat{G}$ maps to a constant sequence (x) . Then ϕ is injective iff G is Hausdorff.

Assume further that $0 \in G$ has a fundamental system of neighborhoods consisting of a sequence of subgroups

$$G = G_0 \supseteq G_1 \supseteq \cdots \supseteq G_n \supseteq \cdots$$

For any $(x_\nu) \in C(G)$ and a fixed n , the image sequence $(\xi_\nu + G_n)$ in G/G_n is ultimately constant. Let ξ_n be this ultimate constant value.

As n varies, the projection $\theta_{n+1} : G/G_{n+1} \rightarrow G/G_n$ satisfies $\theta_{n+1}(\xi_{n+1}) = \xi_n$ for all n . The groups G/G_n combined with maps θ_n form an **inverse system**, and the group of all such coherent sequences (ξ_ν) is called the **inverse limit**, denoted as

$$\hat{G} = \varprojlim G/G_n.$$

A direct (or inverse) system is called **injective/surjective system**, if every transition map μ_{ij} is injective/surjective.

Proposition 10.2. *If $0 \rightarrow \{A_n\} \rightarrow \{B_n\} \rightarrow \{C_n\} \rightarrow 0$ is an exact sequence of inverse system, then*

$$0 \rightarrow \varprojlim A_n \rightarrow \varprojlim B_n \rightarrow \varprojlim C_n$$

is exact. If $\{A_n\}$ is surjective, then

$$0 \rightarrow \varprojlim A_n \rightarrow \varprojlim B_n \rightarrow \varprojlim C_n \rightarrow 0$$

is exact.

Let $A = \prod_{i=1}^{\infty} A_n$ and define $d^A : A \rightarrow A$ by $d^A(a_n) = a_n - \theta_{n+1}(a_{n+1})$. Then $\ker(d^A) \cong \varprojlim A_n$. Similar to B and C , we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \\ & & \downarrow d^A & & \downarrow d^B & & \downarrow d^C \\ 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \end{array}$$

By [Snake lemma](#), we have an exact sequence

$$0 \longrightarrow \ker(d^A) \longrightarrow \ker(d^B) \longrightarrow \ker(d^C) \longrightarrow \operatorname{coker}(d^A) \longrightarrow \operatorname{coker}(d^B) \longrightarrow \operatorname{coker}(d^C) \longrightarrow 0.$$

and the $\operatorname{coker}(d^A)$ is usually denoted by $\varprojlim^1 A_n$.

Corollary 10.3. *Let $0 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 0$ be an exact sequence of groups. Let G have the topology defined by a sequence $\{G_n\}$ of subgroups, and give G' and G'' the topologies by the contraction and extension of $\{G_n\}$, respectively. Then*

$$0 \rightarrow \hat{G}' \rightarrow \hat{G} \rightarrow \hat{G}'' \rightarrow 0$$

is exact.

Corollary 10.4. $\hat{G}_n$ is a subgroup of $\hat{G}$ and $\hat{G}/\hat{G}_n \cong G/G_n$.

Corollary 10.5. $\hat{\hat{G}}_n \cong \hat{G}_n$.

If $G \rightarrow \hat{G}$ is an isomorphism, G is said to be [complete](#).

Let A be a ring, $\mathfrak{a}$ its ideal. The topology defined by $G_n = \mathfrak{a}^n$ ($G_0 = A$) is called the [\$\mathfrak{a}\$ -adic topology](#), and the ring A combined with this topology is a [topological ring](#).

FILTRATIONS

A chain $M = M_0 \supseteq M_1 \supseteq \cdots \supseteq M_n \cdots$ is called a [filtration](#) of M , and denoted by (M_n) . It is an [\$\mathfrak{a}\$ -filtration](#) if $\mathfrak{a}M_n \subseteq M_{n+1}$ for all n , and a [stable \$\mathfrak{a}\$ -filtration](#) if $\mathfrak{a}M_n = M_{n+1}$ for all sufficiently large n .

The [\$\mathfrak{a}\$ -adic completion](#) of M is defined to be

$$\hat{M} = \varprojlim M/\mathfrak{a}^n M.$$

GRADED RINGS AND MODULES

A [graded ring](#) is a ring A together with a family $(A_n)_{n \geq 0}$ of additive subgroups of A , such that $A = \bigoplus_{n=0}^{\infty} A_n$ and $A_m A_n \subseteq A_{m+n}$ for all $m, n \geq 0$.

Proposition 10.7. *A graded ring A is Noetherian, iff A_0 is Noetherian and A is finitely generated as an A_0 -algebra.*

Proposition 10.9 (Artin-Rees lemma). *Let A be a Noetherian ring, $\mathfrak{a}$ an ideal in A , M a finitely-generated A -module, (M_n) a stable $\mathfrak{a}$ -filtration of M . If M' is a submodule of M , then $(M' \cap M_n)$ is a stable $\mathfrak{a}$ -filtration of M' .*

Proposition 10.12. *Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence of finitely-generated modules over a Noetherian ring A , then the sequence of $\mathfrak{a}$ -adic completion $0 \rightarrow \hat{M}' \rightarrow \hat{M} \rightarrow \hat{M}'' \rightarrow 0$ is exact.*

Proposition 10.13. *Let $f : \hat{A} \otimes_A M \rightarrow \hat{A} \otimes_A \hat{M} \rightarrow \hat{A} \otimes_{\hat{A}} \hat{M} \rightarrow \hat{M}$. If M is finitely-generated A -module, f is surjective. If moreover A is Noetherian, f is an isomorphism.*

Proposition 10.14. *If A is a Noetherian ring, the $\mathfrak{a}$ -adic completion $\hat{A}$ is a flat A -algebra.*

Proposition 10.15. *If A is a Noetherian ring, $\hat{A}$ its $\mathfrak{a}$ -adic completion, then*

- (i) $\hat{\mathfrak{a}} = \hat{A}\mathfrak{a} \cong \hat{A} \otimes_A \mathfrak{a}$;
- (ii) $\widehat{(\mathfrak{a}^n)} = (\hat{\mathfrak{a}})^n$;
- (iii) $\hat{\mathfrak{a}}^n / \hat{\mathfrak{a}}^{n+1} \cong \mathfrak{a}^n / \mathfrak{a}^{n+1}$;
- (iv) $\hat{\mathfrak{a}}$ is contained in the Jacobson radical of $\hat{A}$.

Proposition 10.16. *Let A be a Noetherian local ring, $\mathfrak{m}$ its maximal ideal. Then the $\mathfrak{m}$ -adic completion $\hat{A}$ of A is a local ring with maximal ideal $\hat{\mathfrak{m}}$.*

Proposition 10.17 (Krull intersection theorem). *Let A be a Noetherian ring, $\mathfrak{a}$ an ideal, M a finitely-generated A -module and $\hat{M}$ the $\mathfrak{a}$ -adic completion of M . Then the kernel $\bigcap_{n=1}^{\infty} \mathfrak{a}^n M$ of $M \rightarrow \hat{M}$ consists of all $x \in M$ annihilated by some element of $1 + \mathfrak{a}$.*

THE ASSOCIATED GRADED RING

Let A be a ring and $\mathfrak{a}$ an ideal of A . The **associated graded ring** is the graded ring

$$G(A) (= G_{\mathfrak{a}}(A)) = \bigoplus_{n=0}^{\infty} \mathfrak{a}^n / \mathfrak{a}^{n+1} \quad (a^0 = A).$$

Similarly, if M is an A -module and (M_n) is an $\mathfrak{a}$ -filtration of M , then the **associated graded module** is

$$G(M) = \bigoplus_{n=0}^{\infty} M_n / M_{n+1},$$

which is a graded $G(A)$ -module. Let $G_n(M) = M_n / M_{n+1}$.

Proposition 10.22. *Let A be a Noetherian ring, $\mathfrak{a}$ an ideal of A . Then*

- (i) $G_{\mathfrak{a}}(A)$ is Noetherian;
- (ii) $G_{\mathfrak{a}}(A)$ and $G_{\hat{\mathfrak{a}}}(\hat{A})$ are isomorphic as graded rings;
- (iii) If M is a finitely-generated A -module and (M_n) is a stable $\mathfrak{a}$ -filtration of M , then $G(M)$ is a finitely-generated graded $G_{\mathfrak{a}}(A)$ -module.

Proposition 10.23. *Let $\phi : A \rightarrow B$ be a homomorphism of filtered groups, i.e. $\phi(A_n) \subseteq B_n$, and let $G(\phi) : G(A) \rightarrow G(B)$, $\hat{\phi} : \hat{A} \rightarrow \hat{B}$ be the induced homomorphisms of the associated graded and completed groups. Then*

- (i) $G(\phi)$ injective $\Rightarrow \hat{\phi}$ injective;
- (ii) $G(\phi)$ surjective $\Rightarrow \hat{\phi}$ surjective.

Proposition 10.24. *Let A be a ring, $\mathfrak{a}$ an ideal of A , M an A -module, (M_n) an $\mathfrak{a}$ -filtration of M . Suppose that A is complete in the $\mathfrak{a}$ -topology and that $\bigcap_n M_n = 0$. Suppose also $G(M)$ is a finitely generated $G(A)$ -module. Then M is a finitely generated A -module.*

Theorem 10.26. *If A is a Noetherian ring, $\mathfrak{a}$ an ideal of A , then the $\mathfrak{a}$ -adic completion $\hat{A}$ of A is Noetherian.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A Noetherian topological ring in which the topology is defined by an ideal contained in the Jacobson radical is called a **Zariski ring**.

Proposition (Hensel's lemma). *Let A be a local ring, $\mathfrak{m}$ its maximal ideal. Assume that A is $\mathfrak{m}$ -adically complete. If $f \in A[x]$ is monic of degree n , and if there exist coprime monic polynomials $\bar{g}, \bar{h} \in (A/\mathfrak{m})[x]$ of degrees $r, n-r$, with $\bar{f} = \bar{g}\bar{h}$, then we can lift $\bar{g}, \bar{h}$ back to monic polynomials $g, h \in A[x]$ such that $f = gh$.*

11 Dimension Theory

HILBERT FUNCTIONS

Let A be a Noetherian graded ring and M be a finitely-generated graded A -module. Let λ be an additive function on the class of all finitely generated A_0 -modules. The **Poincaré series** of M is the power series

$$P(M, t) = \sum_{n=0}^{\infty} \lambda(M_n) t^n \in \mathbb{Z}[[t]].$$

Theorem 11.1 (Hilbert-Serre theorem). $P(M, t)$ is a rational function in t of the form $f(t) / \prod_{i=1}^d (1 - t^{k_i})$, where $f(t) \in \mathbb{Z}[t]$.

The order of the pole of $P(M, t)$ at $t = 1$ is denoted as $d(M)$.

Corollary 11.2. If each $k_i = 1$, then for all sufficient large n , $\lambda(M_n)$ is a polynomial in n . More specifically, if $f(t) = \sum_{k=0}^N a_k t^k$, we have

$$\lambda(M_n) = \sum_{k=0}^n a_k \binom{d+n-k-1}{d-1}.$$

This is called **Hilbert function** or **Hilbert polynomial** of M , and its leading term is $(\sum a_k) n^{d-1} / (d-1)!$.

Proposition 11.3. If $x \in A_k$ is not a zero-divisor in M then $d(M/xM) = d(M) - 1$.

Proposition 11.4. Let A be a Noetherian local ring, $\mathfrak{m}$ its maximal ideal, $\mathfrak{q}$ an $\mathfrak{m}$ -primary ideal, M a finitely-generated A -module, (M_n) a stable $\mathfrak{q}$ -filtration of M . Then

- (i) M/M_n is of finite length for each $n \geq 0$;
- (ii) for all sufficiently large n this length is a polynomial $\chi_{\mathfrak{q}}^M(n)$ of degree $\leq s$, where s is the least number of generators of $\mathfrak{q}$;
- (iii) the degree and leading coefficient of $\chi_{\mathfrak{q}}^M(n)$ depend only on M and $\mathfrak{q}$, not on the filtration chosen.

If $M = A$, the polynomial $\chi_{\mathfrak{q}}^M(n)$ is called the **characteristic polynomial** of the $\mathfrak{m}$ -primary ideal $\mathfrak{q}$, simply noted as $\chi_{\mathfrak{q}}(n)$.

Proposition 11.6. $\deg \chi_{\mathfrak{q}}(n) = \deg \chi_{\mathfrak{m}}(n)$.

If A is a ring, $\mathfrak{p}$ a prime ideal in A , then the **height** of $\mathfrak{p}$ is defined to be the supremum of chains of prime ideals $\mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \cdots \subsetneq \mathfrak{p}_r = \mathfrak{p}$ which end at $\mathfrak{p}$. Hence $\text{height } \mathfrak{p} = \dim A_{\mathfrak{p}}$.

Likewise we define the **depth** of $\mathfrak{p}$, by considering chains of prime ideals which start at $\mathfrak{p}$. Hence, $\text{depth } \mathfrak{p} = \dim A/\mathfrak{p}$.

DIMENSION THEORY OF NOETHERIAN LOCAL RINGS

Theorem 11.14 (Dimension theorem). For any Noetherian local ring A the following three integers are equal:

- (i) the maximum length of chains of prime ideals in A (**Krull dimension**);
- (ii) the degree of the characteristic polynomial $\chi_{\mathfrak{m}} = l(A/\mathfrak{m}^n)$;

(iii) the least number of generators of an $\mathfrak{m}$ -primary ideal of A .

Corollary 11.17 (Krull's principal ideal theorem). *Let A be a Noetherian ring and let x be an element of A which is neither a zero-divisor nor a unit. Then every minimal prime ideal $\mathfrak{p}$ of (x) has height 1.*

Corollary 11.18. *Let A be a Noetherian local ring, x an element of $\mathfrak{m}$ which is not a zero-divisor. Then $\dim A/(x) = \dim A - 1$.*

Corollary 11.19. *Let $\hat{A}$ be the $\mathfrak{m}$ -adic completion of A . Then $\dim A = \dim \hat{A}$.*

If $x_1, \dots, x_d$ generate an $\mathfrak{m}$ -primary ideal, and $d = \dim A$, we call $x_1, \dots, x_d$ a **system of parameters**.

Proposition 11.20. *Let $x_1, \dots, x_d$ be a system of parameters for A and let $\mathfrak{q} = (x_1, \dots, x_d)$ be the $\mathfrak{m}$ -primary ideal. Let $f(t_1, \dots, t_d)$ be a homogeneous polynomial of degree s with coefficients in A , and assume that $f(x_1, \dots, x_d) \in \mathfrak{q}^{s+1}$. Then all the coefficients of f lie in $\mathfrak{m}$.*

REGULAR LOCAL RINGS

Theorem 11.22. *Let A be a Noetherian local ring of dimension d , $\mathfrak{m}$ its maximal ideal, $k = A/\mathfrak{m}$. Then the following are equivalent:*

- (i) $G_{\mathfrak{m}}(A) \cong k[t_1, \dots, t_d]$ where the t_i are independent indeterminates;
- (ii) $\dim_k(\mathfrak{m}/\mathfrak{m}^2) = d$;
- (iii) $\mathfrak{m}$ can be generated by d elements.

The ring satisfying any of the conditions is called **regular** local ring.

Proposition 11.23. *Let A be a ring, $\mathfrak{a}$ an ideal of A such that $\bigcap_n \mathfrak{a}^n = 0$. Suppose that $G_{\mathfrak{a}}(A)$ is an integral domain. Then A is an integral domain.*

Proposition 11.24. *Let A be a Noetherian local ring. Then A is regular iff $\hat{A}$ is regular.*

TRANSCENDENTAL DIMENSION

Assume that k is an algebraically closed field and let V be an irreducible affine variety over k . Let the coordinate ring of V be $A(V) = k[x_1, \dots, x_n]/I(V)$. Then $k(V) = \text{Frac } A(V)$ is a finitely generated extension of k and so has a finite transcendence degree over k , which is defined to be the **dimension** of V .

If P is a point with maximal ideal $\mathfrak{m}$ we call the Krull dimension $\dim A(V)_{\mathfrak{m}}$ the **local dimension** of V at P .

Theorem 11.25. *For any irreducible variety V over k the local dimension at any point is equal to its dimension.*

EXERCISES

1 Rings and Ideals

1.1. Consider $(1+x)(1-x+x^2-\cdots+(-x)^{n-1})$.

1.2. (i) Passing to $(A/\mathfrak{p})[x]$. $A/\mathfrak{p}$ is an integral domain, so $\deg(\overline{f\bar{g}}) = \deg(\overline{f}) + \deg(\overline{g})$. $\overline{f\bar{g}} = 1$ means $a_1, \dots, a_n \in \mathfrak{p}$. Use [Proposition 1.8](#).

(ii) a, b are nilpotents $\Rightarrow a+b$ is nilpotent. Note that $f-a_0 = xf_1$ is nilpotent.

(iii) Let $gf=0$ and g is of minimal degree. The leading coefficient $a_nb_m=0$, so a_ng has less degree than g and so $a_ng=0$. Same process to the second leading and following coefficients, we have $a_i g=0$ for all i .

(iv) (Gauss's lemma) Let $\mathfrak{a}, \mathfrak{b}$ and $\mathfrak{c}$ are generated by the coefficients of f, g and fg , respectively. We need to proof $\mathfrak{c} = (1) \Leftrightarrow \mathfrak{a}\mathfrak{b} = (1)$. $\mathfrak{c} \subseteq \mathfrak{a}\mathfrak{b}$ by the form of coefficients in fg . If $\mathfrak{c} \subsetneq \mathfrak{a}\mathfrak{b} = (1)$, let $\mathfrak{m}$ be the maximal ideal containing $\mathfrak{c}$, then $(A/\mathfrak{m})[x]$ is an integer domain. $\overline{f\bar{g}} = 0$ but $\overline{f}, \overline{g} \neq 0$.

1.3. Induction on the indeterminates.

1.4. Pick the y in [Proposition 1.9](#) to be the indeterminate, then use [Exercise 1.2\(i\)](#).

1.5. (i) $\Leftarrow$: $fg = \sum_{i=0}^{\infty} \left(\sum_{j=0}^i a_j b_{i-j} \right) x^i$, so we can find g such that $\sum_{j=0}^i a_j b_{i-j} = 0$.

(ii) Similar to [Exercise 1.2\(ii\)](#). Construct a counterexample where a_n have increasing nilpotencies.

(iii) Pick the y in [Proposition 1.9](#) to be 1. Use (i) on $1-f \Leftrightarrow 1-a_0$.

(iv),(v) $A \cong A[[x]]/(x)$. Use [Proposition 1.1](#).

1.6. $\mathfrak{J}$ is an ideal, so there is an idempotent $e \in \mathfrak{J}$ but $e \notin \mathfrak{R}$. Pick the y in [Proposition 1.9](#) to be 1.

1.7. Note that $x(1-x^{n-1}) = 0 \in \mathfrak{p}$.

1.8. Zorn's lemma but define the partial order reversely.

1.9. $\Rightarrow$: Apply [Proposition 1.8](#) to $A/\mathfrak{a}$.

$\Leftarrow$: Use [Proposition 1.11\(ii\)](#).

1.10. Use [Corollary 1.5](#) and [Corollary 1.6\(i\)](#)

1.11. (i) $x+x = (x+x)^2 = x^2+2x+x^2 = x+2x+x$.

(ii) Prove that every element in $A/\mathfrak{p}$ is 1.

(iii) $x+y-xy$.

1.12. Use [Corollary 1.5](#).

1.13. **1 is a monic polynomial but it is neither irreducible nor reducible.** A is an integral domain, so $\deg(fg) = \deg(f) + \deg(g)$. If $\mathfrak{a} = (1)$, apply the equation to $1 = \sum_f y_f f(x_f)$.

1.14. If $xy \in \mathfrak{m}$, xy is zero-divisor, so are x and y . By maximality, $\mathfrak{m} \subseteq \mathfrak{m} + (x) = \mathfrak{m}$.

1.15. (i) Apply [Proposition 1.8](#) to $A/\mathfrak{a}$ to proof the second equation.

(iv) Use $\sqrt{\mathfrak{a} \cap \mathfrak{b}} = \sqrt{\mathfrak{a}\mathfrak{b}}$.

1.16. Irreducible polynomials relate to prime ideals.

1.17. (i)-(iv) Use [Exercise 1.15](#).

(v) Let $(X_{f_i})_{i \in I}$ be an open cover. By [Exercise 1.15](#), $V(\{f_i\}_{i \in I}) = \emptyset$, so $\{f_i\}_{i \in I}$ generates the unit ideal. Let $1 = \sum_{i \in J} g_i f_i$ where J is a finite subset of I . If $\mathfrak{p} \notin \bigcup_{i \in J} X_{f_i}$, then $f_i \in \mathfrak{p}$ for all $i \in J$.

EXERCISES

- (vi) Similar to (v), where the unit ideal shall change to the radical $\sqrt{(f)}$.
 (vii) $\Rightarrow$: The intersection can be reduced because of (i), so we can assume an infinite union $X = \bigcup X_f$.

1.18. (iv) If $V(\mathfrak{p}_x) \subseteq V(\mathfrak{p}_y)$, $\mathfrak{p}_x \in \overline{V(\mathfrak{p}_y)}$. If they don't have containing relation, $\mathfrak{p}_x \in \overline{V(\mathfrak{p}_y)}$ and $\mathfrak{p}_y \in \overline{V(\mathfrak{p}_x)}$.

1.19. Use the closed sets definition. Consider minimal prime ideals.

- 1.20.** (i) Let $\overline{Y} = Z_1 \cup Z_2$, then $Z_i \cap Y = \emptyset$ for an $i \in \{1, 2\}$. Thus $Z_i = \overline{Z_i \cap Y} = \emptyset$.
 (ii) Similar to (i). Construct the maximal element and prove that it is irreducible.
 (iii) If Y Hausdorff but irreducible, every pair of open sets in Y intersect. Thus Y is singleton.
 (iv) Use the closed set definition.

1.21. Simplify the sets: $\phi^{*-1}(E) = \{\mathfrak{q} \in Y : \phi^{-1}(\mathfrak{q}) \in E\}$, $\phi^*(V(E)) = \{\phi^{-1}(\mathfrak{q}) : E \subseteq \mathfrak{q}\}$.

1.22. Use quotient to find the form of prime ideals in A .

- 1.23.** (i) $X_f = V(1 - f)$.
 (ii) $X_{f_1} \cup X_{f_2} = X_{f_1 + f_2 - f_1 f_2}$.
 (iii) A closed set in compact space is compact. By Exercise 1.17, it is a finite union of basic open set. Then apply (ii).
 (iv) By Exercise 1.11 every prime ideal is maximal. Pick $f \in \mathfrak{p}_x \setminus \mathfrak{p}_y$, then X_f and $V(f)$ separate two points.

1.24. $a = a \wedge 1 = a \wedge (a \vee a') = (a \wedge a) \vee (a \wedge a') = (a \wedge a) \vee 0 = a \wedge a$.

1.25. Use Exercise 1.23(iv).

1.26. $\mathfrak{m}_x \in \mu(U_f) \Leftrightarrow f(x) \neq 0 \Leftrightarrow f \notin \mathfrak{m}_x \Leftrightarrow \mathfrak{m}_x \in \tilde{U}_f$.

1.27. If $x \neq y$, there exists a polynomial $f \in P(X)$ such that $f(x) = 0$ but $f(y) \neq 0$, so $\mathfrak{m}_x \neq \mathfrak{m}_y$, and so μ injective. By Nullstellensatz, every maximal ideal in $P(X)$ is of the form $\mathfrak{m} = (t_1 - a_1, \dots, t_n - a_n)$, with $(a_1, \dots, a_n) \in X$, so μ surjective. By Exercise 1.26, μ is a homeomorphism.

1.28. Project to coordinate functions.

2 Modules

2.1. Bézout's identity.

2.2. Tensor the exact sequence $0 \rightarrow \mathfrak{a} \rightarrow A \rightarrow A/\mathfrak{a} \rightarrow 0$. Calculate the image of $\mathfrak{a} \otimes M \rightarrow A \otimes M$.

2.3. Consider the residue field k of A and pass $M \otimes_A N$ to vector space $M_k \otimes_k N_k$. Use Exercise 2.2 and Nakayama's lemma.

2.4. There exist canonical projections.

2.5. The definition of flatness only considers the module structure. Use Exercise 2.4.

2.6. Construct two homomorphisms.

2.7. Prove that $(A/\mathfrak{p})[x]$ and $A[x]/\mathfrak{p}[x]$ are isomorphic.

2.8. (ii) Use Proposition 2.15.

2.9. Pick $y_1, \dots, y_m \in M$ whose images are the generators of M'' . Use diagram chasing for an element of M .

2.10. $u(m) + \mathfrak{a}N \in N$. If $n \in N$, we have $n + \mathfrak{a}N = u(m + \mathfrak{a}M) = u(m) + \mathfrak{a}N$. Apply Nakayama's lemma on $N/u(M)$ where $\mathfrak{a}(N/u(M)) = (\mathfrak{a}N + u(M))/u(M)$.

2.11. (i),(ii) Consider a residue field and pass to vector spaces. Use the right exactness of tensor product.

(iii) If $m > n$, let $\psi : A^m \xrightarrow{\phi} A^n \hookrightarrow A^m$. By [Proposition 2.4](#), ψ satisfies $\psi^n + a_{n-1}\psi^{n-1} + \cdots + a_0 = 0$. Pick such equation to have the minimum degree, then ψ injective implies $a_0 \neq 0$. Apply the equation to element $(0, \dots, 0, 1)$ to obtain contradiction.

2.12. Choose $u_i \in M$ such that $\phi(u_i)$ are a basis of A^n , then prove that $0 \rightarrow \ker(\phi) \rightarrow M \xrightarrow{\phi} A^n \rightarrow 0$ splits.

2.13. $p \circ g = \text{id}_N$ injective $\Rightarrow g$ injective, so $0 \rightarrow N \xrightarrow{g} N_B \rightarrow N_B/g(N) \rightarrow 0$ splits.

2.14. This question asks to prove $\mu_i = \mu_j \circ \mu_{ij}$.

2.15. (i) Use [Exercise 2.14](#) to increase indices.

(ii) $x_i = \sum_k (y_k - \mu_{ki}(y_k))$ is an equation in $\bigoplus M_i$, so μ_{ij} cannot be applied directly. Use canonical projection $\pi_i : \bigoplus M_i \rightarrow M_i$ on the equation first.

2.16 (Universal property of direct limit). Use [Exercise 2.15\(i\)](#) for both existence and uniqueness.

2.17. Similar to [Exercise 2.15\(i\)](#), move components into a same index.

2.18. Use [Exercise 2.16](#).

2.19 ($\varinjlim$ is an exact functor). Diagram chasing. Use [Exercise 2.15](#).

2.20. Draw the diagram. Define two homomorphisms by the universal properties of tensor product and direct limit. Verify their compositions are identity maps.

2.21. Use [Exercise 2.15\(ii\)](#) on $\alpha_i(1)$.

2.22. Use [Exercise 2.15](#). Pick a nilpotent or assume $xy = 0$.

2.23. If $J \subseteq J'$, $B_J \cong B_J \otimes (A)_{i \in J' \setminus J}$. Let Γ be the set of all finite subsets of Λ , then $(B_J)_{J \in \Gamma}$ is the case of [Exercise 2.21](#). We can also use [Exercise 2.17](#) to see its form.

2.24. Directly from the definition of Tor functor.

2.25. Use [Snake lemma](#).

2.26. Let M_n be generated by $\{x_1, \dots, x_n\}$. We have an exact sequence

$$\text{Tor}_1(M_{n-1}, N) \rightarrow \text{Tor}_1(M_n, N) \rightarrow \text{Tor}_1(M_n/M_{n-1}, N).$$

By [Proposition 2.3](#), $M_n/M_{n-1} \cong A/\mathfrak{a}$.

Let $f : \mathfrak{a} \otimes N \rightarrow A \otimes N$, $f(\sum a_i \otimes y_i) = 0$. Let $\mathfrak{a}'$ be generated by a_i . By assumption, $f|_{\mathfrak{a}' \otimes N}$ is injective, so $\sum a_i \otimes y_i = 0$ and thus $\text{Tor}_1(A/\mathfrak{a}, N) = 0$. Since $M_0 = A$ and $\text{Tor}_1(A, N) = 0$, the exact sequence implies $\text{Tor}_1(M_n, N) = 0$ for all n . Then use [Proposition 2.19\(iv\)](#).

Another approach is to use the fact that any module is the direct limit of some finitely generated modules, and direct limit is an exact functor.

2.27. (i) $\Rightarrow$ (ii) $A/(x)$ is a flat A -module, then $\phi : A/(x) \otimes (x) \rightarrow A/(x)$ injective. $\phi(1 \otimes x) = x = 0$, so $A/(x) \otimes (x) = 0$. Then use [Exercise 2.2](#).

(ii) $\Rightarrow$ (iii) Let $x = ax^2$, then ax is an idempotent. $(x) \ni bx = bax^2 = (bx)(ax) \in (ax)$, so $(ax) = (x)$. Similar to [Exercise 1.11\(iii\)](#), every finitely generated ideal is generated by an idempotent e , so $A = (e) \oplus (1 - e)$.

(iii) $\Rightarrow$ (i) $N \otimes_A A = (N \otimes_A \mathfrak{a}) \oplus (N \otimes_A \mathfrak{a}')$, so $(N \otimes_A \mathfrak{a}) \rightarrow N \otimes_A A$ injective. Use [Exercise 2.26](#).

2.28. Use [Exercise 2.27\(ii\)](#).

3 Rings and Modules of Fractions

3.1. The only if part requires the finitely generated property.

3.2. Use [Proposition 1.9](#). Construct the inverse by elementary algebra.

If $M = \mathfrak{a}M$, $S^{-1}M = S^{-1}\mathfrak{a}$, $S^{-1}M = (S^{-1}\mathfrak{a})(S^{-1}M)$. Then use [Nakayama's lemma](#) and [Exercise 3.1](#).

3.3. Construct the isomorphism by $(a/s)/(t/1) \mapsto a/(st)$.

3.4. Construct the isomorphism.

3.5. If $a \in A$ is a nilpotent, let $\mathfrak{p} \supseteq \text{Ann}(a)$. $a/1 \in A_{\mathfrak{p}}$ is zero, so $sa = 0$ for some $s \in A - \mathfrak{p}$.

Counter example is $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.

3.6. Use the fact that for any multiplicatively closed set S , if $A - S$ is an ideal, it's automatically prime.

3.7. (i),(ii) More specifically, $A - \overline{S} = \bigcup_{\mathfrak{p} \cap S = \emptyset} \mathfrak{p}$.

(iii) $\mathfrak{p} \cap (1 + \mathfrak{a}) = \emptyset \Leftrightarrow \mathfrak{a} + \mathfrak{p} \neq (1) \Leftrightarrow \mathfrak{a} + \mathfrak{p} \subseteq \mathfrak{m}$ for some maximal ideal $\mathfrak{m}$. Note that the union of a chain of prime ideals is only a maximum ideal.

3.8. (v) $\Rightarrow$ (ii) If $t/1$ is not a unit in $S^{-1}A$, it is contained in a prime ideal $\mathfrak{q} \subseteq S^{-1}A$, so $t \in T \cap \mathfrak{q}^c$.

3.9. Let S be maximal in [Exercise 3.6](#). If $S_0 \not\subseteq S$, $S \subsetneq SS_0$. Then use [Exercise 3.6](#).

(i) Assume that S_1 is larger, then pick a zero-divisor $x \in S_1$.

(ii) If a/s is not a zero-divisor, a is not a zero-divisor, so $a \in S_0$.

(iii) Note that $s \in S_0$ is a unit.

3.10. (i) [Exercise 2.27](#) says for any element $x \in A$, $x = ax^2$ for some $a \in A$.

(ii) $\Rightarrow$: Use [Exercise 2.28](#). $\Leftarrow$: Consider an $A_{\mathfrak{m}}$ -module. Use [Proposition 3.10](#).

3.11. (i) $\Rightarrow$ (ii) Use [Exercise 2.27](#) on $A/\mathfrak{p}$.

(ii) $\Rightarrow$ (i) Prove that $(A/\mathfrak{A})_{\overline{\mathfrak{p}}}$ has no nilpotent, then use [Exercise 1.10](#) and [Exercise 3.10](#).

(ii) $\Rightarrow$ (iv) Let $f \in \mathfrak{p}_1$ but $f \notin \mathfrak{p}_2$, then $f/1 \in A_{\mathfrak{p}_1}$ is nilpotent, so $sf^n = 0$. We have $\mathfrak{p}_1 \in X_s$ and $\mathfrak{p}_2 \in X_{f^n}$. By [Exercise 1.17](#), $X_s \cap X_{f^n} = X_{sf^n} = \emptyset$.

3.12. (i)-(iii) Use the torsion element definition.

(iv) Use [Proposition 3.5](#).

3.13. Prove they contain in each other. Use [Proposition 3.8](#) for the equivalent conditions.

3.14. Pass to $M/\mathfrak{a}M$. Use [Proposition 3.8](#).

3.15. Let $\{x_i\}_{i=1}^n$ be generators and $\{e_i\}_{i=1}^n$ be the canonical basis. Let $\phi(e_i) = x_i$, so ϕ is surjective.

Consider a residue field $k = A/\mathfrak{m}$, we have $\text{Tor}_1(k, F) \rightarrow k \otimes \ker \phi \rightarrow k \otimes F \xrightarrow{1 \otimes \phi} k \otimes F \rightarrow 0$. $\text{Tor}_1(k, F) = 0$ by balancing Tor theorem; $1 \otimes \phi$ bijective so $k \otimes \ker \phi = 0$. By [Exercise 2.3](#), $\ker \phi = 0$.

If $\{x_i\}_{i=1}^m$ generate F where $m < n$, pick some element $\{x_i\}_{i=m+1}^n$ so we also have $\phi(e_i) = x_i$. By injectiveness of ϕ , $\{e_i\}_{i=1}^n$ are linearly dependent.

3.16. (i) $\Rightarrow$ (ii) $f^*(p^e) = p$.

(ii) $\Rightarrow$ (iii) Use [Proposition 1.17](#).

(iii) $\Rightarrow$ (iv) Every module contains a cyclic submodule, which is isomorphic to $A/\mathfrak{a}$. Then use $B \otimes_A A/\mathfrak{a} \cong B/\mathfrak{a}^e$ and $\mathfrak{a}^e \subseteq \mathfrak{m}^e$.

(iv) $\Rightarrow$ (v) Let the mapping be ϕ . Since B flat, $0 \rightarrow (\ker \phi)_B \rightarrow M_B \rightarrow (M_B)_B$ exact. By [Exercise 2.13](#), $M_B \rightarrow (M_B)_B$ injective, so $(\ker \phi)_B = 0$; by assumption, $\ker \phi = 0$.

(v) $\Rightarrow$ (i) Take $M = A/\mathfrak{a}$. Then $M_B = B/\mathfrak{a}^e$.

3.17. If $M \rightarrow N$ injective, we have a diagram

$$\begin{array}{ccc} B \otimes_A M & \xrightarrow{\phi} & B \otimes_A N \\ \downarrow g \otimes \text{id}_M & & \downarrow g \otimes \text{id}_N \\ C \otimes_A M & \xrightarrow{\psi} & C \otimes_A N \end{array}$$

$g \circ f$ flat, so ψ injective. g faithfully flat, by [Exercise 3.16\(v\)](#), $B \otimes_A M \rightarrow C \otimes_B (B \otimes_A M)$ injective. By [Proposition 2.15](#), $C \otimes_B (B \otimes_A M) = C \otimes_A M$, so $g \otimes \text{id}_M$ injective.

3.18. Use [Exercise 3.3](#) to prove $B_{\mathfrak{q}} \cong (B_{\mathfrak{p}})_{\mathfrak{q}B_{\mathfrak{p}}}$. By [Proposition 3.3](#) and [Proposition 3.5](#), $B_{\mathfrak{q}}$ is flat over $B_{\mathfrak{p}}$. Hence $B_{\mathfrak{q}}$ is flat over $A_{\mathfrak{p}}$ and satisfies [Exercise 3.16\(iii\)](#).

3.19. (v) $M \cong \bigoplus_{i=1}^n A/\mathfrak{a}_i$, so $\text{Supp}(M) = \bigcup_{i=1}^n \text{Supp}(A/\mathfrak{a}_i) = \bigcup_{i=1}^n V(\mathfrak{a}_i) = V(\bigcap_{i=1}^n \mathfrak{a}_i)$. Prove that $\bigcap_{i=1}^n \mathfrak{a}_i = \text{Ann}(M)$.

(vi) Use [Exercise 2.3](#).

(vii) Use [Exercise 2.2](#).

(viii) $\mathfrak{q} \in f^{*-1}(\text{Supp}(M)) \Leftrightarrow M_{\mathfrak{q}^c} \neq 0 \Leftrightarrow M_{\mathfrak{p}} \neq 0$ and $B_{\mathfrak{q}} \neq 0 \Leftrightarrow \mathfrak{q} \in \text{Supp}(M) \cap \text{Supp}(B)$.

3.20. Counter example is $A[x]/(x^2)$.

3.21. (i),(ii) Use [Proposition 3.11\(iv\)](#).

(iii) Use [Exercise 1.21\(iv\)](#).

(iv) To prove $f^{*-1}(\mathfrak{p}) = \text{Spec}(B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}})$, use $\mathfrak{q} \in \text{Spec}(S^{-1}B) \Leftrightarrow \mathfrak{q} \cap f(S) = \emptyset$ and $\mathfrak{q} \in \text{Spec}(C/\mathfrak{p}C) \Leftrightarrow f^{-1}(\mathfrak{q}) \supseteq \mathfrak{p}$. To prove the equation, note that $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} = \text{Frac}(A/\mathfrak{p})$, then apply [Proposition 2.15](#) on $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \otimes_{A/\mathfrak{p}} A/\mathfrak{p} \otimes_A B$.

3.22. Let $\mathfrak{p} \in X_f$, so $f \notin \mathfrak{p}$. Then $\mathfrak{q} \in \text{Spec}(A_{\mathfrak{p}}) \Leftrightarrow \mathfrak{q} \subseteq \mathfrak{p} \Leftrightarrow f \notin \mathfrak{q} \Leftrightarrow \mathfrak{q} \in X_f$.

3.23. (i) Use [Exercise 1.17\(iv\)](#).

(v) Define two homomorphisms by the universal properties of direct limit and local ring.

3.24. [Exercise 1.17\(v\)](#) asserts that there exists a finite open cover (X_{f_i}) such that $\sum_{i=1}^n (f_i) = (f_1, \dots, f_n) = (1)$, hence $\sqrt{\sum_{i=1}^n (f_i^N)} = \sqrt{\sum_{i=1}^n \sqrt{(f_i^N)}} = (1)$ and therefore $\sum_{i=1}^n (f_i^N) = (1)$ for all N . We can find b_i such that $\sum_{i=1}^n b_i f_i^N = 1$.

Pick N large enough and let $s_i = a'_i/f_i^{m_i} = a_i/f_i^N$. Define $s = \sum_{i=1}^n a_i b_i$. Note that $a_i/f_i^N = a_j/f_j^N$ in $A(U_i \cap U_j)$, and use it to verify s satisfies the condition.

If $t = s - s'$, $t/1 = 0$ in all A_f . Pick f_i are those correspond to the open cover, so $f_i^M t = 0$ for a large M . Then $t = \sum_{i=1}^n b_i f_i^N t = 0$.

3.25. The h is not a homomorphism. Redefine $h(x) = f(x) \otimes 1 = 1 \otimes g(x)$. We have fibers

$$\begin{aligned} f^{*-1}(\mathfrak{p}) &= \text{Spec}(B \otimes_A k(\mathfrak{p})) & g^{*-1}(\mathfrak{p}) &= \text{Spec}(C \otimes_A k(\mathfrak{p})) \\ h^{*-1}(\mathfrak{p}) &= \text{Spec}((B \otimes_A C) \otimes_A k(\mathfrak{p})) \cong \text{Spec}((B \otimes_A k(\mathfrak{p})) \otimes_{k(\mathfrak{p})} (C \otimes_A k(\mathfrak{p}))) \end{aligned}$$

Note that $B \otimes_A k(\mathfrak{p})$ and $C \otimes_A k(\mathfrak{p})$ are vector space, so $\mathfrak{p} \in h^*(T)$ iff $(B \otimes_A k) \otimes_k (C \otimes_A k) \neq 0$, iff $B \otimes_A k \neq 0$ and $C \otimes_A k \neq 0$, iff $\mathfrak{p} \in f^*(Y)$ and $\mathfrak{p} \in g^*(Z)$.

3.26. The fiber $f^{*-1}(\mathfrak{p})$ is the spectrum of $B \otimes_A k(\mathfrak{p})$. By [Exercise 2.20](#), $B \otimes_A k(\mathfrak{p}) = \varinjlim (B_{\alpha} \otimes_A k(\mathfrak{p}))$. By [Exercise 2.21](#), $f^{*-1}(\mathfrak{p}) = \emptyset \Rightarrow B_{\alpha} \otimes_A k(\mathfrak{p}) = 0$ for some α , and the reverse is true by the definition of direct limit of rings.

3.27. (i) Use [Exercise 3.25](#) for finite case. Expand to infinite case by [Exercise 2.23](#), then use [Exercise 3.26](#).

(ii) Use [Exercise 1.22](#).

(iii) $V(\mathfrak{a}) = f^*(\text{Spec}(A/\mathfrak{a}))$, but $f^*(\text{Spec}(S^{-1}A))$ is also closed in constructible topology.

(iv) If $\{X \setminus f^*(\text{Spec}(B_\alpha))\}_\alpha$ is an open cover, $\bigcap_\alpha f^*(\text{Spec}(B_\alpha)) = \emptyset$, so $B = 0$. We can still think B is constructed by [Exercise 2.23](#), then apply [Exercise 2.21](#) we have $B_J = 0$ for a finite set J .

3.28. (iii) We claim that any continuous bijection from a compact space to a Hausdorff space is a homeomorphism, so is $X_C \rightarrow X_{C'}$. Let $f : X \rightarrow Y$ be such map and let $C \subseteq X$ closed. Then C is compact, so is $f(C)$. Since Y Hausdorff, $f(C)$ is closed and therefore f maps closed sets to closed sets.

3.29. Consider $g : B \rightarrow C$. Use [Exercise 1.21\(vi\)](#).

3.30. $\leftarrow \text{id} : X_C \rightarrow X$ is a map from a compact space to a Hausdorff space. The topology on X_C is finer than X so the map is continuous. Then use the claim in [Exercise 3.28\(iii\)](#).

4 Primary Decomposition

4.1. Use [Exercise 1.21\(iv\)](#), so $\text{Spec}(A/\mathfrak{a}) = V(\mathfrak{a})$.

4.2. All primary components are prime.

4.3. For any element $x \in A$, $x(1 - ax) = 0 \in \mathfrak{q}$.

4.4. $\mathfrak{m} = \sqrt{\mathfrak{q}} \supsetneq \mathfrak{q} \supsetneq \mathfrak{m}^2$.

4.5. $\mathfrak{m}^2$ is $\mathfrak{m}$ -primary.

4.6. Use Urysohn's lemma to prove that if $f(x_1) = f(x_2) = 0$ for all $f \in \mathfrak{q}$, $\mathfrak{q}$ cannot be primary, so there are infinite many minimal prime ideals. Now use [Proposition 1.11\(ii\)](#) to prove that $(0) = \sqrt{(0)} = \bigcap \mathfrak{p}$ is the intersection of ALL minimal prime ideals.

4.7. (ii),(iv),(v) $(\mathfrak{a}[x])^c = \mathfrak{a}$.

(iii) Let $f, \bar{g} \in (A/\mathfrak{q})[x]$. If $f\bar{g} = 0$ but $\bar{g} \neq 0$, by [Exercise 1.2\(iii\)](#), $\bar{a}\bar{f} = 0$, so $aa_i \in \mathfrak{q}$. Hence $a_i \in \mathfrak{p}$ and $a_i^{n_i} \in \mathfrak{q}$, so $\bar{a}_i^{n_i} = 0$ and use [Exercise 1.2\(ii\)](#).

4.8. $k[x_1, \dots, x_n]/\mathfrak{p}_i = k[x_{i+1}, \dots, x_n]$ integer domain, so $\mathfrak{p}_i$ prime. Note that $\mathfrak{p}_i^n$ is the set of all homogeneous polynomials in $k[x_1, \dots, x_i]$ of degree $\geq n$.

4.9. (i) $\Leftarrow$: x is contained in all prime ideals containing $(0 : a)$, so $x \in \sqrt{(0 : a)}$.

(ii) If $\mathfrak{a} \subseteq \mathfrak{p}$ but $\mathfrak{a}^e \subseteq S^{-1}\mathfrak{p}' \subsetneq S^{-1}\mathfrak{p}$, then $\mathfrak{a} \subseteq \mathfrak{a}^{ec} \subseteq \mathfrak{p}' \subsetneq \mathfrak{p}$. If $\mathfrak{b} \subseteq S^{-1}\mathfrak{p}$ but $\mathfrak{b}^c \subseteq \mathfrak{p}' \subsetneq \mathfrak{p}$, use $\mathfrak{b}^{ce} = \mathfrak{b}$ derived from [Proposition 3.11\(i\)](#). $S^{-1}\mathfrak{p}$ is a prime ideal if S doesn't meet $\mathfrak{p}$.

(iii) $(0 : a) = \bigcap (\mathfrak{q}_i : a) \subseteq \mathfrak{p}$. By [Proposition 1.11\(ii\)](#), $(\mathfrak{q}_i : a) \subseteq \mathfrak{p}$ for some i . Then use [Proposition 4.4](#) and the minimality of $\mathfrak{p}$.

4.10. (iv) Let $x \neq 0$, $(0 : x) \subseteq \mathfrak{p}$ for some $\mathfrak{p}$. If $x \in S_{\mathfrak{p}}(0)$, then $sx = 0$, so $s \in (0 : x)$.

4.11. Note that $x \in S(\mathfrak{a}) \Leftrightarrow$ there exists an $s \in S$ such that $sx \in \mathfrak{a}$. Show that $S_{\mathfrak{p}}(0)$ is contained in all $\mathfrak{p}$ -primary ideals.

If $0 = \bigcap \mathfrak{q}_i$ is irredundant, $\sqrt{(0)}$ is the intersection of minimal prime ideals.

4.12. Use [Proposition 3.11\(ii\)](#). Use [Proposition 4.9](#) to show there are at most 2^n different $S(\mathfrak{a})$.

4.13. (i) Use [Exercise 4.12](#).

(ii) Use [Proposition 4.8](#) and [Proposition 4.9](#).

(iii) Prove that $S_{\mathfrak{p}}(\mathfrak{p}^{(m)}\mathfrak{p}^{(n)}) = \mathfrak{p}^{(m+n)}$.

(iv) $\Leftarrow$: Show that $\mathfrak{p}^{(n)}$ is the smallest $\mathfrak{p}$ -primary ideal containing $\mathfrak{p}^n$.

4.14. Let $x \notin \mathfrak{q}$. If $ab \in (\mathfrak{q} : x)$ but $a \notin (\mathfrak{q} : x)$, we have $xab \in \mathfrak{q}$ but $xa \notin \mathfrak{q}$. Thus, $b \in (\mathfrak{q} : xa) = (\mathfrak{q} : x)$ by maximality, and therefore $(\mathfrak{q} : x)$ is prime.

4.15. Use [Proposition 4.9](#).

4.16. Use [Proposition 3.11\(iv\)](#) and [Proposition 4.9](#).

4.17. Take $\mathfrak{p}_1$ to be a minimal prime ideal containing $\mathfrak{a}_1 = \mathfrak{a}$. By [Exercise 4.11](#), $S_{\mathfrak{p}_1}(\mathfrak{a}_1)$ is $\mathfrak{p}_1$ -primary, and $S_{\mathfrak{p}_1}(\mathfrak{a}_1) = (\mathfrak{a}_1 : x_1)$ for some $x_1 \notin \mathfrak{p}_1$. If $u \in \mathfrak{q}_1 \cap (\mathfrak{a}_1 + (x_1))$, $u = a + tx_1$ and $ux_1 \in \mathfrak{a}_1$, so $ax_1 + tx_1^2 \in \mathfrak{a}_1$ and so $tx_1^2 \in \mathfrak{a}_1$. Since $x_1 \notin \mathfrak{p}_1$, $t \in (\mathfrak{a}_1 : x_1)$, which implies $f \in \mathfrak{a}_1$, and therefore $\mathfrak{q}_1 \cap (\mathfrak{a}_1 + (x_1)) \subseteq \mathfrak{a}_1$.

Let $\mathfrak{a}_n$ be a maximal element of the set $\{\mathfrak{b} \subseteq A : \mathfrak{a}_{n-1} + (x_{n-1}) \subseteq \mathfrak{b} \text{ and } \mathfrak{q}_{n-1} \cap \mathfrak{b} = \mathfrak{a}_{n-1}\}$. If for some n , $\mathfrak{a}_n = (1)$, the step stops, otherwise by transfinite induction.

4.18. (i) $\Rightarrow$ (L1) By [Proposition 4.9](#), $S_{\mathfrak{p}}(\mathfrak{a}) = \bigcap \mathfrak{q}_i$. Pick $x \in S_{\mathfrak{p}}$, then $\mathfrak{q}_i = (\mathfrak{q}_i : x)$ by [Proposition 4.4](#). (i) $\Rightarrow$ (L2) Use [Exercise 4.12](#).

4.19. Similar to the proof of [Exercise 4.11](#).

The goal is to prove that there exists $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ irredundant, so we need to prove $\mathfrak{q}_i \not\supseteq \bigcap_{j \neq i} \mathfrak{q}_j$. Using the fact that $\mathfrak{u}\mathfrak{v} \subseteq \mathfrak{q}$ but $\mathfrak{u} \not\subseteq \mathfrak{q} \Rightarrow \mathfrak{v} \subseteq \mathfrak{p}$, we can assume $\mathfrak{b} = \bigcap_{i=1}^{n-1} \mathfrak{q}_i$ irredundant and $\mathfrak{p}_n$ maximal among $\{\mathfrak{p}_i\}$, so we only need to find $\mathfrak{q}_n \not\supseteq \mathfrak{b}$. If such $\mathfrak{q}_n$ doesn't exist, $\mathfrak{b} \subseteq S_{\mathfrak{p}_n}(0)$. Pick a minimal prime ideal $\mathfrak{p} \subseteq \mathfrak{p}_n$ and use [Exercise 4.10](#) and [Proposition 1.11\(ii\)](#).

4.20. (vi) If P is a prime submodule, $\text{rad}_M((P : M)^n M) = (P : M)$ for all $n > 0$.

4.21. We only show that the definition is equivalent to the [element-wise definition](#).

$\Rightarrow$: Let $xm \in Q$ but $m \notin Q$. The map $\phi_x : M/Q \rightarrow M/Q$ defined by $\overline{m} \mapsto x\overline{m}$ is not injective. Therefore, $(\phi_x)^n = 0$ for some n , and thus $x^n(M/Q) = 0$.

$\Leftarrow$: If x is a zero-divisor of M/Q , $xm \in Q$ for some $m \notin Q$. Therefore $x^n M \subseteq Q$ for some n .

Note that when asked for generalization of $(Q : A) = B$, one of the A and B has to be an ideal.

4.22. We only need to replace $\mathfrak{a}$ to N and pick $x \in M$ in the original proof.

4.23. Replace all *primary ideal* to *primary submodule*, but remain *prime ideal* unchanged.

5 Integral Dependence and Valuations

5.1. $f^*(V(I)) = \{f^{-1}(\mathfrak{q}) : \mathfrak{q} \supseteq I\}$. Use [Theorem 5.10](#) to prove $f^*(V(I)) \supseteq V(f^{-1}(I))$.

5.2. $\mathfrak{p} = \ker f$ is prime. By [Theorem 5.10](#), there is a prime ideal $\mathfrak{q} \subseteq B$. $\text{Frac}(B/\mathfrak{q})$ is an algebraic extension of $\text{Frac}(A/\mathfrak{p})$, so $\text{Frac}(A/\mathfrak{p}) \rightarrow \Omega$ can be extended to $\text{Frac}(B/\mathfrak{q}) \rightarrow \Omega$.

5.3. Use the operator $- \otimes_A c^n$.

5.4. If $\left(\frac{1}{x+1}\right)^n + \sum \frac{f_i(x^2-1)}{g_i(x^2-1)(x+1)^i} = 0$, multiply both side by $(x+1)^n$ and take $x = -1$ to get contradiction.

5.5. (ii) Use [Theorem 5.10](#) to prove that the contraction of maximal ideals in B are maximal in A .

5.6. Let $b_i \in B_i$, $f_i \in A[x_i]$ monic, $f_i(b_i) = 0$. Let $f = (f_1, \dots, f_n)$, then $f(b_1, \dots, b_n) = 0$.

5.7. Let $x \in B$ and x integral over A . Let $x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$ is the equation of the least degree. Analyze $x(x^{n-1} + a_{n-2}x^{n-2} + \dots + a_1) = -a_0$.

5.8. Prove that there is a ring $D \supseteq B$ such that f and g factor into linear factors in D . Use Kronecker's construction.

5.9. Let $f^n + g_{n-1}f^{n-1} + \dots + g_0 = 0$. Let $f_1 = f - x^r$ with r larger than all $\deg g_i$, then $f_1(f_1^{n-1} + h_{n-1}f_1^{n-2} + \dots + h_1) = -h_0$ with h_0 monic. Then use [Exercise 5.8](#).

5.10. (a) $\Rightarrow$ (b) Let $f^*(V(\mathfrak{q}_1)) = V(I)$, so $I \subseteq \mathfrak{p}_1 \subseteq \mathfrak{p}_2$, and we have $\mathfrak{p}_2 \in V(I) = f^*(V(\mathfrak{q}_1))$.

(a') $\Rightarrow$ (c') By [Exercise 3.23\(v\)](#) and [Exercise 3.26](#), $f^*(\text{Spec}(B_{\mathfrak{q}})) = f^*(\varinjlim \text{Spec}(B_t)) = \bigcap_t f_t^*(\text{Spec}(B_t))$. Each $f_t^*(\text{Spec}(B_t))$ is an open neighborhood of $\mathfrak{p}$, so it contains $A_{\mathfrak{p}}$.

5.11. Use [Exercise 3.18](#) and [Exercise 5.10](#).

5.12. Note that G has identity element so x is a root of the polynomial $\prod_{\sigma \in G} (t - \sigma(x))$. The coefficients are symmetric polynomials of σ_i . **G acting on $S^{-1}A$ is defined to be $\sigma(x/s) = \sigma(x)/\sigma(s)$.**

5.13. Let $\mathfrak{q} \in P$. Then $\sigma(\mathfrak{q}) \cap A^G = \mathfrak{q} \cap A^G = \mathfrak{p}$, so $\sigma(\mathfrak{q}) \in P$. Let $\mathfrak{q}_1, \mathfrak{q}_2 \in P$ and $x \in \mathfrak{q}_1$. Then $\prod_{\sigma \in G} \sigma(x) \in \mathfrak{q}_1 \cap A^G = \mathfrak{p} \subseteq \mathfrak{q}_2$, so $x \in \sigma^{-1}(\mathfrak{q}_2)$, and so $\mathfrak{q}_1 \subseteq \bigcup_{\sigma \in G} \sigma(\mathfrak{q}_2)$. Use [Proposition 1.11\(i\)](#) and [Corollary 5.9](#).

5.14. Note that $\sigma \in G$ fixes A .

5.15. If L/K is separable, we can find a finite Galois extension M/K such that $L \subseteq M$. Let C be the integral closure of A in M . By [Exercise 5.14](#) and [Exercise 5.13](#), P_C is finite. To prove $P_B = \{\mathfrak{p} \cap B : \mathfrak{p} \in P_C\}$, we need [Theorem 5.10](#), so $\mathfrak{p}_B \in P_B$ can be lifted to $\mathfrak{p}_C$. Verify $\mathfrak{p}_C \in P_C$.

If L/K purely inseparable, fix a $\mathfrak{p} \subseteq A$ and pick $\mathfrak{q} \subseteq B$ such that $\mathfrak{q} \cap A = \mathfrak{p}$. Let $x \in \mathfrak{q}$, so $x^{p^m} \in K$. x is integral over A , so is x^{p^m} . But A integrally closed, so $x^{p^m} \in A$. Therefore, $x^{p^m} \in \mathfrak{q} \cap A = \mathfrak{p}$. If there is another $\mathfrak{q}'$ whose contraction is $\mathfrak{p}$, then $x^{p^m} \in \mathfrak{q}'$ and so $x \in \mathfrak{q}'$. Thus, such picked $\mathfrak{q}$ is unique.

5.16. We have a homomorphism $f : k[y_1, \dots, y_r] \hookrightarrow k[x_1, \dots, x_n] \twoheadrightarrow A$. Let $a = (a_1, \dots, a_r) \in L$ corresponds to a maximal ideal $\mathfrak{m}_a = (y_1 - a_1, \dots, y_r - a_r)$. By [Theorem 5.10](#), we can find a prime ideal $\mathfrak{p}_a \subset A$ such that $f^{-1}(\mathfrak{p}_a) = \mathfrak{m}_a$, which is contained in a maximal $\mathfrak{n}_a$. Notice that $\mathfrak{n}_a$ correspond a point in X .

5.17. Consider $k[t_1, \dots, t_n]/I(X)$. Use [Exercise 5.16](#).

5.18 (**Zariski's lemma**). Let n be the number of generators. If $n = 1$, x_1 is invertible, so $x_1^{-1} = \sum_{i=1}^n c_i x_1^i$.

If $n > 1$, let $A = k[x_1]$ and $K_A = \text{Frac}(A)$. Then $K_A[x_2, \dots, x_n]$ is a finite algebraic extension of K_A , so there are monic polynomials $f_i \in K_A[y]$ such that $f_i(x_i) = 0$ for $2 \leq i \leq n$. Pick $a \in A$ such that $a f_2 \cdots f_n \in A[y]$, then $x_2, \dots, x_n$ are integral over A_a . Therefore, $K_A \subseteq B$ are integral over A_a .

If x_1 transcendental over k , A is a UFD. Since every UFD is integrally closed, A is integrally closed, so A_a is integrally closed by [Proposition 5.12](#). K_A integral over A_a , so $A_a = K_A$, but such a must be a unit in A .

5.19. By [Exercise 5.18](#), $k[t_1, \dots, t_n]/\mathfrak{m}$ is a finite algebraic extension of k , so $k[t_1, \dots, t_n]/\mathfrak{m} \cong k$. Let $t_i \mapsto a_i$, then $t_i - a_i \in \mathfrak{m}$. Prove that $(t_1 - a_1, \dots, t_n - a_n)$ is maximal.

5.20. Let $S = A - \{0\}$. By [Noether normalization lemma](#), $S^{-1}B$ is integral over $S^{-1}A[y_1, \dots, y_r]$ with $y_1, \dots, y_r \in S^{-1}B$. The denominators of $y_1, \dots, y_r$ can be absorbed to $S^{-1}A$, so we can assume $y_1, \dots, y_r \in B$. Let $B = A[x_1, \dots, x_n]$, then $x_i/1 \in S^{-1}B$ is integral over $S^{-1}A[y_1, \dots, y_r]$. Reduce the coefficients of all integral dependence equations to a common denominator s , we have $x_i/1$ is integral over $A_s[y_1, \dots, y_r]$. Therefore, $B_s = A_s[x_1/1, \dots, x_n/1]$ is integral over $A_s[y_1, \dots, y_r]$.

5.21. We have

$$\begin{array}{ccccccc} A & \xrightarrow{\phi_A} & A_s & \longrightarrow & A_s[y_1, \dots, y_n] & \longrightarrow & B_s \xleftarrow{\phi_B} B \\ & & \searrow f & & \searrow f_s & & \searrow f'_s \\ & & & & & & \downarrow g \\ & & & & & & \Omega \end{array}$$

By [Exercise 5.20](#), B_s integral over $A_s[y_1, \dots, y_n]$, so g is obtained from [Exercise 5.2](#). Then, f_s by universal property of localization; f'_s by defining $y_i \mapsto 0$.

5.22. Let $v \neq 0$ an element of B . Then B_v is integral over A , so $f : A \rightarrow \Omega$ can be extended to $\bar{f} : B_v \rightarrow \Omega$ by [Exercise 5.21](#), with $s \neq 0$ an element of A such that $f(s) \neq 0$. Find an $\mathfrak{m}$ such that $s \notin \mathfrak{m}$ and use [Exercise 5.21](#) again, $g : A \rightarrow \overline{A/\mathfrak{m}}$ can be extended to $\bar{g} : B_v \rightarrow \overline{A/\mathfrak{m}}$. Since $\bar{g}(s) \neq 0$, $\bar{g}$ is not trivial, so $\bar{g}(1/v)\bar{g}(v) = \bar{g}(1) = 1$ means $\bar{g}(v) \neq 0$.

Since $\overline{A/\mathfrak{m}}$ integral over $A/\mathfrak{m}$, $\bar{g}(B_v)$ integral over $A/\mathfrak{m}$, so $\bar{g}(B_v) \cong B_v/\ker(\bar{g})$ is a field and therefore $\ker \bar{g}$ is maximal. By [Theorem 5.10](#), $\ker \bar{g} \cap B$ is maximal in B that doesn't contain v .

5.23. (i) $\Rightarrow$ (ii) Note that $f : A \rightarrow A/\ker f$. Use [Proposition 1.1](#).

(ii) $\Rightarrow$ (iii) Consider $A/\mathfrak{p}$.

(iii) $\Rightarrow$ (i) If $\mathfrak{p}$ is not an intersection of maximal ideals, the Jacobson radical $\mathfrak{J}$ of $A/\mathfrak{p}$ is not zero. Pick $f \in \mathfrak{J}$, then $(A/\mathfrak{p})_f \neq 0$. The contraction of a maximal ideal of $(A/\mathfrak{p})_f$ is $\mathfrak{q} \in A$ which is not maximal and doesn't contain f , but all prime ideals strictly containing $\mathfrak{q}$ contains f .

5.24. (i) Use [Exercise 5.23\(iii\)](#) and [going-up theorem](#).

(ii) Consider $B/\mathfrak{q}$ and $A/\mathfrak{q}^c$. Then use [Exercise 5.22](#), [Exercise 5.23\(i\)](#) and [Proposition 1.1](#).

5.25. (i) $\Rightarrow$ (ii) Similar to the proof of [Exercise 5.22](#), we can find $\bar{g} : B \rightarrow \overline{A/\mathfrak{m}}$. Let $B = A[x_1, \dots, x_n]$, then $\bar{g}(x_i)$ integral over $A/\mathfrak{m}$, and therefore $\bar{g}(B)$ is finitely generated as an $A/\mathfrak{m}$ -algebra. By [Zariski's lemma](#), $\bar{g}(B)$ is finite over $A/\mathfrak{m}$. Note that B is a field, so $\bar{g}$ injective.

(ii) $\Rightarrow$ (i) Take $B = A/\mathfrak{p}$. For any $f \neq 0$ in B , B_f is a finitely generated A -algebra. If B_f is a field, it is finite over A , so integral over A . The integral dependence equation also asserts that B_f is integral over B , so B is a field by [Proposition 5.7](#). If B_f is not a field, it has a non-zero prime ideal whose contraction in B doesn't contain f . Therefore, the intersection of all prime ideals in B is zero.

5.26. (2) $\Rightarrow$ (3) $U_1 \cap X_0 = U_2 \cap X_0 \Rightarrow U_1^c \cap X_0 = U_2^c \cap X_0$, then take closure.

(i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) Consider $V(\mathfrak{a}) \cap X_f \cap \text{Max}(A)$. If all maximal ideals in $V(\mathfrak{a})$ contains f , $V(\mathfrak{a}) \cap X_f = \emptyset$ by [Exercise 5.23\(iii\)](#).

(iii) $\Rightarrow$ (i) If not Jacobson, there is a prime ideal $\mathfrak{p}$ such that $\mathfrak{p} \subsetneq \bigcap_{\mathfrak{q} \supseteq \mathfrak{p}} \mathfrak{q}$. Let $f \in \bigcap_{\mathfrak{q} \supseteq \mathfrak{p}} \mathfrak{q} \setminus \mathfrak{p}$, then $X_f \cap V(\mathfrak{p}) = \{\mathfrak{p}\}$.

5.27. $\Rightarrow$: Use [Theorem 5.21](#) to prove maximal elements are valuation rings.

$\Leftarrow$: If $(A, \mathfrak{m}) \subsetneq (A', \mathfrak{m}')$, pick $x \in A' \setminus A$. Then $x^{-1} \in \mathfrak{m}$ because it is not a unit in A .

5.28. (1) $\Rightarrow$ (2) Pick $a \in \mathfrak{a} \setminus \mathfrak{b}$, and any $b \in \mathfrak{b}$. Consider a/b and b/a .

(2) $\Rightarrow$ (1) Consider $(a) \subseteq (b)$ or $(b) \subseteq (a)$, we have $a = xb$ or $b = ya$, so $y = x^{-1}$.

5.29. *B is a local ring of A means $B = A_{\mathfrak{p}}$ for some prime ideal $\mathfrak{p}$.* Let $\mathfrak{m}$ be the maximal ideal of B and let $\mathfrak{p} = \mathfrak{m} \cap A$. If $s \in A$ but $s \notin \mathfrak{p}$, then $s \in B$ and $s \notin \mathfrak{m}$, so s is a unit in B . Let $A_{\mathfrak{p}} \rightarrow B$ to be $x/s \mapsto s^{-1}x$, which is injective; $\mathfrak{p}A_{\mathfrak{p}} \subseteq B$, so B dominates $A_{\mathfrak{p}}$. By [Exercise 5.28](#), $A_{\mathfrak{p}}$ is a valuation ring, then use [Exercise 5.27](#).

5.30. $xy^{-1} \in A$ or $x^{-1}y \in A$, so $\geq$ is a total order. Also $(xz)(yz)^{-1} = xy^{-1}$.

Treat v as a representation map. If $v(x) \geq v(y)$, $(x+y)y^{-1} = xy^{-1} + 1 \in A$.

5.31. Use the given conditions to verify the set is closed under addition and multiplication. Then verify it is a valuation ring by taking $x = y = 1$ and $y = x^{-1}$ in the first condition.

5.32. $v(x) + v(y) = v(xy)$ implies $v(A - \mathfrak{p})$ is closed under addition. Let $\Delta = v(A - \mathfrak{p}) \cup (-v(A - \mathfrak{p}))$. If $0 \leq v(x) \leq v(y)$ with $y \in A - \mathfrak{p}$, then $x^{-1}y \in A$ and $x(x^{-1}y) \in A - \mathfrak{p}$. Since $A - \mathfrak{p}$ is saturated as a multiplicatively closed set, $x \in A - \mathfrak{p}$.

Let $S = \{x \in A : v(x) \in \Delta, v(x) \geq 0\}$. If $x, y \notin S$, $v(x+y) \geq \min(v(x), v(y))$ means $x+y \notin S$; if $x \notin S$ and $y \in A$, $v(xy) \geq v(x)$ means $xy \notin S$. Hence, $A - S$ is an ideal, and it is automatically prime due to S is multiplicatively closed.

Define $v_{A_{\mathfrak{p}}} : K^* \rightarrow \Gamma/\Delta$ to be $x \mapsto x + \Delta$, then $A_{\mathfrak{p}} = \{x \in K : v_{A_{\mathfrak{p}}}(x) \geq 0\}$.

The fraction field of $A/\mathfrak{p}$ is not K but $(A/\mathfrak{p})_{\bar{\mathfrak{p}}}$. Let $x = \bar{a}/\bar{b} \in (A/\mathfrak{p})_{\bar{\mathfrak{p}}}$ and define $v_{A/\mathfrak{p}}(x) = v(a) - v(b)$, so $v_{A/\mathfrak{p}}(x) \in \Delta$. We have $v_{A/\mathfrak{p}}(A/\mathfrak{p}) = \Delta_+$.

5.33. Let $v(a/b) = v_0(a) - v_0(b)$. Use $v(xy) = v(x) + v(y)$ to prove it is well-defined and unique.

5.34. Since $gf(A) = A$, $g(B) \supseteq A$. Let $\mathfrak{n}$ be a maximal ideal of $g(B)$, by [Exercise 5.10](#), $\mathfrak{m} = \mathfrak{n} \cap A$ is the maximal ideal of A . Also since $(g(B))_{\mathfrak{n}}$ dominates A , by [Exercise 5.27](#), $(g(B))_{\mathfrak{n}} = A$ and therefore $g(B) \subseteq A$.

5.35. Let A' be an arbitrary valuation ring of $\text{Frac}(B)$ containing $f(A)$. By hypothesis, $\text{Spec}(B \otimes_A A') \rightarrow \text{Spec}(A')$ is a closed map, and by [Exercise 5.34](#), $g : B \otimes_A A' \rightarrow \text{Frac}(B)$ defined by $b \otimes a' \mapsto ba'$ satisfies $g(B \otimes_A A') = A'$. In other words, $ba' \in A'$ for all $b \in B$ and all $a' \in A'$. By taking $a' = 1$, we have $B \subseteq A'$. By [Corollary 5.22](#), B is in the integral closure of $f(A)$, so B is integral over $f(A)$.

The composition map $A \rightarrow B \rightarrow B/\mathfrak{p}_i$ is integral. By [Exercise 5.6](#), $A \rightarrow \prod_{i=1}^n B/\mathfrak{p}_i$ is integral. Using integral dependence equation can prove $A \rightarrow A/\mathfrak{R}$ is integral then $A \rightarrow B$ is integral.

6 Chain Conditions

6.1. (i) Let $x \in \ker(u)$. Since u^n surjective, there is a $y \in M$ such that $u^n(y) = x$. But $u^{n+1}(y) = u(x) = 0$, means that $y \in \ker(u^{n+1}) = \ker(u^n)$.

(ii) Let $x \in M$. Then $u^n(x) \in \text{im}(u^n) = \text{im}(u^{n+1})$, so there is a $y \in M$ such that $u^{n+1}(y) = u^n(x)$. But u^n injective, means that $u(y) = x$.

6.2. If M is not Noetherian, there is a submodule N which is not finitely generated. Construct a chain of finitely generated submodules by the generators of N .

6.3. Consider $0 \rightarrow N_2/(N_1 \cap N_2) \rightarrow M/(N_1 \cap N_2) \rightarrow M/N_2 \rightarrow 0$. Use [Proposition 2.1](#).

6.4. Let $m_1, \dots, m_n$ be generators of M , then $\mathfrak{a} = \bigcap \text{Ann}(m_i)$. $A/\text{Ann}(m_i) \cong Am_i \subseteq M$, then use [Exercise 6.3](#).

6.5. Use the open set definition.

6.6. (i) $\Rightarrow$ (iii) Use [Exercise 6.5](#).

(ii) $\Rightarrow$ (i) Consider a chain of ascending open sets in X .

6.7. Let Σ be the collection of closed subsets of X which are not finite union of irreducible closed subspaces. Then Σ has minimal element X_0 . Prove that X_0 cannot be either irreducible or reducible.

6.8. The converse isn't true. The counter example is $k[x_1, \dots]/(x_1^2, \dots)$.

6.9. The irreducible components of $\text{Spec}(A)$ is $V(\mathfrak{p})$ where $\mathfrak{p}$ minimal ([Exercise 1.20\(iv\)](#)).

6.10. Use [Exercise 3.19\(v\)](#).

6.11. By [Exercise 6.7](#), $V(\mathfrak{b})$ is a finite union of $V(\mathfrak{q}_i)$, where $\mathfrak{q}_i$ minimal and containing $\mathfrak{b}$. Then use [Exercise 5.10\(c\)](#).

6.12. The converse isn't true. A counter example is an infinite product $\prod \mathbb{Z}/2\mathbb{Z}$.

7 Noetherian Rings

7.1 (Cohen's theorem). Let $\mathfrak{a}$ be a maximal element in Σ . If $xy \in \mathfrak{a}$ but $x, y \notin \mathfrak{a}$, then $\mathfrak{a} + (x)$ is finitely generated. Let $\{x, a_1, \dots, a_n\}$ be its generators and let $\mathfrak{a}_0 = (a_1, \dots, a_n)$. Prove that $\mathfrak{a}_0 \not\subseteq \mathfrak{a}$ is impossible. Therefore, if $u \in \mathfrak{a} \subseteq \mathfrak{a} + (x) = \mathfrak{a}_0 + (x)$, $u = a_0 + bx$ with $b \in \mathfrak{a}$, so $\mathfrak{a} \subseteq \mathfrak{a}_0 + x \cdot (\mathfrak{a} : x)$; we also have $x \cdot (\mathfrak{a} : x) \subseteq \mathfrak{a}$. Note that $y \in (\mathfrak{a} : x)$.

7.2. $\Rightarrow$: [Exercise 1.5\(ii\)](#).

$\Leftarrow$: $(a_0, a_1, \dots) \subseteq \mathfrak{R}$ and use [Corollary 7.15](#).

7.3. (i) $\Rightarrow$ (ii) Use [Proposition 4.8](#) and [Proposition 4.4](#).

(ii) $\Rightarrow$ (iii) Let $S = \{1, x, x^2, \dots\}$. Prove that $\bigcup (\mathfrak{a} : x^n) = S(\mathfrak{a})$.

(iii) $\Rightarrow$ (i) Similar to the proof of [Proposition 7.12](#).

7.4. (i) $\mathfrak{p} = \{(z - c) : |c| = 1\}$ is prime, and the ring is $(\mathbb{C}[x])_{\mathfrak{p}}$. Use [Proposition 7.3](#).

(ii) Use [Exercise 1.5\(i\)](#). If $f = z^n g$ with g invertible, $(f) = (z^n)$, so the only ideals are (0) and (z^n) .

(iii) $\dots \subsetneq (\sin \pi n z) \subsetneq (\sin \pi(n+1)z) \subsetneq \dots$ is an ascending chain.

- (iv) The ring is $\mathbb{C}[z^{k+1}]$.
 (v) The ring is $\mathbb{C}[z, z^2, \dots, wz, wz^2, \dots, w^2z, w^2z^2, \dots]$.

7.5. Use [Exercise 5.12](#) and [Proposition 7.8](#).

7.6. A ring is finitely generated if it is finitely generated as a $\mathbb{Z}$ -algebra. $\mathbb{Z}$ is a Jacobson ring. By [Exercise 5.25](#), K is finite over $\mathbb{Z}$, so it is of the form $\mathbb{Z}^n \oplus \bigoplus_i \mathbb{Z}/p_i\mathbb{Z}$ due to the addition operation forms a finitely generated abelian group. If $\text{char } K = 0$, $\mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{Z}^n$, then use [Proposition 7.8](#).

7.7. Use ascending chain condition.

7.8. $A \cong A[x]/(x)$.

7.9. Let $\mathfrak{a}_0 \subseteq \mathfrak{a}_1 \subseteq \dots$ be a chain in A . Since $\mathfrak{a}_0$ is contained in only finite many maximal ideals, there is a large enough n such that $S_{\mathfrak{m}}^{-1}\mathfrak{a}_n = S_{\mathfrak{m}}^{-1}\mathfrak{a}_{n+1} = \dots$ for all $\mathfrak{m} \supseteq \mathfrak{a}_0$. We also have $S_{\mathfrak{m}}^{-1}\mathfrak{a}_i = A_{\mathfrak{m}}$ for all i , when $\mathfrak{m} \not\supseteq \mathfrak{a}_0$. Then apply [Proposition 3.8](#) on $S_{\mathfrak{m}}^{-1}(\mathfrak{a}_{n+i}/\mathfrak{a}_n)$.

7.10. $M[x] = M \otimes A[x]$, so it is a homomorphic image of $M \times A[x]$. Use [Proposition 7.1](#).

7.11. Counter example is an infinite product $\prod \mathbb{Z}/2\mathbb{Z}$.

7.12. By [Exercise 3.16\(i\)](#), $\mathfrak{a} \mapsto \mathfrak{a}^e$ is injective. Then use ascending chain condition.

7.13. Transfer the Noetherian property $B \rightarrow B_{\mathfrak{p}} \rightarrow B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ and use [Exercise 6.8](#).

7.14 ([Hilbert's Nullstellensatz](#)). If $f \notin \sqrt{\mathfrak{a}}$, there is a $\mathfrak{p} \supseteq \mathfrak{a}$ such that $f \notin \mathfrak{p}$. $(A/\mathfrak{p})_{\overline{f}}$ is a finitely generated k -algebra, so by [Zariski's lemma](#), $\left((A/\mathfrak{p})_{\overline{f}}\right)/\mathfrak{m} = k$. Let x_i be the images of t_i in $\left((A/\mathfrak{p})_{\overline{f}}\right)/\mathfrak{m}$, then $x = (x_1, \dots, x_n)$ satisfies $x \in V$ but $f(x) \neq 0$.

Rabinowitsch trick: Let $f_1, \dots, f_m$ be generators of $\mathfrak{a}$. If $f \in I(V)$, then $f_1, \dots, f_m, 1 - t_0f$ has no zeros in $k[t_0, t_1, \dots, t_n]$. Apply [Exercise 5.17](#), $1 = g_0(1 - t_0f) + \sum_{i=1}^m g_i f_i$. Now put $t_0 = 1/f$, we have $g_i \in k[t_1, \dots, t_n, 1/f]$. Therefore, $g_i = h_i/f^{N_i}$ with $h_i \in k[t_1, \dots, t_n]$. Hence we have $f^N = \sum_{i=1}^m h_i f^{N-N_i} f_i$ for all $N \geq \max_i \{N_i\}$.

7.15. (iv) $\Rightarrow$ (i) Let $x_1, \dots, x_n$ be elements of M whose images in $M/\mathfrak{m}M$ form a basis of k -vector space. Let $N = (x_1, \dots, x_n)$, so $N + \mathfrak{m}M = M$. Since $\mathfrak{J} = \mathfrak{m}$, apply [Nakayama's lemma](#) on M/N , we have $M = N$. Let F be a free A -module with basis $e_1, \dots, e_n$, and $\phi : F \rightarrow M$ be $\phi(e_i) = x_i$. Then

$$\text{Tor}_1^A(k, M) \rightarrow k \otimes_A \ker \phi \rightarrow k \otimes_A F \xrightarrow{1 \otimes \phi} k \otimes_A M \rightarrow 0$$

exact. Since $k \otimes F$ and $k \otimes M = M/\mathfrak{m}M$ are vector spaces of a same dimension, $1 \otimes \phi$ is an isomorphism, so $k \otimes \ker \phi = 0$. By [Exercise 2.3](#), $\ker \phi = 0$.

7.16. Use [Proposition 3.10](#) and [Exercise 7.15](#).

7.17. Only prove the generalization of [Proposition 7.12](#). Let $ax \in Q$ but $x \notin Q$, we have $(Q : a^n) = (Q : a^{n+1})$ for some n . If $u \in a^n M \cap Ax$, $u = a^n m = bx$, so $a^{n+1}m = bax \in Q$, and so $m \in (Q : a^{n+1}) = (Q : a^n)$, and so $u = a^n m \in Q$. Since Q irreducible, $a^n M = Q$.

7.18. (i) $\Leftrightarrow$ (ii) The generalization of [Proposition 7.17](#).

(ii) $\Leftrightarrow$ (iii) $\text{Ann}(x) = \ker(A \rightarrow Ax)$, and $A/\mathfrak{p} = A/\text{Ann}(x) \cong Ax$.

7.19. Let $\mathfrak{a}_{kl} = \left(\bigcap_{i \neq k}^r \mathfrak{b}_i\right) \cap \mathfrak{c}_l$, then $\bigcap_{l=1}^s \mathfrak{a}_{kl} = \mathfrak{a}$ for all k . Since $\mathfrak{a} \subseteq \mathfrak{b}_k$, the projection $\pi : A \rightarrow A/\mathfrak{b}_k$ maps the equation to $\bigcap_{l=1}^s \pi(\mathfrak{a}_{kl}) = 0$. Since $\mathfrak{b}_k$ irreducible, $\pi(\mathfrak{a}_{kl}) = 0$ for some l and therefore $\mathfrak{a}_{kl} \subseteq \mathfrak{b}_k$. The construction of $\mathfrak{a}_{kl}$ implies $\mathfrak{a} \subseteq \mathfrak{a}_{kl} \subseteq \mathfrak{b}_i$ for all $i \neq k$. Take the intersection of all i , we conclude that for any k , there is an l such that $\mathfrak{a}_{kl} = \mathfrak{a}$. Therefore, $r = s$, and use [Proposition 4.5](#).

7.20. (i) $\Rightarrow$: We have a chain $\mathcal{F}_0 \subseteq \mathcal{C}_1 \subseteq \mathcal{F}_1 \subseteq \mathcal{C}_2 \subseteq \dots \subseteq \mathcal{F}$, where $\mathcal{F}_0$ is the topology, $\mathcal{C}_i = \mathcal{F}_{i-1} \cup \{S^c : S \in \mathcal{F}_{i-1}\}$, and $\mathcal{F}_i$ is a collection of all finite intersections of sets in $\mathcal{C}_i$.

(ii) $\Rightarrow$: Use the open set definition.

7.21. $\Rightarrow$: Use [Exercise 7.20](#).

$\Leftarrow$: If $E \notin \mathcal{F}$, let $\mathcal{C} = \{X' \subseteq X : X' \text{ closed, } E \cap X' \notin \mathcal{F}\}$. By Noetherian property, $\mathcal{C}$ has a minimal element X_0 . Use the closed set definition to show X_0 is irreducible. If $\overline{E \cap X_0} \neq X_0$, $E \cap \overline{E \cap X_0} \in \mathcal{F}$ by minimality, but $E \cap \overline{E \cap X_0} = E \cap X_0$. If $U_0 \subseteq E \cap X_0$, let $U_0 = U \cap X_0$ where U is open in X , then $E \cap X_0 = U_0 \cup (E \cap U^c \cap X_0)$.

7.22. $\Leftarrow$: If E is not open, let $\mathcal{C} = \{X' \subseteq X : X' \text{ closed, } E \cap X' \text{ is not open}\}$ and use the notations similar to [Exercise 7.21](#). If $U_0 \subseteq E \cap X_0$, $U^c \cap X_0$ is closed, so by minimality $E \cap U^c \cap X_0$ is open.

7.23. It is enough to prove $f^*(U \cap C)$ is constructible. Let $C = V(\mathfrak{a})$, by $\text{Spec}(B/\mathfrak{a}) \hookrightarrow \text{Spec}(B) \xrightarrow{f^*} \text{Spec}(A)$, we only need to show $f^*(U)$ is constructible. By [Exercise 6.6](#) and [Exercise 1.17\(vii\)](#), $U = \bigcup_{i=1}^n X_{g_i}$. By $\text{Spec}(B_{g_i}) \hookrightarrow \text{Spec}(B) \xrightarrow{f^*} \text{Spec}(A)$, it suffices to show $f^*(Y)$ is constructible and take $B = B_{g_i}$. By [Exercise 6.7](#), we can assume B is an integral domain.

Let X_0 irreducible and suppose $f^*(Y) \cap X_0$ is dense in X_0 . By [Exercise 1.20\(iv\)](#), $X_0 = V(\mathfrak{p})$, so $f^{*-1}(X_0) = V(f(\mathfrak{p})B) = \text{Spec}(A/\mathfrak{p} \otimes_A B)$ by [Exercise 1.21\(ii\)](#) and [Exercise 2.2](#). For any X_0 , we also have $f^*(Y) \cap X_0 = f^*(f^{*-1}(X_0))$. Using the universal property of quotient rings to restrict f on $A/\mathfrak{p}$, denoted as $\tilde{f}$, so $\tilde{f}^*(Y) = f^*(\text{Spec}(A/\mathfrak{p} \otimes_A B)) = f^*(Y) \cap X_0$ is dense in X_0 . By [Exercise 1.21\(v\)](#), $\ker \tilde{f} \subseteq \mathfrak{R}_{A/\mathfrak{p}} = 0$, so $\tilde{f}$ is injective. Replacing A with $A/\mathfrak{p}$, we may assume A is an integral domain.

Suppose $A \subseteq B$ are integral domains. By [Exercise 5.21](#), there exists $s \neq 0$ such that each $g_{\mathfrak{p}} : A \rightarrow \text{Frac}(A/\mathfrak{p}) \rightarrow \text{Frac}(A/\mathfrak{p})$ can be extended to $\tilde{g}_{\mathfrak{p}} : B \rightarrow \text{Frac}(A/\mathfrak{p})$. If $s \notin \mathfrak{p}$, $\mathfrak{q} = \ker(\tilde{g}_{\mathfrak{p}})$ satisfies $\mathfrak{q} \cap A = \mathfrak{p}$, so $X_s \subseteq \tilde{f}^*(Y)$. By [Exercise 7.21](#), $\tilde{f}^*(Y)$ is constructible.

7.24. Let U be an open set in Y . By [Exercise 7.23](#), $f^*(U)$ is a constructible subset of X . Since f has the going-down property, using $\mathfrak{q} \in U \Rightarrow \text{Spec}(B_{\mathfrak{q}}) \subseteq U$, we have $\mathfrak{p}' \in f^*(U)$ for each $\mathfrak{p}' \subseteq \mathfrak{p} \in f^*(U)$.

Let X_0 be an irreducible closed subset of X and suppose $f^*(U) \cap X_0 \neq \emptyset$. By [Exercise 1.20\(iv\)](#), $X_0 = V(\mathfrak{p}_0)$, so $\mathfrak{p}_0 \in f^*(U)$, and therefore $f^*(U) \cap X_0$ is dense in X_0 . By [Exercise 7.20\(ii\)](#), $f^*(U) \cap X_0$ contains a non-empty open subset, and by [Exercise 7.22](#), $f^*(U)$ is open.

7.25. Use [Exercise 5.11](#) and [Exercise 7.24](#).

7.26. (i) $\lambda(D) = 0$. Use the universal property of quotient groups.

(ii) Use [Exercise 7.18](#) and $0 \rightarrow M_{i-1} \rightarrow M_i \rightarrow A/\mathfrak{p}_i \rightarrow 0$.

(iii) $0 \rightarrow A \xrightarrow{a} A \rightarrow A/(a) \rightarrow 0$ means $\gamma(A/(a)) = 0$ for all $a \neq 0$, so $K(A)$ is generated by a single element $\gamma(A)$. We also have $0 \rightarrow A \rightarrow A^n \rightarrow A^{n-1} \rightarrow 0$, so $K(A)$ infinite cyclic.

(iv) Any finitely generated B -module is also a finitely generated A -module. The map is constructed by using the universal property in (i) on $F(B) \rightarrow F(A) \rightarrow K(A)$.

7.27. (iii) Use [Exercise 7.15](#), so that $F_1(A)$ is the set of all free modules.

(iv) Use [Proposition 2.15](#) to prove M_B is a flat B -module. The rest is similar to [Exercise 7.26\(iv\)](#).

(v) Treat x, y as modules. Use the formulas in [Exercise 7.26](#) and [Exercise 7.27](#).

8 Artinian Rings

8.1. Use [Corollary 7.16](#) on local ring $A_{\mathfrak{p}_i}$. Then apply the contraction. If $\mathfrak{q}_i$ is isolated, then $\mathfrak{p}_i$ is minimal.

8.2. (i) $\Rightarrow$ (ii) Use [Proposition 8.1](#).

(iii) $\Rightarrow$ (i) Use [Proposition 8.5](#).

8.3. (i) $\Rightarrow$ (ii) By [Proposition 8.7](#), we can assume that A is an Artinian local ring. By [Proposition 8.4](#), $\mathfrak{m}^n = 0$ for some n , so we have a chain $A = \mathfrak{m}^0 \supseteq \mathfrak{m}^1 \supseteq \cdots \supseteq \mathfrak{m}^n = 0$.

By [Zariski's lemma](#), $A/\mathfrak{m}$ is a finitely generated k -module. By [Proposition 8.5](#), A is Noetherian, so $\mathfrak{m}^i$ is a finitely generated A -module, and therefore $\mathfrak{m}^i/\mathfrak{m}^{i+1}$ is a finitely generated $A/\mathfrak{m}$ -module. Use

$$\begin{aligned} \dim_k(\mathfrak{m}^i/\mathfrak{m}^{i+1}) &= \dim_{A/\mathfrak{m}}(\mathfrak{m}^i/\mathfrak{m}^{i+1}) \dim_k(A/\mathfrak{m}); \\ \dim_k(\mathfrak{m}^i) &= \dim_k(\mathfrak{m}^{i+1}) + \dim_k(\mathfrak{m}^i/\mathfrak{m}^{i+1}), \end{aligned} \quad \text{where } 0 \rightarrow \mathfrak{m}^{i+1} \rightarrow \mathfrak{m}^i \rightarrow \mathfrak{m}^i/\mathfrak{m}^{i+1} \rightarrow 0$$

to finish the proof.

(ii) $\Rightarrow$ (i) A is a k -vector space of finite dimension. Use [Proposition 6.10](#).

8.4. (i) $\Rightarrow$ (iii) The finitely-generated property is preserved for localization and quotient of modules.
(ii) $\Rightarrow$ (iii) $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ is a finitely generated $k(\mathfrak{p})$ -algebra. By [Hilbert's basis theorem](#), any finitely generated $k(\mathfrak{p})$ -algebra is Noetherian. Then use [Exercise 8.2](#) and [Exercise 8.3](#).

(iii) $\Rightarrow$ (ii),(iv) $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ is Noetherian and integral over the field $k(\mathfrak{p})$. If $\mathfrak{q} \in B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ is prime, $(B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}})/\mathfrak{q}$ is an integral domain, so it is a field by [Proposition 5.7](#). Hence, every prime ideal in $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ is maximal, which means $\dim(B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}) = 0$. Thus, $B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$ is Artinian by [Proposition 8.5](#). Then use [Proposition 8.3](#).

A counter example is $f : \mathbb{Q} \hookrightarrow \overline{\mathbb{Q}}$ where $\overline{\mathbb{Q}}$ is an algebraic closure of $\mathbb{Q}$.

8.5. Use [Noether normalization lemma](#) and let $\mathfrak{m}$ be a maximal ideal in $k[y_1, \dots, y_r]$. By [Exercise 8.4](#), $A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}$ is finite over $k[y_1, \dots, y_r]/\mathfrak{m} = k$, so by [Exercise 8.3](#) and [Proposition 8.3](#), $A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}$ only has finite many maximal ideals, correspond to finite many points in X .

Let N be the number of generators of A over $k[y_1, \dots, y_r]$. The k -dimension of k -vector space $A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}$ is uniformly bounded by N . If $\mathfrak{n}_1, \dots, \mathfrak{n}_M$ are maximal ideals in $A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}$, by [Proposition 1.10](#), $A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}} \rightarrow \prod_{i=1}^M (A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}})/\mathfrak{n}_i$ is surjective, so $\dim_k(A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}}) \geq \sum_{i=1}^M \dim_k((A_{\mathfrak{m}}/\mathfrak{m}A_{\mathfrak{m}})/\mathfrak{n}_i) \geq M$.

8.6. Consider $(A/\mathfrak{q})_{\overline{\mathfrak{p}}}$, which is an Artinian local ring. Use the following claims then [Proposition 6.7](#).

- If A is an Artinian local ring, $\mathfrak{m}$ its only maximal ideal. Then every ideal in A is $\mathfrak{m}$ -primary. Radical is the intersection of the only prime ideal. Then use [Proposition 4.2](#).
- There is a 1-1 correspondence between $\mathfrak{p}$ -primary ideals in A and $S^{-1}\mathfrak{p}$ -primary ideals in $S^{-1}A$. Use [Proposition 3.11\(i\)](#) and [Proposition 4.9](#).

9 Discrete Valuation Rings and Dedekind Domain

9.1. [Proposition 3.11\(iv\)](#) asserts that $\dim S^{-1}A$ can only be 0 and 1. If $\dim S^{-1}A = 0$, it is $\text{Frac } A$; if $\dim S^{-1}A = 1$, use [Proposition 5.12](#).

By [Corollary 3.15](#), $S^{-1}(M^{-1}) = S^{-1}(A : M) = (S^{-1}A : S^{-1}M) = (S^{-1}M)^{-1}$. We also have $S^{-1}(MN) = (S^{-1}M)(S^{-1}N)$, so the map $M \mapsto S^{-1}M$ is a homomorphism. Use [Proposition 3.11\(i\)](#). Note that $(u) \mapsto S^{-1}(u) = (S^{-1}A)u$ maps P into P' , so $H \rightarrow H'$ is surjective.

9.2. Localize and use [Proposition 3.8](#) to the final move, we can assume that A is a DVR.

[Proposition 9.2\(iii\)\(v\)](#) assert that $c(f) = (x^n)$ where x is the generator of $\mathfrak{m}$, so $f = x^n f_0$ where f_0 is primitive. Apply [Exercise 1.2\(iv\)](#) to f_0, g_0 and note that for any $a \in A$, $ac(f) = c(af)$.

9.3. $\Rightarrow$: $(x_1, x_2) = (x_1)$ if $x_1 x_2^{-1} \in A$, so every ideal is principal. Then use [Proposition 9.2\(iii\)](#).
 $\Leftarrow$: If $v(y) \geq v(x)$, $y = (yx^{-1})x$, so every ideal in DVR is of the form $\mathfrak{m}_k = \{z \in A : v(z) \geq k\}$.

9.4. For any non-zero $x \in A$, $x \in \mathfrak{m}^n$ but $x \notin \mathfrak{m}^{n+1}$ for some n . Define a discrete valuation $v(x) = n$.

9.5. By [Exercise 3.13](#) and [Exercise 7.16](#), and notice that torsion free module over a PID is flat.

9.6. From [Theorem 9.3\(iii\)](#), [Proposition 9.2\(v\)](#), and the structure theorem for finitely generated modules over a PID, $M_{\mathfrak{p}} = \bigoplus_{i=1}^N A_{\mathfrak{p}}/\mathfrak{p}^{n_i} A_{\mathfrak{p}} = \bigoplus_{i=1}^N (A/\mathfrak{p}^{n_i})_{\mathfrak{p}}$. Define a homomorphism $\phi : M \rightarrow \bigoplus_{\mathfrak{p}} M_{\mathfrak{p}}$ by $m \mapsto (m/1, \dots)$, so each $\phi_{\mathfrak{p}}$ is an isomorphism. By [Proposition 3.9](#), ϕ is an isomorphism.

By [Exercise 3.19\(v\)](#), $\text{Supp}(M) = V(\text{Ann}(M))$. From [Proposition 9.1](#) and [Theorem 9.3\(ii\)](#), $\text{Ann}(M) = \mathfrak{p}_1^{n_1} \cdots \mathfrak{p}_k^{n_k}$. Use $\dim A = 1$ to prove that $V(\text{Ann}(M))$ is finite.

9.7. From [Proposition 9.1](#) and [Theorem 9.3\(ii\)](#), $\mathfrak{a} = \mathfrak{p}_1^{n_1} \cdots \mathfrak{p}_k^{n_k}$. By [Proposition 1.10](#), $A \rightarrow \prod_{i=1}^k A/\mathfrak{p}_i^{n_i}$ is a surjective with kernel $\mathfrak{a}$. Note that each component $A/\mathfrak{p}_i^{n_i}$ is local so is PID.

If $a \in \mathfrak{b}$, $\mathfrak{b}/(a) = (\bar{b})$ and therefore $\mathfrak{b} = (a, b)$.

9.8. Localize and note that all ideals in DVR are $\{z \in A : v(z) \geq k\}$. Use [Proposition 3.8](#) to pull back.

9.9. This is equivalent to say that

$$A \xrightarrow{\phi} \bigoplus_{i=1}^n A/\mathfrak{a}_i \xrightarrow{\psi} \bigoplus_{i < j} A/(\mathfrak{a}_i + \mathfrak{a}_j)$$

is exact, where $\phi(x) = (x + \mathfrak{a}_i)_{i=1}^n$, and $\psi(x_i + \mathfrak{a}_i)_{i=1}^n = (x_i - x_j + \mathfrak{a}_i + \mathfrak{a}_j)_{i < j}$. If A is DVR, we have $\mathfrak{a}_i + \mathfrak{a}_j = \mathfrak{a}_{\min(i,j)}$. If $(x_i + \mathfrak{a}_i)_{i=1}^n \in \ker \psi$, $x_i + \mathfrak{a}_i = x_j + \mathfrak{a}_j$ for all $i < j$, so take $j = n$ for all i and we have $\phi(x_n) \in \ker \psi$. Then use [Theorem 9.3\(iii\)](#) and [Proposition 3.8](#) for Dedekind domain.

10 Completion

10.1. Some completions are calculated as follows

- $A/p^n A = \bigoplus_{k \geq 0} \mathbb{Z}/p\mathbb{Z}$, where the transition maps are all identity.
- $A_n = \alpha^{-1}(p^n B) = \bigoplus_{k > n} \mathbb{Z}/p\mathbb{Z}$, so $A/A_n = \bigoplus_{k=0}^n \mathbb{Z}/p\mathbb{Z}$. The transition maps mean, if $(x^{(1)}, x^{(2)}, \dots) \in \varprojlim A/A_n$, then $x_i^{(i)} = x_i^{(i+1)} = \dots$ for all i .
- $B/p^n B = (\bigoplus_{k=0}^n \mathbb{Z}/p^k \mathbb{Z}) \oplus (\bigoplus_{k > n} \mathbb{Z}/p^n \mathbb{Z})$. The transition maps are similar to A/A_n .

The sequence $0 \rightarrow A \xrightarrow{\alpha} B \xrightarrow{p} pB \rightarrow 0$ is exact. Since p -adic topology and induced topology on pB coincide, by [Corollary 10.3](#), $\ker(\hat{p}) = \varprojlim A/A_n$.

10.2. The explicit form of the given exact sequence is

$$0 \rightarrow \bigoplus_{k > n} \mathbb{Z}/p\mathbb{Z} \rightarrow \bigoplus_{k \geq 0} \mathbb{Z}/p\mathbb{Z} \rightarrow \bigoplus_{k=0}^n \mathbb{Z}/p\mathbb{Z} \rightarrow 0.$$

Use [Exercise 10.1](#), and note that $\varprojlim \bigoplus_{k > n} \mathbb{Z}/p\mathbb{Z} = 0$. [The definition of \$\varprojlim^1 A\$ in this book is here.](#)

10.3. Denote $N = \bigcap_{\mathfrak{m} \supseteq \mathfrak{a}} \ker(M \rightarrow M_{\mathfrak{m}})$. Let $x \in \bigcap_{n=1}^{\infty} \mathfrak{a}^n M$. By [Proposition 10.17](#), we have $(1+a)x = 0$. If $\mathfrak{m} \supseteq \mathfrak{a}$, $1+a \in A - \mathfrak{m}$, so $x \in N$. Since $(\ker(M \rightarrow M_{\mathfrak{m}}))_{\mathfrak{m}} = 0$, we have $N_{\mathfrak{m}} = 0$ for all $\mathfrak{m} \supseteq \mathfrak{a}$. By [Exercise 3.14](#), $N = \mathfrak{a}N$, and therefore $N = \mathfrak{a}N = \mathfrak{a}^2 N = \dots$. Thus, $N = \bigcap_{n=1}^{\infty} \mathfrak{a}^n N \subseteq \bigcap_{n=1}^{\infty} \mathfrak{a}^n M$.

We can see N is also equal to $\bigcap_{\mathfrak{p} \supseteq \mathfrak{a}} \ker(M \rightarrow M_{\mathfrak{p}})$. Therefore, $\text{Supp}(M) \cap V(\mathfrak{a}) = 0$ iff $N = M$, iff $M = \bigcap_{n=1}^{\infty} \mathfrak{a}^n M$, iff $M = \mathfrak{a}M$. By [Exercise 2.2](#), [Proposition 10.13](#) and [Proposition 10.15\(iii\)](#), $\hat{M}/\hat{\mathfrak{a}}\hat{M} = \hat{A}/\hat{\mathfrak{a}}\hat{A} \otimes_{\hat{A}} \hat{M} = A/\mathfrak{a} \otimes_A (M \otimes_A \hat{A}) = M/\mathfrak{a}M \otimes_A \hat{A}$. Since $\hat{A} \neq 0$, by [Exercise 2.3](#), $M/\mathfrak{a}M = 0$ iff $\hat{M}/\hat{\mathfrak{a}}\hat{M} = 0$. By [Proposition 10.15\(iv\)](#) and [Nakayama's lemma](#), $\hat{M} = \hat{\mathfrak{a}}\hat{M}$ iff $\hat{M} = 0$.

10.4. x not a zero-divisor, so $A \xrightarrow{x} A$ is injective. Use [Proposition 10.14](#) and apply the functor $\hat{A} \otimes_A -$.

Counter example is $k[x, y]/f$, where $f = y^2 - x^2 - x^3$. The completion is $k[[x, y]]/f$. f is irreducible in $k[x, y]$, but $f = (y - x\sqrt{1+x})(y + x\sqrt{1+x})$ is 0 in $k[[x, y]]/f$.

10.5. We can deduce from [Proposition 10.2](#) that taking $\mathfrak{a}$ -adic completion on $0 \rightarrow b^m M \rightarrow M \rightarrow M/b^m M \rightarrow 0$ preserves exactness. Thus, $(M/b^m M)^{\mathfrak{a}} \cong M^{\mathfrak{a}}/b^m M^{\mathfrak{a}}$, and therefore

$$(M^{\mathfrak{a}})^{\mathfrak{b}} = \varprojlim_m \left(\varprojlim_n (M/\mathfrak{a}^n M)/(\mathfrak{b}^m (M/\mathfrak{a}^n M)) \right) = \varprojlim_m \left(\varprojlim_n M/(\mathfrak{a}^n + \mathfrak{b}^m) M \right).$$

On the other hand, we have $(\mathfrak{a} + \mathfrak{b})^{2n} \subseteq \mathfrak{a}^n + \mathfrak{b}^n \subseteq (\mathfrak{a} + \mathfrak{b})^n$. If $x \in (\mathfrak{a} + \mathfrak{b})^n$, $x + \mathfrak{a}^n + \mathfrak{b}^n \subseteq (\mathfrak{a} + \mathfrak{b})^n$, which means each set in the basis $\{(\mathfrak{a} + \mathfrak{b})^n\}$ is open in the topology generated by the basis $\{\mathfrak{a}^n + \mathfrak{b}^n\}$, and vice versa. Thus, they generate a same topology, and therefore

$$M^{\mathfrak{a}+\mathfrak{b}} = \varprojlim M/(\mathfrak{a} + \mathfrak{b})^n M = \varprojlim M/(\mathfrak{a}^n + \mathfrak{b}^n) M.$$

We construct two homomorphisms $\varprojlim_m \left(\varprojlim_n M/(\mathfrak{a}^n + \mathfrak{b}^m) M \right) \xrightarrow[\Psi]{\Phi} \varprojlim M/(\mathfrak{a}^n + \mathfrak{b}^n) M$. When

$k \geq \max(m, n)$, $\mathfrak{a}^k + \mathfrak{b}^k \subseteq \mathfrak{a}^n + \mathfrak{b}^m$, so there is a canonical quotient map $\psi_{nm} : M/(\mathfrak{a}^k + \mathfrak{b}^k) M \rightarrow M/(\mathfrak{a}^n + \mathfrak{b}^m) M$. Let $\Phi((x)_{nm}) = (x)_{kk}$ be the diagonal map, and let $\Psi((y)_k) = \psi_{nm}((y)_k)$.

10.6. If $\mathfrak{a} \subseteq \mathfrak{J}$, let $x \notin \mathfrak{m}$. We can find an open neighborhood $x + \mathfrak{a}$ of x such that $(x + \mathfrak{a}) \cap \mathfrak{m} = \emptyset$.

By [Proposition 10.1](#), the closure of $\mathfrak{m}$ is $\bigcap_{n \geq 1} (\mathfrak{m} + \mathfrak{a}^n)$. If $\mathfrak{a} \not\subseteq \mathfrak{m}$, we have $\mathfrak{m} + \mathfrak{a}^n = A$, otherwise $\mathfrak{a} \subseteq \sqrt{\mathfrak{a}^n} \subseteq \mathfrak{m}$.

10.7. Let M be a finitely generated A -module. By [Proposition 10.13](#), $\hat{M} = M_{\hat{A}}$, so $\ker(M \rightarrow M_{\hat{A}}) = \bigcap_{n=1}^{\infty} \mathfrak{a}^n M$.

If $\mathfrak{a} \subseteq \mathfrak{J}$, every element in $1 + \mathfrak{a}$ is a unit, and by [Proposition 10.17](#), $\bigcap_{n=1}^{\infty} \mathfrak{a}^n M = 0$. If N is an arbitrary A -module and $x \in \ker(N \rightarrow N_{\hat{A}})$, then $x \in \ker(Ax \rightarrow (Ax)_{\hat{A}})$ and Ax is finitely generated.

If $\mathfrak{a} \not\subseteq \mathfrak{m}$, we have $\mathfrak{m} + \mathfrak{a}^n = A$. $(A/\mathfrak{m})/(\mathfrak{a}^n(A/\mathfrak{m})) = A/(\mathfrak{m} + \mathfrak{a}^n) = 0$, so the $\mathfrak{a}$ -adic completion of $A/\mathfrak{m}$ is 0. Hence, $A/\mathfrak{m} \rightarrow (A/\mathfrak{m})_{\hat{A}}$ isn't injective.

10.8. By [Exercise 1.5\(i\)](#) and [Corollary 1.6\(i\)](#), the only maximal ideal of B is $\mathfrak{m} = (z_1, \dots, z_n)$.

The $\mathfrak{m}$ -adic topologies on A , B and C are respectively equal to the induced topologies from the $\mathfrak{m}$ -adic topology on $\mathbb{C}[z_1, \dots, z_n]$. Apply [Proposition 10.2](#) to $\mathbb{C}[z_1, \dots, z_n] \subseteq A \subseteq B \subseteq C$.

By [Proposition 10.14](#), C is a flat A -algebra. Then use [Exercise 10.7](#) and [Exercise 3.17](#).

10.9. Assume that we have constructed g_k and h_k such that $f - g_k h_k \in \mathfrak{m}^k[x]$. Denote $w_k = f - g_k h_k$. Since there exist u_k and v_k such that $1 - u_k g_k - v_k h_k \in \mathfrak{m}[x]$, we have $w_k - w_k u_k g_k - w_k v_k h_k \in \mathfrak{m}^{k+1}[x]$.

Applying polynomial division algorithm, we have $w_k u_k = q_k h_k + a_k$ with $\deg a_k < \deg h_k$. Since h_k monic, the division process shows that $q_k \in \mathfrak{m}^k[x]$, so $a_k \in \mathfrak{m}^k[x]$. Let $b_k = w_k v_k + q_k g_k$, then $b_k \in \mathfrak{m}^k[x]$. Note that $\deg w_k < n$, we have $\deg b_k < \deg g_k$ in the polynomial ring $(A/\mathfrak{m}^{k+1})[x]$.

Let $g_{k+1} = g_k + b_k$, $h_{k+1} = h_k + a_k$, then $f - g_{k+1} h_{k+1} \in \mathfrak{m}^{k+1}[x]$. Since A complete, the coefficients of g_k and h_k converge, so we have $g, h \in A[x]$ such that $g \equiv g_k \pmod{\mathfrak{m}^k}$ and $h \equiv h_k \pmod{\mathfrak{m}^k}$. Therefore, $f - gh \in \bigcap_{k=1}^{\infty} \mathfrak{m}^k[x]$. By [Proposition 10.17](#), $\bigcap_{k=1}^{\infty} \mathfrak{m}^k[x] = 0$.

10.10. (i) $\bar{f}(x) = (x - \alpha)\bar{g}(x)$.

(ii) $\mathbb{Z}_7/7\mathbb{Z}_7 = \mathbb{Z}/7\mathbb{Z}$ by [Proposition 10.15\(iii\)](#). we also have $(x^2 - \bar{2}) = (x + \bar{3})(x - \bar{3})$ in $\mathbb{Z}/7\mathbb{Z}$.

(iii) $A = k[[x]]$ is a local ring with $\mathfrak{m} = (x)$. $\bar{f}(\bar{0}, y) = (y - \bar{a})\bar{g}(\bar{0}, y) \in A[y]$, so $a \in A$. Lift back by [Exercise 10.9](#) and let $y(x) = a$.

10.11. Let A be the ring of germs of $C^\infty(\mathbb{R})$ functions. If $f(0) \neq 0$, then $f(x)$ is always not 0 in A , so $\mathfrak{m} = \{f(x) \in A : f(0) = 0\}$ is its only maximal ideal.

$\mathfrak{m}^n = \{f(x) \in A : f^{(0)}(0) = \dots = f^{(n-1)}(0) = 0\}$, so $A/\mathfrak{m}^n \cong \mathbb{R}[x]/(x^n)$ and therefore $\hat{A} = \mathbb{R}[[x]]$. By Borel's theorem, $A \rightarrow \hat{A}$ is surjective, so $\hat{A}$ is a finitely generated A -module.

If A is Noetherian, by [Proposition 10.17](#), $\bigcap_{n=1}^{\infty} \mathfrak{m}^n = 0$, but $e^{-1/x^2} \in \bigcap_{n=1}^{\infty} \mathfrak{m}^n$.

10.12. By [Proposition 10.14](#), $A[[x_1, \dots, x_n]]$ is flat over $A[x_1, \dots, x_n]$, but from [Exercise 2.5](#) and [Exercise 2.8\(ii\)](#), $A[x_1, \dots, x_n]$ is flat over A . Then use [Exercise 2.8\(ii\)](#) again.

11 Dimension Theory

11.1. Move P to the origin to remove the constant term of f . $\mathfrak{m} = (x_1, \dots, x_n)/(f)$, so $\mathfrak{m}/\mathfrak{m}^2 = (x_1, \dots, x_n)/((x_1, \dots, x_n)^2 + (f))$. $\{x_i\}_{i=1}^n$ is a set of generators $\mathfrak{m}/\mathfrak{m}^2$. If $f \in (x_1, \dots, x_n)^2$, $\{x_i\}_{i=1}^n$ is a basis. If $f \notin (x_1, \dots, x_n)^2$, the existence of linear term of f means $\{x_i\}_{i=1}^n$ is linearly dependent.

Use [Corollary 11.18](#) on $A_{\mathfrak{m}} = k[x_1, \dots, x_n]_{\mathfrak{m}}/(f)_{\mathfrak{m}}$, then [Theorem 11.22](#).

11.2. The map $k[t_1, \dots, t_d]/(t_1, \dots, t_d)^n \rightarrow A/(x_1, \dots, x_d)^n$ is injective. Use [Corollary 7.16\(iii\)](#) and a similar step as [Exercise 10.5](#) to prove that $\mathfrak{m}$ -adic topology and $(x_1, \dots, x_d)$ -adic topology are same.

From [Proposition 10.22\(ii\)](#), $G_{(t_1, \dots, t_d)}(k[[t_1, \dots, t_d]]) \cong G_{(t_1, \dots, t_d)}(k[t_1, \dots, t_d]) \cong k[t_1, \dots, t_d]$. Note that every homogeneous element in $G_{(x_1, \dots, x_d)}(A)$ has a preimage in $k[t_1, \dots, t_d]$, so $G_{(x_1, \dots, x_d)}(A)$ is a finitely generated $k[t_1, \dots, t_d]$ -module. Then use [Proposition 10.24](#).

11.3. Let $\dim V = d$. From [Noether normalization lemma](#), $A(V)$ integral over $B = k[x_1, \dots, x_d]$. Since B is UFD, it is integrally closed. Apply [Exercise 5.3](#) on $k \otimes_k k[x] \rightarrow \bar{k} \otimes_k k[x]$, we also have $C = \bar{k}[x_1, \dots, x_d]$ integral over B .

Let $\mathfrak{m}$ be a maximal ideal in $A(V)$. By [Theorem 5.10](#), $\mathfrak{m} \cap B$ is maximal in B , and can be lifted to a prime ideal $\mathfrak{p}$ in C , and $\mathfrak{p}$ is contained in a maximal ideal $\mathfrak{n}$, with $\mathfrak{n} \cap B = \mathfrak{m} \cap B$. If there is a chain in $A(V)$ which ends at $\mathfrak{m}$, intersect with B and use [Theorem 5.16](#) to lift to C , we have a chain ends at $\mathfrak{n}$. Therefore, $\dim C_{\mathfrak{n}} \geq \dim A(V)_{\mathfrak{m}}$. Note that $\dim C_{\mathfrak{n}} = d$ by [Theorem 11.25](#).

11.4. The maximal ideals in $S^{-1}A$ is $S^{-1}\mathfrak{p}_i$. Localize at $S^{-1}\mathfrak{p}_i$ we have a ring $K[x_{m_i+1}, \dots, x_{m_{i+1}}]$, where $K = k[x_1, x_1^{-1}, \dots]$ without indeterminates $x_{m_i+1}, \dots, x_{m_{i+1}}$, so it is a field. Use [Exercise 7.9](#).

11.5. Let $\gamma : (M) \rightarrow K(A_0)$. Define the Poincaré series of M to be

$$P(M, t) = \sum_{n=0}^{\infty} \gamma(M_n) t^n \in K(A_0)[[t]].$$

11.6. From [Exercise 4.7\(ii\)](#), if $\mathfrak{p}_0 \subsetneq \dots \subsetneq \mathfrak{p}_n$ is a chain in A , then $\mathfrak{p}_0[x] \subsetneq \dots \subsetneq \mathfrak{p}_n[x] \subsetneq \mathfrak{p}_n[x] + (x)$ is a chain in $A[x]$.

Let $f : A \rightarrow A[x]$ be the embedding. The fiber of f^* at $\mathfrak{p}$ is $\text{Spec}(k(\mathfrak{p}) \otimes_A A[x]) = \text{Spec}(k(\mathfrak{p})[x])$. Since $\dim(k(\mathfrak{p})[x]) = 1$, a chain in $A[x]$ whose contraction is $\mathfrak{p}$ contains no more than 2 primes, so a chain in $A[x]$ contains at most $2 + 2 \dim A$ prime ideals.

11.7. Localize at $\mathfrak{p}$, and denote $r = \text{height } \mathfrak{p} = \dim A_{\mathfrak{p}}$. By [Theorem 11.14](#), there is a $\mathfrak{p}$ -primary ideal $\mathfrak{q}$ generated by r elements, so $\mathfrak{q}[x]$ also generated by r elements. From [Exercise 4.7\(iii\)](#), $\mathfrak{q}[x]$ is $\mathfrak{p}[x]$ -primary, then use [Theorem 11.14](#) again we have $\text{height } \mathfrak{p}[x] \leq r$. Then use [Exercise 11.6](#).

SUMMARY

Chapter 1 - Rings and Ideals

This chapter provides a broad overview of commutative algebra and introduces many of the themes developed later in the book.

The exercises introduce the spectrum of a ring, which becomes a central object in later studies.

Chapter 2 - Modules

Important results and ideas: [Nakayama's Lemma](#), changing basis ($M_B = B \otimes_A M$).

The exercises also introduce direct limits and absolutely flat rings.

[Exercise 2.15](#) is important in direct limit calculations.

Chapter 3 - Rings and Modules of Fractions

The notion of $S(\mathfrak{a})$ is particularly important for primary decomposition.

[Exercise 3.8](#) shows that saturated multiplicatively closed subsets of A are in one-to-one correspondence with distinct localizations $S^{-1}A$.

Chapter 4 - Primary Decomposition

[Proposition 4.4](#) and [Proposition 4.8](#) are fundamental for understanding primary ideal structure.

[Proposition 4.9](#) is useful for computations.

Chapter 5 - Integral Dependence and Valuations

The most important chapter in the book.

Important results: [Lying Over Theorem](#), [Noether Normalization Lemma](#).

[Exercise 5.10](#) connects the going-up and going-down properties of a ring homomorphism with topological properties of the induced map on spectra, but the ideas are not used further in the book.

[Exercise 5.25](#) uses Jacobson rings to generalize Zariski's Lemma.

Chapter 6 & 7 - Chain Conditions & Noetherian Rings

Most exercises further refine old concepts rather than introducing new ideas.

Chapter 8 - Artinian Rings

An important result is [Proposition 8.5](#).

[Exercise 8.3](#) provides another generalization of Zariski's Lemma.

Chapter 9 - Discrete Valuation Rings and Dedekind Domains

This chapter is closely connected to algebraic number theory. [Proposition 9.2](#) and [Theorem 9.3](#) appear repeatedly throughout the exercises.

Chapter 10 - Completions

Introduces completions and topological rings. For Noetherian rings, [Proposition 10.13](#) is important. [Proposition 10.15](#) and [Krull intersection theorem](#) are useful in computations.

The exercises show that two linear topologies defined by two filtrations are equivalent when each filtration eventually dominates the other.

The proof of [Hensel's lemma](#) is straightforward, but it involves many details that must be handled carefully.

Chapter 11 - Dimension Theory

Important result: [Dimension Theorem for Noetherian local rings](#).

This chapter mainly serves as an introduction to the subject, and a deeper understanding generally requires further study.