

1 Rings and Ideals

IDEALS. PRIME AND MAXIMAL IDEALS

Proposition 1.1. *There is a one-to-one order-preserving correspondence between the ideals $\mathfrak{b}$ of A which contain $\mathfrak{a}$, and the ideals $\bar{\mathfrak{b}}$ of $A/\mathfrak{a}$, given by $\mathfrak{b} = \phi^{-1}(\bar{\mathfrak{b}})$.*

Corollary 1.5. *Every non-unit of A is contained in a maximal ideal.*

Corollary 1.6. (i) *Let A be a ring and $\mathfrak{m} \neq (1)$ an ideal of A such that every $x \in A - \mathfrak{m}$ is a unit in A . Then A is a local ring and $\mathfrak{m}$ its maximal ideal.*

(ii) *Let A be a ring and $\mathfrak{m}$ a maximal ideal of A , such that every element of $1 + \mathfrak{m}$ is a unit in A , then A is a local ring.*

NILRADICAL AND JACOBSON RADICAL

We denote nilradical to be $\mathfrak{N}$ and Jacobson radical to be $\mathfrak{J}$.

Proposition 1.8. *The nilradical of A is the intersection of all the prime ideals of A .*

Sketch proof. Assume that $f \notin \mathfrak{N}$. Let Σ be a set of ideals $\mathfrak{a}$ with property $f^n \notin \mathfrak{a}$. Then Σ has maximal element which is prime. $\square$

Proposition 1.9. $x \in \mathfrak{J} \Leftrightarrow 1 - xy$ is a unit in A for all $y \in A$.

OPERATIONS ON IDEALS

Two ideals $\mathfrak{a}, \mathfrak{b}$ are said to be **coprime** if $\mathfrak{a} + \mathfrak{b} = 1$. Thus we have $\mathfrak{a} \cap \mathfrak{b} = \mathfrak{a}\mathfrak{b}$.

Proposition 1.10 (Chinese Remainder Theorem). *Define ϕ to be*

$$\phi : A \rightarrow \prod_{i=1}^n (A/\mathfrak{a}_i), \quad \phi(x) = (x + \mathfrak{a}_1, \dots, x + \mathfrak{a}_n).$$

(i) *If $\mathfrak{a}_i$ and $\mathfrak{a}_j$ are coprime whenever $i \neq j$, then $\mathfrak{a}_1 \cdots \mathfrak{a}_n = \bigcap_{i=1}^n \mathfrak{a}_i$.*

(ii) *ϕ is surjective $\Leftrightarrow \mathfrak{a}_i$ and $\mathfrak{a}_j$ are coprime whenever $i \neq j$.*

(iii) *ϕ is injective $\Leftrightarrow \bigcap_{i=1}^n \mathfrak{a}_i = (0)$.*

Proposition 1.11. *Denote $\mathfrak{a}$ to be ideal and $\mathfrak{p}$ to be prime ideal.*

(i) *If $\mathfrak{a} \subseteq \bigcup_{i=1}^n \mathfrak{p}_i$, then $\mathfrak{a} \subseteq \mathfrak{p}_i$ for some i .*

(ii) *If $\mathfrak{p} \supseteq \bigcap_{i=1}^n \mathfrak{a}_i$, then $\mathfrak{p} \supseteq \mathfrak{a}_i$ for some i . If $\mathfrak{p} = \bigcap_{i=1}^n \mathfrak{a}_i$, then $\mathfrak{p} = \mathfrak{a}_i$ for some i .*

The **ideal quotient** of $\mathfrak{a}$ and $\mathfrak{b}$ is

$$(\mathfrak{a} : \mathfrak{b}) = \{x \in A : x\mathfrak{b} \subseteq \mathfrak{a}\}.$$

The annihilator of $\mathfrak{b}$ is $\text{Ann}(\mathfrak{b}) = (0 : \mathfrak{b})$.

The **radical** of $\mathfrak{a}$ is

$$\sqrt{\mathfrak{a}} = \{x \in A : x^n \in \mathfrak{a} \text{ for some } n > 0\}.$$

EXTENSION AND CONTRACTION

Let $f : A \rightarrow B$ be a ring homomorphism. Let $\mathfrak{a} \in A$ and $\mathfrak{b} \in B$. Define

$$\mathfrak{a}^e = Bf(\mathfrak{a}), \quad \mathfrak{b}^c = f^{-1}(\mathfrak{b})$$

to be the **extension** of $\mathfrak{a}$ and the **contraction** of $\mathfrak{b}$, respectively. If $\mathfrak{b}$ is prime, $\mathfrak{b}^c$ is prime; if $\mathfrak{a}$ is prime, $\mathfrak{a}^e$ need not be prime.

Proposition 1.17. $\mathfrak{a} \subseteq \mathfrak{a}^{ec}$, $\mathfrak{b} \supseteq \mathfrak{b}^{ce}$.

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

Idempotent is an element e which satisfies $e^2 = e$.

A ring in which every element is an idempotent is a **Boolean ring**.

A topological space X is **reducible** if it can be written as a union $X = X_1 \cup X_2$ of two closed proper subsets X_1, X_2 of X . Equivalently, X is **irreducible** if every pair of non-empty open sets in X intersect, or if every non-empty open set is dense in X .

Let A be a ring and let $\text{Spec}(A)$ be the set of all prime ideals of A . For each subset E of A , let $V(E)$ denote the set of all prime ideals of A which contain E . By [Exercise 1.15](#), the sets $V(E)$ satisfy the axioms for closed sets in a topological space. The resulting topology on $\text{Spec}(A)$ is called **Zariski topology**, and $\text{Spec}(A)$ is called the **prime spectrum** of A .

For each $f \in A$, the complement of $V(f)$ in $\text{Spec}(A)$, denote to be X_f , is called **basic open set** of $\text{Spec}(A)$.

The subspace of $\text{Spec}(A)$ consisting of the maximal ideals of A with the induced topology, is called the **maximal spectrum** of A and is denoted by $\text{Max}(A)$.

Proposition ([Exercise 1.27](#)). *Let k be algebraically closed. Then the affine space k^n with its classical Zariski topology is homeomorphic to $\text{Max}(k[x_1, \dots, x_n])$.*

2 Modules

Proposition 2.1. (i) If $L \supseteq M \supseteq N$ are A -modules, then $(L/N)/(M/N) \cong L/M$.

(ii) If M_1, M_2 are submodules of M , then $(M_1 + M_2)/M_1 \cong M_2/(M_1 \cap M_2)$.

An A -module is **faithful** if $\text{Ann}(M) = 0$.

FINITELY GENERATED MODULES

Proposition 2.3. M is a finitely generated A -module $\Leftrightarrow M$ is isomorphic to a quotient of A^n for some integer $n > 0$. More precisely, n is the number of generators of M and may not be unique.

Proposition 2.4 (**Cayley–Hamilton theorem**). Let M be a finitely generated A -module, let $\mathfrak{a}$ be an ideal of A , and let ϕ be an A -module endomorphism of M such that $\phi(M) \subseteq \mathfrak{a}M$. Then ϕ satisfies an equation of the form

$$\phi^n + a_1\phi^{n-1} + \cdots + a_n = 0$$

where the a_i are in $\mathfrak{a}$.

Proposition 2.6 (**Nakayama's lemma**). Let M be a finitely generated A -module and $\mathfrak{a}$ an ideal of A contained in the Jacobson radical of A . Then $\mathfrak{a}M = M$ implies $M = 0$.

EXACT SEQUENCES

Proposition 2.10 (Snake lemma). Let

$$\begin{array}{ccccccc} 0 & \longrightarrow & M' & \xrightarrow{u} & M & \xrightarrow{v} & M'' \longrightarrow 0 \\ & & \downarrow f' & & \downarrow f & & \downarrow f'' \\ 0 & \longrightarrow & N' & \xrightarrow{u'} & N & \xrightarrow{v'} & N'' \longrightarrow 0 \end{array}$$

be a commutative diagram of A -modules and homomorphisms, with the rows exact. Then there exists an exact sequence

$$0 \longrightarrow \ker(f') \xrightarrow{\bar{u}} \ker(f) \xrightarrow{\bar{v}} \ker(f'') \xrightarrow{\delta} \text{coker}(f') \xrightarrow{\bar{u}'} \text{coker}(f) \xrightarrow{\bar{v}'} \text{coker}(f'') \longrightarrow 0.$$

Let C be a class of A -modules and let λ be a function on C with values in $\mathbb{Z}$. The function λ is **additive** if for each short exact sequence

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

in which all terms belong to C , we have $\lambda(M') - \lambda(M) + \lambda(M'') = 0$.

TENSOR PRODUCT

Proposition 2.12 (Universal property). Let M, N be A -modules. The tensor product of M and N over A is an A -module $M \otimes_A N$ together with an A -bilinear map

$$f : M \times N \rightarrow M \otimes_A N, \quad (m, n) \mapsto m \otimes n,$$

such that, for every A -module P and every A -bilinear map $\tau : M \times N \rightarrow P$, there exists a unique A -linear map $\tilde{\tau} : M \otimes_A N \rightarrow P$ such that $\tau = \tilde{\tau} \circ f$. Equivalently, the following diagram commutes:

$$\begin{array}{ccc} M \times N & \xrightarrow{f} & M \otimes_A N \\ & \searrow \tau & \downarrow \tilde{\tau} \\ & & P \end{array}$$

Proposition 2.15. Let A, B be rings, let M be an A -module, P a B -module and N an (A, B) -bimodule. then

$$(M \otimes_A N) \otimes_B P \cong M \otimes_A (N \otimes_B P).$$

Let M be an A -module and $f : A \rightarrow B$ be a homomorphism of rings. Define $M_B = B \otimes_A M$.

If the functor $T_N : M \mapsto M \otimes_A N$ on the category of A -modules is exact, then N is said to be a **flat** A -module.

Proposition 2.19. For an A -module N , the following are equivalent:

- (i) N is flat.
- (ii) If $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is any exact sequence of A -modules, then $0 \rightarrow M' \otimes N \rightarrow M \otimes N \rightarrow M'' \otimes N \rightarrow 0$ is exact.
- (iii) If $f : M' \rightarrow M$ is injective, then $f \otimes 1 : M' \otimes N \rightarrow M \otimes N$ is injective.
- (iv) If $f : M' \rightarrow M$ is injective and M, M' are finitely generated, then $f \otimes 1 : M' \otimes N \rightarrow M \otimes N$ is injective.

ALGEBRA

Let $f : A \rightarrow B$ be a ring homomorphism. If $a \in A$ and $b \in B$, define a product

$$ab = f(a)b.$$

Thus, B has an A -module structure. The ring B is said to be an **A -algebra**.

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A partially ordered set I is said to be a **directed** set if for each pair $i, j \in I$ there exists $k \in I$ such that $i \leq k$ and $j \leq k$.

Let A be a ring, let I be a directed set and let $(M_i)_{i \in I}$ be a family of A -modules indexed by I . For each pair $i, j \in I$ such that $i \leq j$, let $\mu_{ij} : M_i \rightarrow M_j$ be an A -homomorphism, and suppose:

- (1) μ_{ii} is the identity mapping of M_i , for all $i \in I$;
- (2) $\mu_{ik} = \mu_{jk} \circ \mu_{ij}$ whenever $i \leq j \leq k$.

Then the modules M_i and homomorphisms μ_{ij} are said to form a **direct system** $\mathbf{M} = (M_i, \mu_{ij})$ over the directed set I .

Let D be the submodule of $\bigoplus M_i$ generated by all elements of the form $x_i - \mu_{ij}(x_i)$ whenever $i \leq j$ and $x_i \in M_i$. The **direct limit** of the direct system $\mathbf{M}$ is defined to be

$$\varinjlim M_i = \bigoplus_{i \in I} M_i / D.$$

Let $\mu : \bigoplus M_i \rightarrow \varinjlim M_i$ be the projection of quotient module. Define μ_i to be the restriction of μ to M_i .

A ring A (not necessarily commutative) is **von Neumann regular** if for every $a \in A$, there is an $x \in A$ with $a = axa$. A commutative ring A is **absolutely flat** if every A -module is flat. By [Exercise 2.27](#), absolute flat and von Neumann regular are identical properties.

3 Rings and Modules of Fractions

MULTIPLICATIVELY CLOSED SETS

Proposition 3.3. *The operation S^{-1} is exact.*

Proposition 3.5. *Let M be an A -module. Then there exists a unique $S^{-1}A$ module isomorphism $f : S^{-1}A \otimes_A M \rightarrow S^{-1}M$.*

LOCAL PROPERTIES

Proposition. *Let P be a property of a ring A (or of an A -module M). Then the following are equivalent:*

- (i) A (or M) has $P \Leftrightarrow A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has P for each prime ideal $\mathfrak{p}$ of A ;
- (ii) A (or M) has $P \Leftrightarrow A_{\mathfrak{m}}$ (or $M_{\mathfrak{m}}$) has P for each maximal ideal $\mathfrak{m}$ of A .

*The property P is said to be a **local property**.*

Sketch proof. $(A_{\mathfrak{p}})_{\mathfrak{m}A_{\mathfrak{p}}} \cong A_{\mathfrak{m}}$

□

Proposition 3.8. *Let M, N be A -modules. Then the following are equivalent:*

- (i) $M = 0$;
- (ii) $M_{\mathfrak{p}} = 0$ for all prime ideals $\mathfrak{p}$ of A ;
- (iii) $M_{\mathfrak{m}} = 0$ for all maximal ideals $\mathfrak{m}$ of A ;

Proposition 3.10. *Let M be A -modules. Then the following are equivalent:*

- (i) M is a flat A -module;
- (ii) $M_{\mathfrak{p}}$ is a flat $A_{\mathfrak{p}}$ -module for each prime ideal $\mathfrak{p}$;
- (iii) $M_{\mathfrak{m}}$ is a flat $A_{\mathfrak{m}}$ -module for each maximal ideal $\mathfrak{m}$;

EXTENSIONS AND CONTRACTIONS OF FRACTIONS

If $\mathfrak{a}$ is an ideal in A , its extension $\mathfrak{a}^e$ in $S^{-1}A$ is $S^{-1}\mathfrak{a}$. Denote the contraction of $S^{-1}\mathfrak{a}$ in A to be $S(\mathfrak{a})$.

Proposition 3.11. (i) *Every ideal in $S^{-1}A$ is an extended ideal.*

(ii) *If $\mathfrak{a}$ is an ideal in A , then $S(\mathfrak{a}) = \mathfrak{a}^{ec} = \bigcup_{s \in S} (\mathfrak{a} : s)$.*

(iii) *$\mathfrak{a}$ is a contracted ideal $\Leftrightarrow$ no element of S is a zero-divisor in $A/\mathfrak{a}$.*

(iv) *Them prime ideal of $S^{-1}A$ are in one-to-one correspondence ($\mathfrak{p} \leftrightarrow S^{-1}\mathfrak{p}$) with the prime ideal of A which don't meet S .*

(v) *The operation S^{-1} commutes with formation of finite sums, products, intersections and radicals.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A multiplicatively closed subset S of a ring A is said to be **saturated** if

$$xy \in S \Leftrightarrow x \in S \text{ and } y \in S.$$

Let M be an A -module. An element $x \in M$ is a **torsion** element of M if there exist a non-zero-divisor $a \in A$ such that $ax = 0$. All torsion elements of M form a submodule of M , which is called the **torsion submodule** of M and denote by $T(M)$. If $T(M) = 0$, the module M is said to be **torsion free**.

Let M be an A -module. The **support** of M is defined to be the set $\text{Supp}(M)$ of prime ideals $\mathfrak{p}$ such that $M_{\mathfrak{p}} \neq 0$.

$\text{Spec}(A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} \otimes_A B)$ is called the **fiber** of $f^* : \text{Spec}(B) \rightarrow \text{Spec}(A)$ over $\mathfrak{p}$.

4 Primary Decomposition

PRIMARY IDEALS

An ideal $\mathfrak{q} \in A$ is **primary** if $\mathfrak{q} \neq A$ and if either

- (i) $x \notin \mathfrak{q} \implies y^n \in \mathfrak{q}$ for some $n > 0$;
- (ii) $x \notin \sqrt{\mathfrak{q}} \implies y \in \mathfrak{q}$.

Proposition 4.4. *Let $\mathfrak{q}$ be a $\mathfrak{p}$ -primary ideal, $x \in A$ an element. Then*

- (i) *if $x \in \mathfrak{q}$ then $(\mathfrak{q} : x) = (1)$;*
- (ii) *if $x \notin \mathfrak{q}$ then $(\mathfrak{q} : x)$ is $\mathfrak{p}$ -primary;*
- (iii) *if $x \notin \mathfrak{p}$ then $(\mathfrak{q} : x) = \mathfrak{q}$;*

Proposition 4.8. *Let S be a multiplicatively closed subset of A , and let $\mathfrak{q}$ be a $\mathfrak{p}$ -primary ideal.*

- (i) *If $S \cap \mathfrak{p} \neq \emptyset$, then $S^{-1}\mathfrak{q} = S^{-1}A$.*
- (ii) *If $S \cap \mathfrak{p} = \emptyset$, then $S^{-1}\mathfrak{q}$ is $S^{-1}\mathfrak{p}$ -primary and its contraction in A is $\mathfrak{q}$.*

UNIQUENESS

Proposition 4.5 (1st uniqueness theorem). *Let $\mathfrak{a}$ be a decomposable ideal and let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ be a minimal primary decomposition of $\mathfrak{a}$. Let $\mathfrak{p}_i = \sqrt{\mathfrak{q}_i}$. Then $\mathfrak{p}_i$ are precisely the prime ideals which occur in the set of ideals $\sqrt{(\mathfrak{a} : x)}$, and hence are independent of the particular decomposition of $\mathfrak{a}$.*

The prime ideals $\mathfrak{p}_i$ are said to belong to $\mathfrak{a}$, or to be **associated** with $\mathfrak{a}$. The minimal elements of the set $\{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$ are called the minimal or **isolated** prime ideals belonging to $\mathfrak{a}$. The others are called **embedded** prime ideals.

Proposition 4.9. *Let S be a multiplicatively closed subset of A and let $\mathfrak{a}$ be a decomposable ideal. Let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ be a minimal primary decomposition of $\mathfrak{a}$. Let $\mathfrak{p}_i = \sqrt{\mathfrak{q}_i}$ and suppose the $\mathfrak{q}_i$ numbered so that S meets $\mathfrak{p}_{m+1}, \dots, \mathfrak{p}_n$ but not $\mathfrak{p}_1, \dots, \mathfrak{p}_m$. Then we have minimal primary decompositions*

$$S^{-1}\mathfrak{a} = \bigcap_{i=1}^m S^{-1}\mathfrak{q}_i, \quad S(\mathfrak{a}) = \bigcap_{i=1}^m \mathfrak{q}_i.$$

A set Σ of prime ideals belonging to $\mathfrak{a}$ is said to be **isolated** if it satisfies the following condition: if $\mathfrak{p}'$ is a prime ideal belonging to $\mathfrak{a}$ and $\mathfrak{p}' \subseteq \mathfrak{p}$ for some $\mathfrak{p} \in \Sigma$, then $\mathfrak{p}' \in \Sigma$. Thus, every isolated set contains an isolated prime ideal.

Proposition 4.10 (2nd uniqueness theorem). *Let $\mathfrak{a}$ be a decomposable ideal, let $\mathfrak{a} = \bigcap_{i=1}^n \mathfrak{q}_i$ be a minimal primary decomposition of $\mathfrak{a}$, and let $\{\mathfrak{p}_{i_1}, \dots, \mathfrak{p}_{i_m}\}$ be an isolated set of prime ideals of $\mathfrak{a}$. Then the ideal $\mathfrak{q}_{i_1} \cap \dots \cap \mathfrak{q}_{i_m}$ is independent of the decomposition.*

Corollary 4.11. *The isolated primary components are uniquely determined by $\mathfrak{a}$.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

Let A be a ring and $\mathfrak{p}$ a prime ideal of A . The n th **symbolic power** of $\mathfrak{p}$ is defined to be the ideal $\mathfrak{p}^{(n)} = S_{\mathfrak{p}}(\mathfrak{p}^n)$.

5 Integral Dependence and Valuations

INTEGRAL DEPENDENCE

Proposition 5.1. *The following are equivalent:*

- (i) $x \in B$ is integral over A ;
- (ii) $A[x]$ is a finitely generated A -module;
- (iii) $A[x]$ is contained in a subring C of B such that C is a finitely generated A -module;
- (iv) There exist a faithful $A[x]$ -module M which is finitely generated as an A -module.

The set $\bar{A}$ of elements of B which are integral over A is called the **integral closure** of A in B . If $\bar{A} = A$, then A is said to be **integrally closed in B** . If $\bar{A} = B$, the ring B is said to be **integral over A** .

An integral domain is said to be **integrally closed** (without qualification) if it is integrally closed in its field of fractions.

THE GOING-UP THEOREM

Theorem 5.10 (**Lying over property**). *Let $A \subseteq B$ be rings, B integral over A , and let $\mathfrak{p}$ be a prime ideal of A . Then there exists a prime ideal $\mathfrak{q}$ of B such that $\mathfrak{q} \cap A = \mathfrak{p}$.*

Theorem 5.11 (Going-up theorem). *Let $A \subseteq B$ be rings, B integral over A . Let $\mathfrak{p}_1 \subseteq \cdots \subseteq \mathfrak{p}_n$ be a chain of prime ideals of A and $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_m$ ($m < n$) a chain of prime ideals of B such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq m$. Then the chain $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_m$ can be extended to a chain $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_n$ such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq n$.*

THE GOING-DOWN THEOREM

Proposition 5.13. *Let A be an integral domain. Then " A is integrally closed" is a local property.*

Theorem 5.16 (Going-down theorem). *Let $A \subseteq B$ be integral domains, A integrally closed, B integral over A . Let $\mathfrak{p}_1 \subseteq \cdots \subseteq \mathfrak{p}_n$ be a chain of prime ideals of A and $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_m$ ($m < n$) a chain of prime ideals of B such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq m$. Then the chain $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_m$ can be extended to a chain $\mathfrak{q}_1 \subseteq \cdots \subseteq \mathfrak{q}_n$ such that $\mathfrak{q}_i \cap A = \mathfrak{p}_i$ for $1 \leq i \leq n$.*

VALUATION RINGS

Let B be an integral domain. B is a **valuation ring** of $\text{Frac}(B)$ if, for each $x \neq 0$, either $x \in B$ or $x^{-1} \in B$ (or both).

Proposition 5.18. *Let B be a valuation ring, $K = \text{Frac}(B)$ its field of fractions. Then*

- (i) B is a local ring;
- (ii) If B' is a ring such that $B \subseteq B' \subseteq K$, then B' is a valuation ring of K ;
- (iii) B is integrally closed.

Theorem 5.21. *Let K be a field, Ω an algebraically closed field. Let Σ be the set of all pairs (A, f) , where A is a subring of K and $f : A \rightarrow \Omega$ a homomorphism. Define a partial order of Σ to be:*

$$(A, f) \leq (B, g) \Leftrightarrow A \subseteq B \text{ and } g|_A = f.$$

Then the maximal element of Σ is a valuation ring of K .

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A ring homomorphism $f : A \rightarrow B$ is said to have the **going-up property** if for $\mathfrak{p}_1 \subseteq \mathfrak{p}_2$ and $\mathfrak{p}_1 = f^{-1}(\mathfrak{q}_1)$, there exist a $\mathfrak{q}_2$ such that $\mathfrak{q}_1 \subseteq \mathfrak{q}_2$ and $f^{-1}(\mathfrak{q}_2) = \mathfrak{p}_2$. Reversing the inclusions of prime ideals we have the **going-down property**.

Proposition (Noether normalization lemma). *Let k be a field and let $A \neq 0$ be a finitely generated k -algebra. Then there exist elements $y_1, \dots, y_r \in A$ which are algebraically independent over k and such that A is integral over $k[y_1, \dots, y_r]$.*

A ring A satisfied the following equivalent property is a **Jacobson ring**.

- (i) Every prime ideal in A is an intersection of maximal ideals.
- (ii) In every homomorphic image of A the nilradical is equal to the Jacobson radical.
- (iii) Every prime ideal in A which is not maximal is equal to the intersection of the prime ideals which contain it strictly.

Let A be a valuation ring of a field K . The group U of units of A is a subgroup of the multiplicative group K^* of K . Let $\Gamma = K^*/U$. If $\xi, \iota \in \Gamma$ are represented by $x, y \in K$, define $\xi \geq \iota$ to mean $xy^{-1} \in A$. Then Γ is a totally ordered abelian group. It is called the **value group** of A .

Conversely, let Γ be a totally ordered abelian group. A **valuation** of K with values in Γ is a mapping $v : K^* \rightarrow \Gamma$ such that

- (1) $v(xy) = v(x) + v(y)$,
- (2) $v(x + y) \geq \min(v(x), v(y))$,

for all $x, y \in K^*$. Then the set of elements $x \in K^*$ such that $v(x) \geq 0$ is a valuation ring of K . This ring is called the **valuation ring** of v , and the subgroup $v(K^*)$ of Γ is the **value group** of v . Thus the concepts of valuation ring and valuation are equivalent.

6 Chain Conditions

Let Σ be a partially ordered set. Σ said to be satisfying the **ascending chain condition** (a.c.c) if, for every increasing sequence $x_1 \leq x_2 \leq \dots$ in Σ there exists n such that $x_n = x_{n+1} = \dots$.

Similarly, Σ said to be satisfying the **descending chain condition** (d.c.c) if, for every decreasing sequence $x_1 \geq x_2 \geq \dots$ in Σ there exists n such that $x_n = x_{n+1} = \dots$.

A module satisfying the ascending chain condition is said to be **Noetherian**. A module satisfying the descending chain condition is said to be **Artinian**.

Proposition 6.3. *Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence of A -modules. Then*

- (i) M is Noetherian $\Leftrightarrow M'$ and M'' are Noetherian.
- (ii) M is Artinian $\Leftrightarrow M'$ and M'' are Artinian.

Proposition 6.5. *Let A be a Noetherian (resp. Artinian) ring, M a finitely generated A -module. Then M is Noetherian (resp. Artinian).*

A **chain** of submodules of a module M is a sequence (M_i) of submodules of M such that

$$M = M_0 \supsetneq M_1 \supsetneq \dots \supsetneq M_n = 0.$$

The **length** of the chain is n . A **composition series** of M is a maximal chain, that is one which no extra submodules can be inserted, or equivalently M_{i-1}/M_i is simple.

Proposition 6.7. *Suppose that M has a composition series of length n . Then every composition series of M has length n , and every chain can be extended to a composition series.*

The length of a composition series of M is called the **length** of M , denoted as $l(M)$.

Proposition 6.10. *For a vector space, finite dimension $\Leftrightarrow$ finite length $\Leftrightarrow$ a.c.c. $\Leftrightarrow$ d.c.c.. If the conditions are satisfied, length equals dimension.*

DEFINITIONS AND PROPOSITIONS FROM EXERCISES

A topological space X is said to be **Noetherian** if the open subsets of X satisfy the ascending chain condition. Or equivalently, the closed subsets satisfy descending chain condition.

7 Noetherian Rings

Proposition 7.1. *Any homomorphic image of Noetherian ring is Noetherian.*

Proposition 7.2. *Let A be a subring of B ; suppose that A is Noetherian and that B is finitely generated as an A -module. Then B is Noetherian.*

Theorem 7.5 (Hilbert's basis theorem). *If A is Noetherian, then the polynomial ring $A[x]$ is Noetherian.*

Proposition 7.8. *Let $A \subseteq B \subseteq C$ be rings. Suppose that A is Noetherian, that C is finitely generated as an A -algebra and that C is either (i) finitely generated as a B -module or (ii) integral over B . Then B is finitely generated as an A -algebra.*

Proposition 7.9 (Zariski's lemma). *Let k be a field, E a finitely generated k -algebra. If E is a field then it is a finite algebraic extension of k .*

PRIMARY DECOMPOSITION IN NOETHERIAN RINGS

Proposition 7.11. *In a Noetherian ring every ideal is a finite intersection of irreducible ideals.*

Proposition 7.12. *In a Noetherian ring every irreducible ideal is primary.*

Proposition 7.14. *In a Noetherian ring every ideal contains a power of its radical.*

Proposition 7.15. *In a Noetherian ring the nilradical is nilpotent.*

Proposition 7.16. *Let A be a Noetherian ring, $\mathfrak{m}$ a maximal ideal of A , $\mathfrak{q}$ any ideal of A . Then the following are equivalent:*

- (i) $\mathfrak{q}$ is $\mathfrak{m}$ -primary;
- (ii) $\sqrt{\mathfrak{q}} = \mathfrak{m}$;
- (iii) $\mathfrak{m}^n \subseteq \mathfrak{q} \subseteq \mathfrak{m}$ for some $n > 0$.

Proposition 7.17. *Let $\mathfrak{a} \neq (1)$ be an ideal in a Noetherian ring. Then the prime ideals which belong to $\mathfrak{a}$ are precisely the prime ideals which occur in the set of ideals $(\mathfrak{a} : x)$.*

8 Artinian Rings

Proposition 8.1. *In an Artinian ring every prime ideal is maximal.*

Proposition 8.3. *An Artinian ring has only a finite number of maximal ideals.*

The **dimension** of A is the supremum of the length of all chains of prime ideals in A .

Proposition 8.5. *A ring A is Artinian $\Leftrightarrow A$ is Noetherian and $\dim(A) = 0$.*

Proposition 8.6. *Let A be a Noetherian local ring, $\mathfrak{m}$ its maximal ideal. Then either $\mathfrak{m}^n \neq \mathfrak{m}^{n+1}$ for all n , or $\mathfrak{m}^n = 0$ for some n . In the latter case A is an Artinian local ring.*

Proposition 8.7 (Structure theorem of Artinian rings). *An Artinian ring A is uniquely (up to isomorphism) a finite direct product of Artinian local rings.*

Proposition 8.8. *Let A be an Artinian local ring. Then the following are equivalent:*

- (i) *every ideal in A is principal;*
- (ii) *the maximal ideal $\mathfrak{m}$ is principal;*
- (iii) $\dim_{A/\mathfrak{m}} (\mathfrak{m}/\mathfrak{m}^2) \leq 1$.

9 Discrete Valuation Rings and Dedekind Domain

10 Completion

TOPOLOGIES AND COMPLETIONS

A **topological abelian group** G is both a topological space and an abelian group, with two structural mappings

$$\begin{aligned} G \times G &\rightarrow G : (x, y) \mapsto x + y \\ G &\rightarrow G : x \mapsto -x \end{aligned}$$

being continuous.

If $\{0\}$ is closed in G , then G is Hausdorff.

Since $T_a(x) = x + a$ is a homeomorphism, the topology of G is uniquely determined by the neighborhoods of 0 in G .

Proposition 10.1. *Let H be the intersection of all neighborhoods of 0 in G . Then*

- (i) H is a subgroup.
- (ii) H is the closure of $\{0\}$.
- (iii) G/H is Hausdorff.
- (iv) G is Hausdorff $\Leftrightarrow H = 0$.

Assume that $0 \in G$ has a countable fundamental system of neighborhoods. A **Cauchy sequence** (x_ν) of G is that, for any neighborhood U of 0, there exists an integer $s(U)$ such that $x_m - x_n \in U$ for all $m, n \geq s(U)$.

Let $C(G)$ be the set of all Cauchy sequences in G . Define an equivalence relation $\sim$ on $C(G)$ where $(x_\nu) \sim (y_\nu)$ means $x_\nu - y_\nu \rightarrow 0$ in G . The **completion** $\hat{G}$ of G is the set of all equivalence classes in $C(G)/\sim$.

For each $x \in G$, define a homomorphism $\phi : G \rightarrow \hat{G}$ maps to a constant sequence (x) . Then ϕ is injective iff G is Hausdorff.

Assume further that $0 \in G$ has a fundamental system of neighborhoods consisting of a sequence of subgroups

$$G = G_0 \supseteq G_1 \supseteq \cdots \supseteq G_n \supseteq \cdots$$

For any $(x_\nu) \in C(G)$ and a fixed n , the image sequence $(\xi_\nu + G_n)$ in G/G_n is ultimately constant. Let ξ_n be this ultimate constant value.

As n varies, the projection $\theta_{n+1} : G/G_{n+1} \rightarrow G/G_n$ satisfies $\theta_{n+1}(\xi_{n+1}) = \xi_n$ for all n . The groups G/G_n combined with maps θ_n form an **inverse system**, and the group of all such coherent sequences (ξ_ν) is called the **inverse limit**, denoted as

$$\hat{G} = \varprojlim G/G_n.$$

A direct (or inverse) system is called **injective/surjective system**, if every transition map μ_{ij} is injective/surjective.

Proposition 10.2. *If $0 \rightarrow \{A_n\} \rightarrow \{B_n\} \rightarrow \{C_n\} \rightarrow 0$ is an exact sequence of inverse system, then*

$$0 \rightarrow \varprojlim A_n \rightarrow \varprojlim B_n \rightarrow \varprojlim C_n$$

is exact. If $\{A_n\}$ is surjective, then

$$0 \rightarrow \varprojlim A_n \rightarrow \varprojlim B_n \rightarrow \varprojlim C_n \rightarrow 0$$

is exact.

Corollary 10.3. Let $0 \rightarrow G' \rightarrow G \rightarrow G'' \rightarrow 0$ be an exact sequence of groups. Let G have the topology defined by a sequence $\{G_n\}$ of subgroups, and give G' and G'' the topologies by the contraction and extension of $\{G_n\}$, respectively. Then

$$0 \rightarrow \hat{G}' \rightarrow \hat{G} \rightarrow \hat{G}'' \rightarrow 0$$

is exact.

Corollary 10.4. $\hat{G}_n$ is a subgroup of $\hat{G}$ and $\hat{G}/\hat{G}_n \cong G/G_n$.

Corollary 10.5. $\hat{\hat{G}}_n \cong \hat{G}_n$.

If $G \rightarrow \hat{G}$ is an isomorphism, G is said to be **complete**.

Let A be a ring, $\mathfrak{a}$ its ideal. The topology defined by $G_n = \mathfrak{a}^n$ ($G_0 = A$) is called the **$\mathfrak{a}$ -adic topology**, and the ring A combined with this topology is a **topological ring**.

FILTRATIONS

A chain $M = M_0 \supseteq M_1 \supseteq \cdots \supseteq M_n \cdots$ is called a **filtration** of M , and denoted by (M_n) . It is an **$\mathfrak{a}$ -filtration** if $\mathfrak{a}M_n \subseteq M_{n+1}$ for all n , and a **stable $\mathfrak{a}$ -filtration** if $\mathfrak{a}M_n = M_{n+1}$ for all sufficiently large n .

The **$\mathfrak{a}$ -adic completion** of M is defined to be

$$\hat{M} = \varprojlim M/\mathfrak{a}^n M.$$

GRADED RINGS AND MODULES

A **graded ring** is a ring A together with a family $(A_n)_{n \geq 0}$ of additive subgroups of A , such that $A = \bigoplus_{n=0}^{\infty} A_n$ and $A_m A_n \subseteq A_{m+n}$ for all $m, n \geq 0$.

Proposition 10.7. A graded ring A is Noetherian, iff A_0 is Noetherian and A is finitely generated as an A_0 -algebra.

Proposition 10.9 (Artin-Rees lemma). Let A be a Noetherian ring, $\mathfrak{a}$ an ideal in A , M a finitely-generated A -module, (M_n) a stable $\mathfrak{a}$ -filtration of M . If M' is a submodule of M , then $(M' \cap M_n)$ is a stable $\mathfrak{a}$ -filtration of M' .

Proposition 10.12. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be an exact sequence of finitely-generated modules over a Noetherian ring A , then the sequence of $\mathfrak{a}$ -adic completion $0 \rightarrow \hat{M}' \rightarrow \hat{M} \rightarrow \hat{M}'' \rightarrow 0$ is exact.

Proposition 10.13. If M is finitely-generated A -module, $\hat{A} \otimes_A M \rightarrow \hat{M}$ is surjective. If A is Noetherian, then $\hat{A} \otimes_A M \rightarrow \hat{M}$ is an isomorphism.

Proposition 10.14. If A is a Noetherian ring, the $\mathfrak{a}$ -adic completion $\hat{A}$ is a flat A -algebra.

Proposition 10.15. If A is a Noetherian ring, $\hat{A}$ its $\mathfrak{a}$ -adic completion, then

- (i) $\hat{\mathfrak{a}} = \hat{A}\mathfrak{a} \cong \hat{A} \otimes_A \mathfrak{a}$;
- (ii) $\widehat{(\mathfrak{a}^n)} = (\hat{\mathfrak{a}})^n$;
- (iii) $\hat{\mathfrak{a}}^n / \hat{\mathfrak{a}}^{n+1} \cong \mathfrak{a}^n / \mathfrak{a}^{n+1}$;
- (iv) $\hat{\mathfrak{a}}$ is contained in the Jacobson radical of $\hat{A}$.

Proposition 10.16. Let A be a Noetherian local ring, $\mathfrak{m}$ its maximal ideal. Then the $\mathfrak{m}$ -adic completion $\hat{A}$ of A is a local ring with maximal ideal $\hat{\mathfrak{m}}$.

Proposition 10.17 (Kull's theorem). Let A be a Noetherian ring, $\mathfrak{a}$ an ideal, M a finitely-generated A -module and $\hat{M}$ the $\mathfrak{a}$ -adic completion of M . Then the kernel $E = \bigcap_{n=1}^{\infty} \mathfrak{a}^n M$ of $M \rightarrow \hat{M}$ consists of those $x \in M$ annihilated by some element of $1 + \mathfrak{a}$.

THE ASSOCIATED GRADED RING

Let A be a ring and $\mathfrak{a}$ an ideal of A . The **associated graded ring** is the graded ring

$$G(A) (= G_{\mathfrak{a}}(A)) = \bigoplus_{n=0}^{\infty} \mathfrak{a}^n / \mathfrak{a}^{n+1} \quad (a^0 = A).$$

Similarly, if M is an A -module and (M_n) is an $\mathfrak{a}$ -filtration of M , then the **associated graded module** is

$$G(M) = \bigoplus_{n=0}^{\infty} M_n / M_{n+1},$$

which is a graded $G(A)$ -module. Let $G_n(M) = M_n / M_{n+1}$.

Proposition 10.22. *Let A be a Noetherian ring, $\mathfrak{a}$ an ideal of A . Then*

- (i) $G_{\mathfrak{a}}(A)$ is Noetherian;
- (ii) $G_{\mathfrak{a}}(A)$ and $G_{\hat{\mathfrak{a}}}(\hat{A})$ are isomorphic as graded rings;
- (iii) If M is a finitely-generated A -module and (M_n) is a stable $\mathfrak{a}$ -filtration of M , then $G(M)$ is a finitely-generated graded $G_{\mathfrak{a}}(A)$ -module.

Proposition 10.23. *Let $\phi : A \rightarrow B$ be a homomorphism of filtered groups, i.e. $\phi(A_n) \subseteq B_n$, and let $G(\phi) : G(A) \rightarrow G(B)$, $\hat{\phi} : \hat{A} \rightarrow \hat{B}$ be the induced homomorphisms of the associated graded and completed groups. Then*

- (i) $G(\phi)$ injective $\Rightarrow \hat{\phi}$ injective;
- (ii) $G(\phi)$ surjective $\Rightarrow \hat{\phi}$ surjective.

Proposition 10.26. *If A is a Noetherian ring, $\mathfrak{a}$ an ideal of A , then the $\mathfrak{a}$ -adic completion $\hat{A}$ of A is Noetherian.*

11 Dimension Theory

HILBERT FUNCTIONS

Let A be a Noetherian graded ring and M be a finitely-generated graded A -module. Let λ be an additive function on the class of all finitely generated A_0 -modules. The **Poincaré series** of M is the power series

$$P(M, t) = \sum_{n=0}^{\infty} \lambda(M_n) t^n \in \mathbb{Z}[[t]].$$

Theorem 11.1 (Hilbert-Serre theorem). $P(M, t)$ is a rational function in t of the form $f(t) / \prod_{i=1}^d (1 - t^{k_i})$, where $f(t) \in \mathbb{Z}[[t]]$.

The order of the pole of $P(M, t)$ at $t = 1$ is denoted as $d(M)$.

Corollary 11.2. If each $k_i = 1$, then for all sufficient large n , $\lambda(M_n)$ is a polynomial in n . More specifically, if $f(t) = \sum_{k=0}^N a_k t^k$, we have

$$\lambda(M_n) = \sum_{k=0}^n a_k \binom{d+n-k-1}{d-1}.$$

This is called **Hilbert function** or **Hilbert polynomial** of M , and its leading term is $(\sum a_k) n^{d-1} / (d-1)!$.

Proposition 11.3. If $x \in A_k$ is not a zero-divisor in M then $d(M/xM) = d(M) - 1$.

Proposition 11.4. Let A be a Noetherian local ring, $\mathfrak{m}$ its maximal ideal, $\mathfrak{q}$ an $\mathfrak{m}$ -primary ideal, M a finitely-generated A -module, (M_n) a stable $\mathfrak{q}$ -filtration of M . Then

- (i) M/M_n is of finite length for each $n \geq 0$;
- (ii) for all sufficiently large n this length is a polynomial $\chi_{\mathfrak{q}}^M(n)$ of degree $\leq s$, where s is the least number of generators of $\mathfrak{q}$;
- (iii) the degree and leading coefficient of $\chi_{\mathfrak{q}}^M(n)$ depend only on M and $\mathfrak{q}$, not on the filtration chosen.

If $M = A$, the polynomial $\chi_{\mathfrak{q}}^M(n)$ is called the **characteristic polynomial** of the $\mathfrak{m}$ -primary ideal $\mathfrak{q}$, simply noted as $\chi_{\mathfrak{q}}(n)$.

Proposition 11.6. $\deg \chi_{\mathfrak{q}}(n) = \deg \chi_{\mathfrak{m}}(n)$.

DIMENSION THEORY OF NOETHERIAN LOCAL RINGS

Theorem 11.14 (Dimension theorem). For any Noetherian local ring A the following three integers are equal:

- (i) the maximum length of chains of prime ideals in A (**Krull dimension**);
- (ii) the degree of the characteristic polynomial $\chi_{\mathfrak{m}} = l(A/\mathfrak{m}^n)$;
- (iii) the least number of generators of an $\mathfrak{m}$ -primary ideal of A .

Corollary 11.17 (Krull's principal ideal theorem). Let A be a Noetherian ring and let x be an element of A which is neither a zero-divisor nor a unit. Then every minimal prime ideal $\mathfrak{p}$ of (x) has height 1.

REGULAR LOCAL RINGS

Theorem 11.22. *Let A be a Noetherian local ring of dimension d , $\mathfrak{m}$ its maximal ideal, $k = A/\mathfrak{m}$. Then the following are equivalent:*

- (i) $G_{\mathfrak{m}}(A) \cong k[t_1, \dots, t_d]$ where the t_i are independent indeterminates;
- (ii) $\dim_k(\mathfrak{m}/\mathfrak{m}^2) = d$;
- (iii) $\mathfrak{m}$ can be generated by d elements.

The ring satisfying any of the conditions is called *regular* local ring.

Proposition 11.23. *Let A be a ring, $\mathfrak{a}$ an ideal of A such that $\bigcap_n \mathfrak{a}^n = 0$. Suppose that $G_{\mathfrak{a}}(A)$ is an integral domain. Then A is an integral domain.*

Proposition 11.24. *Let A be a Noetherian local ring. Then A is regular iff $\hat{A}$ is regular.*

TRANSCENDENTAL DIMENSION

Assume that k is an algebraically closed field and let V be an irreducible affine variety over k . Let the coordinate ring of V be $A(V) = k[x_1, \dots, x_n]/I(V)$. Then $k(V) = \text{Frac } A(V)$ is a finitely generated extension of k and so has a finite transcendence degree over k , which is defined to be the *dimension* of V .

If P is a point with maximal ideal $\mathfrak{m}$ we call the Krull dimension $\dim A(V)_{\mathfrak{m}}$ the *local dimension* of V at P .

Theorem 11.25. *For any irreducible variety V over k the local dimension at any point is equal to its dimension.*

EXERCISES

1 Rings and Ideals

1.1. Consider $(1+x)(1-x+x^2-\cdots+(-x)^{n-1})$.

1.2. (i) Passing to $(A/\mathfrak{p})[x]$. $A/\mathfrak{p}$ is an integral domain, so $\deg(\overline{f\bar{g}}) = \deg(\overline{f}) + \deg(\overline{g})$. $\overline{f\bar{g}} = 1$ means $a_1, \dots, a_n \in \mathfrak{p}$. Use [Proposition 1.8](#).

(ii) a, b are nilpotents $\Rightarrow a+b$ is nilpotent. Note that $f-a_0 = xf_1$ is nilpotent.

(iii) Let $gf=0$ and g is of minimal degree. The leading coefficient $a_nb_m=0$, so a_ng has less degree than g and so $a_ng=0$. Same process to the second leading and following coefficients, we have $a_i g=0$ for all i .

(iv) (Gauss's lemma) Let $\mathfrak{a}, \mathfrak{b}$ and $\mathfrak{c}$ are generated by the coefficients of f, g and fg , respectively. We need to proof $\mathfrak{c} = (1) \Leftrightarrow \mathfrak{a}\mathfrak{b} = (1)$. $\mathfrak{c} \subseteq \mathfrak{a}\mathfrak{b}$ by the form of coefficients in fg . If $\mathfrak{c} \subsetneq \mathfrak{a}\mathfrak{b} = (1)$, let $\mathfrak{m}$ be the maximal ideal containing $\mathfrak{c}$, then $(A/\mathfrak{m})[x]$ is an integer domain. $\overline{f\bar{g}} = 0$ but $\overline{f}, \overline{g} \neq 0$.

1.3. Induction on the indeterminates.

1.4. Pick the y in [Proposition 1.9](#) to be the indeterminate, then use [Exercise 1.2\(i\)](#).

1.5. (i) $\Leftarrow$: $fg = \sum_{i=0}^{\infty} \left(\sum_{j=0}^i a_j b_{i-j} \right) x^i$, so we can find g such that $\sum_{j=0}^i a_j b_{i-j} = 0$.

(ii) Similar to [Exercise 1.2\(ii\)](#). Construct a counterexample where a_n have increasing nilpotencies.

(iii) Pick the y in [Proposition 1.9](#) to be 1. Use (i) on $1-f \Leftrightarrow 1-a_0$.

(iv),(v) $A \cong A[[x]]/(x)$. Use [Proposition 1.1](#).

1.6. $\mathfrak{J}$ is an ideal, so there is an idempotent $e \in \mathfrak{J}$ but $e \notin \mathfrak{R}$. Pick the y in [Proposition 1.9](#) to be 1.

1.7. Note that $x(1-x^{n-1}) = 0 \in \mathfrak{p}$.

1.8. Zorn's lemma but define the partial order reversely.

1.9. $\Rightarrow$: Apply [Proposition 1.8](#) to $A/\mathfrak{a}$.

$\Leftarrow$: Use [Proposition 1.11\(ii\)](#).

1.10. Use [Corollary 1.5](#) and [Corollary 1.6\(i\)](#)

1.11. (i) $x+x = (x+x)^2 = x^2+2x+x^2 = x+2x+x$.

(ii) Proof that every element in $A/\mathfrak{p}$ is 1.

(iii) $x+y-xy$.

1.12. Use [Corollary 1.5](#).

1.13. **1 is a monic polynomial but it is neither irreducible nor reducible.** A is an integral domain, so $\deg(fg) = \deg(f) + \deg(g)$. If $\mathfrak{a} = (1)$, apply the equality to $1 = \sum_f y_f f(x_f)$.

1.14. If $xy \in \mathfrak{m}$, xy is zero-divisor, so are x and y . By maximality, $\mathfrak{m} \subseteq \mathfrak{m} + (x) = \mathfrak{m}$.

1.15. (i) Apply [Proposition 1.8](#) to $A/\mathfrak{a}$ to proof the second equality.

(iv) Use $\sqrt{\mathfrak{a} \cap \mathfrak{b}} = \sqrt{\mathfrak{a}\mathfrak{b}}$.

1.16. Irreducible polynomials relate to prime ideals.

1.17. (i)-(iv) Use [Exercise 1.15](#).

(v) Let $(X_{f_i})_{i \in I}$ be an open cover. By [Exercise 1.15](#), $V(\{f_i\}_{i \in I}) = \emptyset$, so $\{f_i\}_{i \in I}$ generates the unit ideal. Let $1 = \sum_{i \in J} g_i f_i$ where J is a finite subset of I . If $\mathfrak{p} \notin \bigcup_{i \in J} X_{f_i}$, then $f_i \in \mathfrak{p}$ for all $i \in J$.

EXERCISES

(vi) Similar to (v), where the unit ideal shall change to the radical $\sqrt{(f)}$.

1.18. (iv) If $V(\mathfrak{p}_x) \subseteq V(\mathfrak{p}_y)$, $\mathfrak{p}_x \in \overline{V(\mathfrak{p}_y)}$. If they don't have containing relation, $\mathfrak{p}_x \in \overline{V(\mathfrak{p}_y)}$ and $\mathfrak{p}_y \in \overline{V(\mathfrak{p}_x)}$.

1.19. Use the closed sets definition. Consider minimal prime ideals.

1.20. (i) Let $\overline{Y} = Z_1 \cup Z_2$, then $Z_i \cap Y = \emptyset$ for an $i \in \{1, 2\}$. Thus $Z_i = \overline{Z_i \cap Y} = \emptyset$.

(ii) Similar to (i). Construct the maximal element and proof that it is irreducible.

(iii) If Y Hausdorff but irreducible, every pair of open sets in Y intersect. Thus Y is singleton.

(iv) Use the closed set definition.

1.21. Simplify the sets: $\phi^{*-1}(E) = \{\mathfrak{q} \in Y : \phi^{-1}(\mathfrak{q}) \in E\}$, $\phi^*(V(E)) = \{\phi^{-1}(\mathfrak{q}) : E \subseteq \mathfrak{q}\}$.

1.22. Use quotient to find the form of prime ideals in A .

1.23. (i) $X_f = V(1 - f)$.

(ii) $X_{f_1} \cup X_{f_2} = X_{f_1 + f_2 - f_1 f_2}$.

(iii) A closed set in compact space is compact. By [Exercise 1.17](#), it is a finite union of basic open set. Then apply (ii).

(iv) By [Exercise 1.11](#) every prime ideal is maximal. Pick $f \in \mathfrak{p}_x \setminus \mathfrak{p}_y$, then X_f and $V(f)$ separate two points.

1.24. $a = a \wedge 1 = a \wedge (a \vee a') = (a \wedge a) \vee (a \wedge a') = (a \wedge a) \vee 0 = a \wedge a$.

1.25. Use [Exercise 1.23\(iv\)](#).

1.26. $\mathfrak{m}_x \in \mu(U_f) \Leftrightarrow f(x) \neq 0 \Leftrightarrow f \notin \mathfrak{m}_x \Leftrightarrow \mathfrak{m}_x \in \tilde{U}_f$.

1.27. If $x \neq y$, there exists a polynomial $f \in P(X)$ such that $f(x) = 0$ but $f(y) \neq 0$, so $\mathfrak{m}_x \neq \mathfrak{m}_y$, and so μ injective. By Nullstellensatz, every maximal ideal in $P(X)$ is of the form $\mathfrak{m} = (t_1 - a_1, \dots, t_n - a_n)$, with $(a_1, \dots, a_n) \in X$, so μ surjective. By [Exercise 1.26](#), μ is a homeomorphism.

1.28. Project to coordinate functions.

2 Modules

2.1. Bézout's identity.

2.2. Tensor the exact sequence $0 \rightarrow \mathfrak{a} \rightarrow A \rightarrow A/\mathfrak{a} \rightarrow 0$. Calculate the image of $\mathfrak{a} \otimes M \rightarrow A \otimes M$.

2.3. Consider the residue field k of A and passing $M \otimes_A N$ to vector space $M_k \otimes_k N_k$. Use [Exercise 2.2](#) and [Nakayama's lemma](#).

2.4. There exist canonical projections.

2.5. The definition of flatness only considers the module structure. Use [Exercise 2.4](#).

2.6. Construct two homomorphisms.

2.7. Proof that $(A/\mathfrak{p})[x]$ and $A[x]/\mathfrak{p}[x]$ are isomorphic.

2.8. (ii) Use [Proposition 2.15](#).

2.9. Pick $y_1, \dots, y_m \in M$ whose images are the generators of M'' . Use diagram chasing for an element of M .

2.10. $u(m) + \mathfrak{a}N \in N$, then use surjectiveness. Apply [Nakayama's lemma](#) on $(u(M) + \mathfrak{a}N)/N$.

2.11. (i),(ii) Consider a residue field and passing to vector spaces. Use the right exactness of tensor product.

(iii) If $m > n$, let $\psi : A^m \xrightarrow{\phi} A^n \hookrightarrow A^m$. By [Proposition 2.4](#), ψ satisfies $\psi^n + a_{n-1}\psi^{n-1} + \cdots + a_0 = 0$. Pick such equation to have the minimum degree, then ψ injective implies $a_0 \neq 0$. Apply the equation to element $(0, \dots, 0, 1)$ to obtain contradiction.

2.12. Choose $u_i \in M$ such that $\phi(u_i)$ are a basis of A^n , then proof that $0 \rightarrow \ker(\phi) \rightarrow M \xrightarrow{\phi} A^n \rightarrow 0$ splits.

2.13. $p \circ g = \text{id}_N$ injective $\Rightarrow g$ injective, so $0 \rightarrow N \xrightarrow{g} N_B \rightarrow N_B/g(N) \rightarrow 0$ splits.

2.14. This question asks to proof $\mu_i = \mu_j \circ \mu_{ij}$.

2.15. (i) Use [Exercise 2.14](#) to increase indices.

(ii) $x_i = \sum_k (y_k - \mu_{kl}(y_k))$ is an equation in $\bigoplus M_i$, so μ_{ij} cannot be applied directly. Use canonical projection $\pi_i : \bigoplus M_i \rightarrow M_i$ on the equation first.

2.16 (Universal property of direct limit). Use [Exercise 2.15\(i\)](#) for both existence and uniqueness.

2.17. Similar to [Exercise 2.15\(i\)](#), move components into a same index.

2.18. Use [Exercise 2.16](#).

2.19 ($\varinjlim$ is an exact functor). Diagram chasing. Use [Exercise 2.15](#).

2.20. Draw the diagram. Define two homomorphisms by the universal properties of tensor product and direct limit. Verify their compositions are identity maps.

2.21. Use [Exercise 2.15\(ii\)](#) on $\alpha_i(1)$.

2.22. Use [Exercise 2.15](#). Pick a nilpotent or assume $xy = 0$.

2.23. If $J \subseteq J'$, $B_J \cong B_J \otimes (A)_{i \in J' \setminus J}$. Let Γ be the set of all finite subsets of Λ , then $(B_J)_{J \in \Gamma}$ is the case of [Exercise 2.21](#). We can also use [Exercise 2.17](#) to see its form.

2.24. Directly from the definition of Tor functor.

2.25. Use [Snake lemma](#).

2.26. Let M_n be generated by $\{x_1, \dots, x_n\}$. We have an exact sequence

$$\text{Tor}_1(M_{n-1}, N) \rightarrow \text{Tor}_1(M_n, N) \rightarrow \text{Tor}_1(M_n/M_{n-1}, N).$$

By [Proposition 2.3](#), $M_n/M_{n-1} \cong A/\mathfrak{a}$.

Let $f : \mathfrak{a} \otimes N \rightarrow A \otimes N$, $f(\sum a_i \otimes y_i) = 0$. Let $\mathfrak{a}'$ be generated by a_i . By assumption, $f|_{\mathfrak{a}' \otimes N}$ is injective, so $\sum a_i \otimes y_i = 0$ and thus $\text{Tor}_1(A/\mathfrak{a}, N) = 0$. Since $M_0 = A$ and $\text{Tor}_1(A, N) = 0$, the exact sequence implies $\text{Tor}_1(M_n, N) = 0$ for all n . Then use [Proposition 2.19\(iv\)](#). **Another approach is to use the fact that any module is the direct limit of some finitely generated modules, and direct limit is an exact functor.**

2.27. (i) $\Rightarrow$ (ii) $A/(x)$ is a flat A -module, then $\phi : A/(x) \otimes (x) \rightarrow A/(x)$ injective. $\phi(1 \otimes x) = x = 0$, so $A/(x) \otimes (x) = 0$. Then use [Exercise 2.2](#).

(ii) $\Rightarrow$ (iii) Let $x = ax^2$, then ax is an idempotent. $(x) \ni bx = bax^2 = (bx)(ax) \in (ax)$, so $(ax) = (x)$. Similar to [Exercise 1.11\(iii\)](#), every finitely generated ideal is generated by an idempotent e , so $A = (e) \oplus (1 - e)$.

(iii) $\Rightarrow$ (i) $N \otimes_A A = (N \otimes_A \mathfrak{a}) \oplus (N \otimes_A \mathfrak{a}')$, so $(N \otimes_A \mathfrak{a}) \rightarrow N \otimes_A A$ injective. Use [Exercise 2.26](#).

2.28. Use [Exercise 2.27\(ii\)](#).

3 Rings and Modules of Fractions

3.1. Element-wise. Only if part requires the finitely generated property.

3.2. Element-wise. Hint.

3.3. Element-wise.

3.4. Element-wise.

3.5. Element-wise. Counter example is Boolean ring.

3.6. Use the fact that for any multiplicatively closed set S , if $A - S$ is an ideal, it's automatically prime.

3.7. (i),(ii) More specifically, $A - S = \bigcup_{\mathfrak{p} \cap S = \emptyset} \mathfrak{p}$.

(iii) $\mathfrak{p} \cap (1 + \mathfrak{a}) = \emptyset \Leftrightarrow \mathfrak{a} + \mathfrak{p} \neq (1) \Leftrightarrow \mathfrak{a} + \mathfrak{p} \subseteq \mathfrak{m}$ for some maximal ideal $\mathfrak{m}$. Note that the union of a chain of prime ideals is only a maximum ideal.

3.8. Element-wise. (v) $\Rightarrow$ (ii): If $t/1$ is not a unit in $S^{-1}A$, it contains in a prime ideal $\mathfrak{q} \subseteq S^{-1}A$, so $t \in T \cap \mathfrak{q}^c$.

3.9. Use [Exercise 3.6](#). Element-wise.

3.10. (i) [Exercise 2.27](#) says for any element $x \in A$, $x = ax^2$ for some $a \in A$. (ii) $\Rightarrow$: Use [Exercise 2.28](#). $\Leftarrow$: Consider an $A_{\mathfrak{m}}$ -module. Use [Proposition 3.10](#).

3.11. Let A be a ring. Then the following are equivalent:

(i) $A/\mathfrak{R}$ is absolutely flat.

(ii) Every prime ideal of A is maximal.

(iv) $\text{Spec}(A)$ is Hausdorff.

(i) $\Rightarrow$ (ii): Use [Exercise 2.27](#) on $A/\mathfrak{p}$. (ii) $\Rightarrow$ (i): Proof that $(A/\mathfrak{R})_{\overline{\mathfrak{p}}}$ has no nilpotent, then use [Exercises 1.10](#) and [3.10](#). (iv) $\Rightarrow$ (ii): There exist $f_1 \in \mathfrak{p}_1$ and $f_2 \in \mathfrak{p}_2$ such that $f_1 + f_2 = 1$, so $f_2 \in X_{f_1}$ and $f_1 \in X_{f_2}$. **need to finish the proof**

3.12. Element-wise. (iv) Use [Proposition 3.5](#).

3.13. Element-wise.

3.14. Hint. Use [Proposition 3.8](#).

3.15. Hint.

3.16. (ii) $\Rightarrow$ (iii): [Proposition 1.17](#). (iii) $\Rightarrow$ (iv) $\Rightarrow$ (v): Use hint. (v). Take $M = A/\mathfrak{a}$. Then $M_B = B/\mathfrak{a}^e$.

3.17.

3.18. Use [Exercise 3.3](#) to proof $B_{\mathfrak{q}} \cong (B_{\mathfrak{p}})_{\mathfrak{q}B_{\mathfrak{p}}}$. By [Proposition 3.3](#) and [Proposition 3.5](#), $B_{\mathfrak{q}}$ is flat over $B_{\mathfrak{p}}$.

3.19. (v) $M \cong \bigoplus_{i=1}^n A/\mathfrak{a}_i$, so $\text{Supp}(M) = \bigcup_{i=1}^n \text{Supp}(A/\mathfrak{a}_i) = \bigcup_{i=1}^n V(\mathfrak{a}_i) = V(\bigcap_{i=1}^n \mathfrak{a}_i)$. Proof that $\bigcap_{i=1}^n \mathfrak{a}_i = \text{Ann}(M)$.

(vii) Use [Exercise 2.2](#).

(viii)

3.20. Counter example is $A[x]/(x^2)$.

3.21. (i)-(iii) Use one-to-one order preserving correspondences. (iv) $A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} = (A/\mathfrak{p})_{\mathfrak{p}}$.

3.22. Element-wise.

3.23. (i) Use [Exercise 1.17](#). (v) Define two homomorphisms by the universal properties of direct limit and local ring.

3.24.

4 Primary Decomposition

4.1. Use [Exercise 3.21](#) so $\text{Spec}(A/\mathfrak{a}) = V(\mathfrak{a})$.

4.2. Omitted.

4.3. For any element $x \in A$, $x(1 - ax) = 0 \in \mathfrak{q}$.

4.4. Omitted.

4.5. Omitted.

4.6.

4.7. Element-wise.

4.8. Use [Exercise 4.7](#).

4.9. (i) $\Leftarrow$: x is contained in all prime ideal containing $(0 : \mathfrak{a})$, so $x \in \sqrt{(0 : \mathfrak{a})}$. (iii) We have a relation $(0 : \mathfrak{a}) \subseteq \bigcap \sqrt{(\mathfrak{q}_i : \mathfrak{a})} \subseteq \mathfrak{p}$. Then use minimality to proof that $\mathfrak{p}$ is an associated prime ideal.

4.10. (iv) doesn't hold.

4.11. Note that $x \in S(\mathfrak{a}) \Leftrightarrow$ there exists an $s \in S$ such that $sx \in \mathfrak{a}$. Show that $S_{\mathfrak{p}}(0)$ is contained in all $\mathfrak{p}$ -primary ideals.

If $0 = \bigcap \mathfrak{q}_i$ is minimal, $\sqrt{(0)}$ is the intersection of minimal prime ideals.

4.12. Use [Proposition 3.11](#). Use [Proposition 4.9](#) to show there are at most 2^n different $S(\mathfrak{a})$.

4.13. (i) Use [Exercise 4.12](#). (ii) Use [Proposition 4.8](#) and [Proposition 4.9](#). (iii) Proof that $S_{\mathfrak{p}}(\mathfrak{p}^{(m)}\mathfrak{p}^{(n)}) = \mathfrak{p}^{(m+n)}$. (iv) $\Leftarrow$: Show that $\mathfrak{p}^{(n)}$ is the smallest $\mathfrak{p}$ -primary ideal containing $\mathfrak{p}^n$.

4.14. Let $x \notin \mathfrak{q}$. If $ab \in (\mathfrak{q} : x)$ but $a \notin (\mathfrak{q} : x)$, we have $xab \in \mathfrak{q}$ but $xa \notin \mathfrak{q}$. Thus, $b \in (\mathfrak{q} : xa) = (\mathfrak{q} : x)$ by maximality, and therefore $(\mathfrak{q} : x)$ is prime.

4.15.

4.16. Use [Proposition 3.11](#) and [Proposition 4.9](#).

4.17. Use hint.

4.18. (i) $\Rightarrow$ (L1): By [Proposition 4.9](#), $S_{\mathfrak{p}}(\mathfrak{a}) = \bigcap \mathfrak{q}_i$. Pick $x \in S_{\mathfrak{p}}$, then $\mathfrak{q}_i = (\mathfrak{q}_i : x)$ by [Proposition 4.4](#). (i) $\Rightarrow$ (L2): Use [Exercise 4.12](#).

4.19. Use hint.

5 Integral Dependence and Valuations

5.1. $f^*(V(I)) = \{f^{-1}(\mathfrak{q}) : \mathfrak{q} \supseteq I\}$. Use [Theorem 5.10](#) to proof $f^*(V(I)) \supseteq V(f^{-1}(I))$.

5.2.

5.3. Use the operator $- \otimes_A c^n$ to get the result.

5.4. If $\left(\frac{1}{x+1}\right)^n + \sum \frac{f_i(x^2-1)}{g_i(x^2-1)(x+1)^i} = 0$, multiply both side by $(x+1)^n$ and take $x = -1$ to get contradiction.

5.5. (i) Element-wise. (ii) Use [Theorem 5.10](#) to proof that the contraction of maximal ideals in B are maximal in A

5.6. Let $b_i \in B_i$, $f_i \in A[x]$ monic, $f_i(b_i) = 0$. Let $f = f_1 \cdots f_n$, then $f(b_1, \dots, b_n) = 0$.

EXERCISES

5.7. Let $x \in B$ and x integral over A . Let $x^n + a_{n-1}x^{n-1} + \cdots + a_0 = 0$ is the equation of the least degree. Analyze $x(x^{n-1} + a_{n-2}x^{n-2} + \cdots + a_1) = -a_0$.

5.8. (ii) Proof that there is a ring $D \supseteq B$ such that f factors into linear factors in D . Use Kronecker's construction.

5.9. Hint.

5.10. (a) $\Rightarrow$ (b): Let $f^*(V(\mathfrak{q}_1)) = V(I)$, so $I \subseteq \mathfrak{p}_1 \subseteq \mathfrak{p}_2$, and we have $\mathfrak{p}_2 \in V(I) = f^*(V(\mathfrak{q}_1))$. (a') $\Rightarrow$ (b'): Hint.

5.11. Use [Exercises 3.18](#) and [5.10](#).

5.12. Note that G has identity element so x is a root of the polynomial $\prod_{\sigma \in G} (t - \sigma(x))$. The coefficients are symmetric polynomials of σ_i . We don't know how G acts on $(S^{-1}A)^G$, so we define $\sigma(x/s) = \sigma(x)/\sigma(s)$.

5.13.

5.14. $\sigma \in G$ fixes A .

5.15.

5.16.

5.17. Hint. Using Zariski's lemma is the easiest way.

5.18 ([Zariski's lemma](#)).

5.19. Omitted.

5.20. Hint.

5.21. Hint.

5.22. Hint.

5.23. (i) $\Rightarrow$ (ii): Note that $f : A \rightarrow A/\ker f$. Use [Proposition 1.1](#). (ii) $\Rightarrow$ (iii): Consider $A/\mathfrak{p}$. (iii) $\Rightarrow$ (i): If there is a prime ideal $\mathfrak{p}$ which is not an intersection of maximal ideals, the Jacobson radical $\mathfrak{J}$ of $A/\mathfrak{p}$ is not zero. Pick $f \in \mathfrak{J}$, then $(A/\mathfrak{p})_f \neq 0$. The contraction of a maximal ideal of $(A/\mathfrak{p})_f$ is maximal among all prime ideals which doesn't contain f , but by (iii), the intersection has to contain f .

5.24. (i) Use [Exercise 5.23](#)(iii) and going-up theorem. (ii) Consider $B/\mathfrak{q}$ and $A/\mathfrak{q}^c$. Then use [Exercise 5.22](#), [Exercise 5.23](#)(i) and [Proposition 1.1](#).

5.25. Hint.

5.26. (2) $\Rightarrow$ (3): $U_1 \cap X_0 = U_2 \cap X_0 \Rightarrow U_1^c \cap X_0 = U_2^c \cap X_0$, then take closure. (i) $\Rightarrow$ (ii) and (ii) $\Rightarrow$ (iii): Consider $V(\mathfrak{a}) \cap X_f \cap \text{Max}(A)$. If all maximal ideals in $V(\mathfrak{a})$ contains f , $V(\mathfrak{a}) \cap X_f = \emptyset$ by [Exercise 5.23](#)(iii). (iii) $\Rightarrow$ (i): If not Jacobson, there is a prime ideal $\mathfrak{p}$ such that $\mathfrak{p} \subsetneq \bigcap_{\mathfrak{q} \supseteq \mathfrak{p}} \mathfrak{q}$. Let $f \in \bigcap_{\mathfrak{q} \supseteq \mathfrak{p}} \mathfrak{q} \setminus \mathfrak{p}$, then $X_f \cap V(\mathfrak{p}) = \{\mathfrak{p}\}$.

5.27. Use [Theorem 5.21](#) to proof maximal elements are valuation rings. Conversely, if $(A, \mathfrak{m}) \subsetneq (A', \mathfrak{m}')$, pick $x \in A' \setminus A$. Then $x^{-1} \in \mathfrak{m}$ because it is not a unit in A .

5.28. (1) $\Rightarrow$ (2): Pick $a \in \mathfrak{a} \setminus \mathfrak{b}$, and any $b \in \mathfrak{b}$. Consider a/b and b/a . (2) $\Rightarrow$ (1): Consider $(a) \subseteq (b)$ or $(b) \subseteq (a)$, we have $a = xb$ or $b = ya$, so $y = x^{-1}$.

5.29.

6 Chain Conditions

- 6.1.** (i) Let $x \in \ker(u)$. Since u^n surjective, there is a $y \in M$ such that $u^n(y) = x$. But $u^{n+1}(y) = u(x) = 0$, means that $y \in \ker(u^{n+1}) = \ker(u^n)$.
(ii) Let $x \in M$. Then $u^n(x) \in \operatorname{im}(u^n) = \operatorname{im}(u^{n+1})$, so there is a $y \in M$ such that $u^{n+1}(y) = u^n(x)$. But u^n injective, means that $u(y) = x$.
- 6.2.** If M is not Noetherian, there is a submodule N which is not finitely generated. Construct a chain of finitely generated submodules by the generators of N .
- 6.3.** Consider $0 \rightarrow N_2 / (N_1 \cap N_2) \rightarrow M / (N_1 \cap N_2) \rightarrow M / N_2 \rightarrow 0$. Use [Proposition 2.1](#).
- 6.4.** Let $m_1, \dots, m_n$ be generators of M , then $\mathfrak{a} = \bigcap \operatorname{Ann}(m_i)$. $A / \operatorname{Ann}(m_i) \cong Am_i \subseteq M$, then use [Exercise 6.3](#).
- 6.5.** Use the open set definition.
- 6.6.** (i) $\Rightarrow$ (iii) Use [Exercise 6.5](#).
(ii) $\Rightarrow$ (i) Consider a chain of ascending open sets in X .
- 6.7.** Let Σ be the collection of closed subsets of X which are not finite union of irreducible closed subspaces. Then Σ has minimal element X_0 . Proof that X_0 cannot be either irreducible or reducible.
- 6.8.** The converse isn't true. The counter example is $k[x_1, \dots] / (x_1^2, \dots)$.
- 6.9.** The irreducible components of $\operatorname{Spec}(A)$ is $V(\mathfrak{p})$ where $\mathfrak{p}$ minimal ([Exercise 1.20\(iv\)](#)).
- 6.10.** Use [Exercise 3.19\(v\)](#).
- 6.11.** By [Exercise 6.7](#), $V(\mathfrak{b})$ is a finite union of $V(\mathfrak{q}_i)$, where $\mathfrak{q}_i$ minimal and containing $\mathfrak{b}$. Then use [Exercise 5.10\(c\)](#).
- 6.12.** The converse isn't true. The counter example is an infinite product $\prod \mathbb{Z}/2\mathbb{Z}$.

7 Noetherian Rings

- 7.1.**
- 7.2.**
- 7.3.** (i) $\Rightarrow$ (ii): Use [Proposition 4.8](#) and [Proposition 4.4](#). (ii) $\Rightarrow$ (iii): Let $S = \{1, x, x^2, \dots\}$. Proof that $\bigcup (\mathfrak{a} : x^n) = S(\mathfrak{a})$. (iii) $\Rightarrow$ (i): Similar to the proof of [Proposition 7.12](#).
- 7.4.**
- 7.5.** Use [Exercise 5.12](#) and [Proposition 7.8](#).
- 7.6.** Hint.
- 7.7.** Use a.c.c..
- 7.8.** $A \cong A[x]/(x)$.
- 7.9.** Hint.
- 7.10.** $M[x] = M \otimes A[x]$, so it is a homomorphic image of $M \times A[x]$.
- 7.11.** Counter example is an infinite product $\prod \mathbb{Z}/2\mathbb{Z}$.
- 7.12.** By [Exercise 3.16\(i\)](#), $\mathfrak{a} \mapsto \mathfrak{a}^e$ is injective. Then use a.c.c..
- 7.13.** Transfer the Noetherian property $B \rightarrow B_{\mathfrak{p}} \rightarrow B_{\mathfrak{p}}/\mathfrak{p}B_{\mathfrak{p}}$.

7.14 (Hilbert's Nullstellensatz). The hint uses a method other than Rabinowitsch trick.

7.15. Hint.

7.16. Use [Exercise 7.15](#).

7.17.

7.18.

7.19.

7.20. (i) $\Rightarrow$: Rigorous proof requires using induction on $\mathcal{F}_i$, where $\mathcal{F}_i$ is formed by applying complement i times on the topology $\tau = \mathcal{F}_0$, and satisfies $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \cdots \subseteq \mathcal{F}$. (ii) $\Rightarrow$: Use [Exercise 1.19](#).

7.21. $\Rightarrow$: Use [Exercise 7.20](#). $\Leftarrow$: Use hint. If $\overline{E \cap X_0} \neq X_0$, $E \cap \overline{E \cap X_0} \in \mathcal{F}$ by minimality, but $E \cap \overline{E \cap X_0} = E \cap X_0$. If $U_0 \subseteq E \cap X_0$, let $U_0 = U \cap X_0$ where U is open in X , then $E \cap X_0 = U_0 \cup (E \cap U^c \cap X_0)$.

7.22. If $U_0 \subseteq E \cap X_0$, use the notations similar to [Exercise 7.21](#). $U^c \cap X_0$ is closed, so by minimality $E \cap U^c \cap X_0$ is open.

7.23. Hint.

7.24. Hint.

7.25. Use [Exercises 5.11](#) and [7.24](#).

7.26.

7.27.

8 Artinian Rings

8.1. Use [Proposition 7.16](#) on local ring $A_{\mathfrak{p}_i}$. Then apply the contraction. If $\mathfrak{q}_i$ is isolated, then $\mathfrak{p}_i$ is minimal.

8.2. Omitted.

8.3.

9 Discrete Valuation Rings and Dedekind Domain

10 Completion

10.1. $B = \bigoplus_{n \geq 0} \mathbb{Z}/p^n \mathbb{Z}$, so $A_k = \alpha^{-1}(p^k B) = \bigoplus_{n \geq k} \mathbb{Z}/p \mathbb{Z}$. The sequence $0 \rightarrow A \rightarrow B \xrightarrow{p} B \rightarrow 0$ is exact but $\hat{A} \rightarrow \hat{B} \xrightarrow{\hat{p}} \hat{B} \rightarrow 0$ is not, due to the kernel of $\hat{p}$ is $\prod_{k \geq 0} \mathbb{Z}/p \mathbb{Z}$.

10.2. The explicit form of the given exact sequence is

$$0 \rightarrow \bigoplus_{k > n} \mathbb{Z}/p \mathbb{Z} \rightarrow \bigoplus_{n \geq 0} \mathbb{Z}/p \mathbb{Z} \rightarrow \bigoplus_{n=0}^k \mathbb{Z}/p \mathbb{Z} \rightarrow 0.$$

Then use [Exercise 10.1](#). Analyze definitions in the proof of [Proposition 10.2](#) to claim that $\text{coker } d^B = \varprojlim^1 A = 0$.

10.3.

11 Dimension Theory