

MATH.APP.790 : Topics in Mathematics, Nonlinear time series analysis

Narayan P. Subramaniam

Tampere University (MET Faculty)

January 2021

Lecture overview

- Estimation of embedding dimension
- Fractal dimensions
- Lyapunov exponents
- Entropy measures

Embedding dimension

- In order to perform phase space reconstruction from a time series, apart from τ , another important parameter to be estimated is the embedding dimension (k).

Embedding dimension

- In order to perform phase space reconstruction from a time series, apart from τ , another important parameter to be estimated is the embedding dimension (k).
- There are various methods in the literature on the selection of k and many of these methods are based on false-nearest neighbor (FNN) principle.

Embedding dimension

- In order to perform phase space reconstruction from a time series, apart from τ , another important parameter to be estimated is the embedding dimension (k).
- There are various methods in the literature on the selection of k and many of these methods are based on false-nearest neighbor (FNN) principle.
- The rationale : Examine if the points along a trajectory in dimension k are also neighbors in dimension $k + 1$.

Embedding dimension

- Due to projection, in dimensions lower than the actual dimension, many points on the trajectory will be close to each other (**false neighbors**).

Embedding dimension

- Due to projection, in dimensions lower than the actual dimension, many points on the trajectory will be close to each other (**false neighbors**).
- Starting with an initial dimension of $k = 1$, increment k by one and for each state vector x_i , the nearest neighbor is computed.

Embedding dimension

- Due to projection, in dimensions lower than the actual dimension, many points on the trajectory will be close to each other (**false neighbors**).
- Starting with an initial dimension of $k = 1$, increment k by one and for each state vector x_i , the nearest neighbor is computed.
- For every increase in k , the percentage of false neighbors is also computed.

Embedding dimension

- Due to projection, in dimensions lower than the actual dimension, many points on the trajectory will be close to each other (**false neighbors**).
- Starting with an initial dimension of $k = 1$, increment k by one and for each state vector x_i , the nearest neighbor is computed.
- For every increase in k , the percentage of false neighbors is also computed.
- The value of k at which the percentage of false neighbors becomes zero (or arbitrarily small due to the effect of noise) is considered as an appropriate choice for k .

Embedding dimension

Given a scalar time series and τ , one reconstructs the phase space vector at an initial dimension k . Thus we have a state vector at time instance t_n ,

$$x_n^k = (x_{t_n}, x_{t_n-\tau}, \dots, x_{t_n-(k-1)\tau})$$

which has a nearest neighbor

$$x_{\mathcal{N}(n)}^k = (x_{t_{\mathcal{N}(n)}}, x_{t_{\mathcal{N}(n)}-\tau}, \dots, x_{t_{\mathcal{N}(n)}-(k-1)\tau}).$$

Embedding dimension

Given a scalar time series and τ , one reconstructs the phase space vector at an initial dimension k . Thus we have a state vector at time instance t_n ,

$$x_n^k = (x_{t_n}, x_{t_n-\tau}, \dots, x_{t_n-(k-1)\tau})$$

which has a nearest neighbor

$$x_{\mathcal{N}(n)}^k = (x_{t_{\mathcal{N}(n)}}, x_{t_{\mathcal{N}(n)}-\tau}, \dots, x_{t_{\mathcal{N}(n)}-(k-1)\tau}).$$

If the above reconstruction occurred in an insufficient dimension k , then this closeness could be a result of trajectories crossing.

Embedding dimension

- Increase the dimension to $k + 1$.

Embedding dimension

- Increase the dimension to $k + 1$.
- Reconstruct the phase space and denote the reconstructed vector and its nearest neighbors as x_n^{k+1} and $x_{\mathcal{N}(n)}^{k+1}$ respectively.

Embedding dimension

- Increase the dimension to $k + 1$.
- Reconstruct the phase space and denote the reconstructed vector and its nearest neighbors as x_n^{k+1} and $x_{\mathcal{N}(n)}^{k+1}$ respectively.
- The increase in the distance from k to $k + 1$ can be given as,

Embedding dimension

- Increase the dimension to $k + 1$.
- Reconstruct the phase space and denote the reconstructed vector and its nearest neighbors as x_n^{k+1} and $x_{\mathcal{N}(n)}^{k+1}$ respectively.
- The increase in the distance from k to $k + 1$ can be given as,

$$D_n = ||x_n^{k+1} - x_{\mathcal{N}(n)}^{k+1}|| - ||x_n^k - x_{\mathcal{N}(n)}^k||, \quad (1)$$

Embedding dimension

- Increase the dimension to $k + 1$.
- Reconstruct the phase space and denote the reconstructed vector and its nearest neighbors as x_n^{k+1} and $x_{\mathcal{N}(n)}^{k+1}$ respectively.
- The increase in the distance from k to $k + 1$ can be given as,

$$D_n = ||x_n^{k+1} - x_{\mathcal{N}(n)}^{k+1}|| - ||x_n^k - x_{\mathcal{N}(n)}^k||, \quad (1)$$

which can be normalized and computed for all the points as

$$\xi = \sum_{n=1}^{N-k-1} \Theta \left(\frac{D_n}{||x_n^k - x_{\mathcal{N}(n)}^k||} - r \right). \quad (2)$$

Embedding dimension

- One can set a suitable threshold (r) to determine if x_n^k and $x_{\mathcal{N}(n)}^k$ are neighbors.

Embedding dimension

- One can set a suitable threshold (r) to determine if x_n^k and $x_{\mathcal{N}(n)}^k$ are neighbors.
- In other words, if ξ is greater than r then these two points can be called false neighbors.

Embedding dimension

- One can set a suitable threshold (r) to determine if x_n^k and $x_{\mathcal{N}(n)}^k$ are neighbors.
- In other words, if ξ is greater than r then these two points can be called false neighbors.
- Thus, ξ is the amount of false neighbors that one would find in dimension k .

Embedding dimension

- One can set a suitable threshold (r) to determine if x_n^k and $x_{\mathcal{N}(n)}^k$ are neighbors.
- In other words, if ξ is greater than r then these two points can be called false neighbors.
- Thus, ξ is the amount of false neighbors that one would find in dimension k .
- The optimal embedding dimension is then defined as the dimension for which ξ becomes zero (or very small in case of noisy time series).

Embedding dimension

Hegger and Kantz proposed a modification to the FNN method by excluding points whose initial distance was already greater σ/r , where σ is the standard deviation of the data. The modified FNN method is given by

$$\xi_{mod} = \frac{\sum_{n=1}^{N-k-1} \Theta\left(\frac{\|x_n^{k+1} - x_{\mathcal{N}(n)}^{k+1}\|}{\|x_n^k - x_{\mathcal{N}(n)}^k\|} - r\right) \Theta\left(\frac{\sigma}{r} - \|x_n^k - x_{\mathcal{N}(n)}^k\|\right)}{\sum_{n=1}^{N-k-1} \Theta\left(\frac{\sigma}{r} - \|x_n^k - x_{\mathcal{N}(n)}^k\|\right)}.$$

Embedding dimension

Figure below shows the minimum embedding dimension using the x -component of the Lorenz system with the modified FNN method. It can be seen at $k = 3$, already the percentage of false nearest neighbors is very close to zero and for $k > 3$, this value falls to zero.

Dynamic invariants

- A **dynamic invariant** describes the behaviour of a dynamical system.

Dynamic invariants

- A **dynamic invariant** describes the behaviour of a dynamical system.
- It does not depend on the coordinate system.

Dynamic invariants

- A **dynamic invariant** describes the behaviour of a dynamical system.
- It does not depend on the coordinate system.
- Value of dynamic invariant obtained from original dynamic system is equal to the one obtained from time delay embedding!

Why are dynamic invariants important ?

- We are mostly interested in **bounded** dynamical systems.

Why are dynamic invariants important ?

- We are mostly interested in **bounded** dynamical systems.
- Bounded dynamics can be periodic or non-periodic.

Why are dynamic invariants important ?

- We are mostly interested in **bounded** dynamical systems.
- Bounded dynamics can be periodic or non-periodic.
- The average rate of divergence of nearby trajectories, i.e., the **Lyapunov exponent** can differentiate between these two dynamics.

Why are dynamic invariants important ?

- We are mostly interested in **bounded** dynamical systems.
- Bounded dynamics can be periodic or non-periodic.
- The average rate of divergence of nearby trajectories, i.e., the **Lyapunov exponent** can differentiate between these two dynamics.
- **Chaotic dynamics** : deterministic, bounded and aperiodic.

Why are dynamic invariants important ?

- We are mostly interested in **bounded** dynamical systems.
- Bounded dynamics can be periodic or non-periodic.
- The average rate of divergence of nearby trajectories, i.e., the **Lyapunov exponent** can differentiate between these two dynamics.
- **Chaotic dynamics** : deterministic, bounded and aperiodic.
- Chaotic dynamics can also be described in terms of dynamic invariants.

Why are dynamic invariants important ?

- We are mostly interested in **bounded** dynamical systems.
- Bounded dynamics can be periodic or non-periodic.
- The average rate of divergence of nearby trajectories, i.e., the **Lyapunov exponent** can differentiate between these two dynamics.
- **Chaotic dynamics** : deterministic, bounded and aperiodic.
- Chaotic dynamics can also be described in terms of dynamic invariants.
- Positive Lyapunov exponent \rightarrow Bounded system with deterministic chaotic dynamics

The Lyapunov exponent

- Chaos in a dynamical system can be characterized by the exponential separation of nearby trajectories.

The Lyapunov exponent

- Chaos in a dynamical system can be characterized by the exponential separation of nearby trajectories.
- The Lyapunov exponent can be used to measure this separation.

The Lyapunov exponent

- Chaos in a dynamical system can be characterized by the exponential separation of nearby trajectories.
- The Lyapunov exponent can be used to measure this separation.
- Since chaotic orbits are bounded, this separation and hence the mutual distance between two diverging points cannot tend to infinity.

The Lyapunov exponent

- Chaos in a dynamical system can be characterized by the exponential separation of nearby trajectories.
- The Lyapunov exponent can be used to measure this separation.
- Since chaotic orbits are bounded, this separation and hence the mutual distance between two diverging points cannot tend to infinity.
- Consider two neighboring points x_0 and $x_0 + \delta x_0$, with initial separation δx_0 . The divergence of these two points occurs at a rate given by

$$\delta x(t) \approx e^{\lambda t} |\delta x_0|$$

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.
- An k -dimensional system has k Lyapunov exponents.

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.
- An k -dimensional system has k Lyapunov exponents.
- The behavior of such an k -dimensional subspace is given by the sum of k Lyapunov exponents.

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.
- An k -dimensional system has k Lyapunov exponents.
- The behavior of such an k -dimensional subspace is given by the sum of k Lyapunov exponents.
- Evidently, positive Lyapunov exponents result in exponential separation of trajectories (hallmark of chaotic behavior).

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.
- An k -dimensional system has k Lyapunov exponents.
- The behavior of such an k -dimensional subspace is given by the sum of k Lyapunov exponents.
- Evidently, positive Lyapunov exponents result in exponential separation of trajectories (hallmark of chaotic behavior).
- If all the Lyapunov exponents are negative, it means that all the orbits are stable as the separation will exponentially decrease to zero, i.e., contraction occurs.

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.
- An k -dimensional system has k Lyapunov exponents.
- The behavior of such an k -dimensional subspace is given by the sum of k Lyapunov exponents.
- Evidently, positive Lyapunov exponents result in exponential separation of trajectories (hallmark of chaotic behavior).
- If all the Lyapunov exponents are negative, it means that all the orbits are stable as the separation will exponentially decrease to zero, i.e., contraction occurs.
- For dissipative systems, the sum of all the Lyapunov exponents must be negative.

The Lyapunov exponent

- The parameter λ is known as the Lyapunov exponent.
- An k -dimensional system has k Lyapunov exponents.
- The behavior of such an k -dimensional subspace is given by the sum of k Lyapunov exponents.
- Evidently, positive Lyapunov exponents result in exponential separation of trajectories (hallmark of chaotic behavior).
- If all the Lyapunov exponents are negative, it means that all the orbits are stable as the separation will exponentially decrease to zero, i.e., contraction occurs.
- For dissipative systems, the sum of all the Lyapunov exponents must be negative.
- Flows have one zero exponent \rightarrow for the occurrence of chaos at least three dimensions are required!

The Lyapunov exponent

- The Lyapunov exponent characterizes the exponential growth between two points on a chaotic attractor as $t \rightarrow \infty$.

The Lyapunov exponent

- The Lyapunov exponent characterizes the exponential growth between two points on a chaotic attractor as $t \rightarrow \infty$.
- The ordering of Lyapunov exponents in a k -dimensional system : $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \cdots \geq \lambda_k$

The Lyapunov exponent

- The Lyapunov exponent characterizes the exponential growth between two points on a chaotic attractor as $t \rightarrow \infty$.
- The ordering of Lyapunov exponents in a k -dimensional system : $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \cdots \geq \lambda_k$
- A chaotic system has **at least one positive** Lyapunov exponent.


The Lyapunov exponent

- The Lyapunov exponent characterizes the exponential growth between two points on a chaotic attractor as $t \rightarrow \infty$.
- The ordering of Lyapunov exponents in a k -dimensional system : $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \cdots \geq \lambda_k$
- A chaotic system has **at least one positive** Lyapunov exponent.
- **Note** : Lyapunov exponents are invariant under deformations of the attractor that preserve its topology.


The Lyapunov exponent

- The Lyapunov exponent characterizes the exponential growth between two points on a chaotic attractor as $t \rightarrow \infty$.
- The ordering of Lyapunov exponents in a k -dimensional system : $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_k$
- A chaotic system has **at least one positive** Lyapunov exponent.
- **Note** : Lyapunov exponents are invariant under deformations of the attractor that preserve its topology.
- If the reconstruction is done correctly, λ obtained from reconstructed attractor is equal to that of the original system.



The Lyapunov exponent



The Lyapunov exponent



The Lyapunov exponent



How to compute Lyapunov exponents from time series data ?

- Perform embedding after selecting τ and k

Entropy measures