

CHAPTER 3. PLANT KINGDOM

INTRODUCTION

In the previous chapter, we looked at the broad classification of living organisms under the system proposed by Whittaker (1969) wherein he suggested the Five Kingdom classification viz. Monera, Protista, Fungi, Animalia and Plantae. In this chapter, we will deal in detail with further classification within Kingdom Plantae popularly known as the 'plant kingdom'.

We must stress here that our understanding of the plant kingdom has changed over time. Fungi and members of the Monera and Protista having cell walls have now been excluded from Plantae though earlier classifications put them in the same kingdom. So, the cyanobacteria that are also referred to as blue green algae are not 'algae' any more. In this chapter, we will describe Plantae under

1. **Algae**
2. **Bryophytes**
3. **Pteridophytes**
4. **Gymnosperms and**
5. **Angiosperms.**

Let us also look at classification within angiosperms to understand some of the concerns that influenced the classification systems.

Classification within Angiosperms

1. Artificial system of classification

- Given by Linnaeus
- Based on superficial morphological characters such as habit, color, number and shape of leaves, etc.
- They were based on vegetative characters and androecium structures.
- Gave equal importance to vegetative and sexual characteristics

This is not acceptable since we know that often the vegetative characters are more easily affected by environment. As against this Natural Classification System developed.

2. Natural system of classification

- Based on morphology, anatomy, embryology, phytochemistry, cytological, physiological and reproductive characters.
- Given by George Bentham and Joseph Dalton Hooker

3. Phylogenetic system of classification - based on evolutionary relationship. Given by Englar & Prantl

4. Numerical Taxonomy

- Based on all observable characteristics
- Numbers and codes assigned to all characters
- Easily carried out using computers

5. Cytotaxonomy – Based on cytological information such as chromosome number, structure, behavior

6. Chemotaxonomy – Based on chemical constituents of plant to resolve doubts and confusions

7. **Cladistics taxonomy** – is an approach to biological classification in which organisms are categorized based on shared derived characteristics that can be traced to a groups most recent common ancestor and are not present in more distant ancestors

I) Division: - Algae

- Includes chlorophyll-bearing, simple, thalloid, autotrophic, and largely aquatic (freshwater and marine) organisms
- They occur in a variety of other habitats such as moist stones, soils, wood.
- Some occur in association with fungi (lichens) and animals (on sloth bear).
- Size ranges from microscopic unicellular forms such as *Chlamydomonas* to colonial forms such as *Volvox* and to filamentous forms such as *Ulothrix* and *Spirogyra*.
- Massive plant-like bodies are seen in some marine forms (such as **kelps**).
- Algae reproduce by **vegetative**, **asexual** and **sexual** methods.
- Vegetative reproduction is by **fragmentation**. Each fragment develops into a Thallus (as in *Spirogyra*).
- Asexual reproduction is by the production of different of spores, called as **zoospores**. They are flagellated (motile) and on germination gives rise to new plants.
- Sexual reproduction takes place through fusion of two gametes.
 - These gametes can be flagellated and similar in size (as in *Chlamydomonas*) or non-flagellated (non-motile) but similar in size (as in *Spirogyra*). Such reproduction is called **isogamous**.
- Fusion of two gametes dissimilar in size, as in some species of *Chlamydomonas* is termed as **anisogamous**.
- Fusion between one large, non-motile (static) female gamete and a smaller, motile male gamete is termed **oogamous**, e.g., *Volvox*, *Fucus*.

Economic Importance of Algae

- Carbon dioxide fixation on earth is majorly carried out by algae.
- Important as primary producers of energy-rich compounds
- Many species of *Sargassum*, *Laminaria*, and *Porphyra* are marine algae used as food
- Some brown and red algae species produce water-holding hydrocolloids.
Example – **Algin** (brown algae) and **carrageen** (red algae)
- Agar obtained from *Gelidium* and *Gracilaria* is used to grow microbes and in preparation of ice creams and jellies.
- Chlorella* and *Spirulina* are protein-rich unicellular algae, used as food supplements. They are also known as **space food**.
- The algae are divided into three main classes:-
 - Chlorophyceae** (Green algae)
 - Phaeophyceae** (Brown algae)
 - Rhodophyceae** (Red algae)

i) Chlorophyceae (Green algae)

- The members of Chlorophyceae are commonly called **green algae**.
- Occur in fresh water, brackish water and salt water.
- The plant body may be unicellular, colonial, or filamentous.
- They are usually Grass green in color due to abundance of pigment **chlorophyll a** and **b**.
- The chloroplasts may be discoid, plate like, reticulate, cup shaped, spiral or ribbon shaped in different sps.
- Chloroplast of most of the Chlorophyceae contains **pyrenoids**.

- **Pyrenoids** – Are storage bodies containing proteins in addition to starch.
- Food storage occurs in the form of oil droplets in some algae.
- Cells have rigid cell wall: inner layer made of cellulose, outer layer made of pectose.
- Examples include *Volvox*, *Chara*, *Chlamydomonas*, *Ulothrix* and *Spirogyra*
- Vegetative reproduction takes place by **fragmentation**.
- Asexual reproduction takes place by **flagellated (2 – 8) Zoospores** produced in **Zoosporangia**.
- The sexual reproduction shows variation in the type & formation of sex cells and may be **Isogamous, Anisogamous, or Oogamous**.

ii) Phaeophyceae (Brown algae)

- The member of Phaeophyceae or brown algae is primarily marine forms.
- They show great variation in size and form.
- They range from simple-branched, filamentous forms (*Ectocarpus*) to profusely branched forms such as **kelps** (may reach a height of 100 m).
- Possess **chlorophyll a, c, carotenoids, and xanthophylls**.
- Vary in colour from olive green to various shades of brown (depending on amount of **xanthophylls** and **fucoxanthin**).
- Food stored as complex carbohydrates such as **laminarin** or **mannitol**.
- Vegetative cells have cellulosic wall covered on the outside by gelatinous coating of **algin**.
- Cell contains a centrally located vacuole and nucleus in addition to plastids.
- The plant body is usually attached to the substratum by a **holdfast** and has a stalk the **Stipe** and leaf like photosynthetic organ **Fron**d.
- Vegetative reproduction takes place by **fragmentation**.
- Asexual reproduction in most brown algae is by **biflagellate zoospores** that are pear-shaped and have two unequal laterally attached flagella.
- Sexual reproduction may be isogamous, anisogamous or oogamous.
- Union of gametes takes place in water or within oogonium (oogamous species).
- The gametes are pyriform (pear-shaped) and bear two laterally attached flagella.
- Example – *Ectocarpus*, *Dictyota*, *Laminaria*, *Sargassum*, and *Fucus*.

iii) Rhodophyceae (Red algae)

- Commonly called red algae due to the presence of red pigment, **r-phycoerythrin**. Also contains chlorophyll a and d.
- Mainly marine forms with bulk mass inhabiting warmer areas.
- Occur in well-lighted regions i.e., close to the surface of water and also in deeper areas.
- Red thalli of most of these species are multicellular. Some have complex body organization.
- Cell wall is made up of cellulose.
- Food is stored as **Floridian starch** similar to amylopectin and glycogen in structure.
- The red algae usually reproduce vegetatively by **fragmentation**.
- They reproduce asexually by **non-motile spores** and sexually by **non-motile gametes**.
- Sexual reproduction is **oogamous**. Complex post fertilization developments
- Example – *Polysiphonia*, *Gelidium*, *Gracilaria*, *Porphyra*

II) Division: - Bryophyta

- Bryophytes are also called **amphibians of plant kingdom** since they live on land, but depend on water for sexual reproduction.
- Usually occur in cool, damp, and shady areas.
- Play an important role in plant succession on bare rocks/soils.
- Plant body more differentiated than algae.
- Plant body is thallus-like and prostrate or erect and attached to substratum by unicellular or multicellular **rhizoids**.
- They lack true roots, stem and leaves; may possess root-like, stem-like, and leaf-like structures.
- The plant body of the Bryophytes is **haploid**. It produces gametes, hence is called **Gametophyte**.
- The sex organs in bryophytes are multicellular.
- The male sex organ is called **Antheridium**. It produces biflagellate **Antherozoid**.
- The female sex organ is called **Archegonium**. It is flask shaped and produces a **single egg**.
- The antherozoids are released into the water where they come in contact with Archegonium. An antherozoid fuses with egg to produce **Zygote**. (Zygote do not undergo reduction division i.e. meiosis immediately.)
- The zygote grows into a multicellular body called **Sporophyte**.
- Sporophyte is dependent on gametophyte for food. Hence, it remains attached to the gametophyte.
- Few cells of sporophyte undergo meiosis to produce **haploid spores**.
- Spores germinate to form gametophyte.

Classes of bryophytes

Bryophytes are divided into two classes: - **1) liverworts, 2) Mosses**

Class: - 1. Liverworts

- Grow in moist, shady habitats such as banks of streams, marshy ground, damp soil, bark of the tree and deep in the wood.
- Plant body of the liverwort is thalloid, e.g., **Marchantia**.
- The thallus is dorsiventrally appressed to the substrate.
- Leafy members have tiny leaf-like appendages in two rows on the stem-like structures.
- Asexual reproduction takes place by **fragmentation** of thalli, or by the formation of specialized structure called **Gemmae** (sing: - Gemma).
- The Gemmae are green, multicellular, asexual buds, which develop in small receptacles called **Gemma cups** located on the thalli.
- The gemmae become detached from the plant body and germinate to form new individual.
- In sexual reproduction, male and female sex organs are produced either on the same or different thalli.
- The sporophyte is differentiated into a **foot, seta** and **capsule**.
- After meiosis spores are produced within the capsule. These spores germinate to form free-living gametophyte.

Class: - 2. Mosses

- The predominant stage of the life cycle of a moss is the gametophyte

Gametophyte

- Consist of two stages.
- The first stage is the **protonema stage**, which develops directly from the spore.
 - It is a creeping, green, branched and frequently filamentous stage.
- The second stage is the **Leafy stage**, which develops from the secondary protonema as a lateral bud
 - It consists of upright, slender axes bearing spirally arranged leaves.
 - It is attached to the soil through multicellular and branched rhizoid.
 - This stage bears sex organ.
- Vegetative reproduction in mosses is by **fragmentation** and **budding** in the secondary protonema
- In sexual reproduction, the sex organ antheridia and archegonia are produced at the apex of the leafy shoots
- After fertilization, the zygote develops into a sporophyte,

Sporophyte

- More elaborate than liverworts
- Consists of foot, seta, and capsule
- Capsule contains spores.
- Spores formed by meiosis
- Elaborate mechanism of spore dispersal
- Example – *Funaria*, *Polytrichum*, and *Sphagnum*

Economic Importance

- Provide food for herbaceous mammals, birds, and insects
- Peat provided by *Sphagnum* (a moss) is used as fuel.
- *Sphagnum* is also used as packing material in trans-shipment of living material because of their water-holding capacity.
- They form dense mats on the soil and hence prevent soil erosion.
- Mosses along with lichens form the pioneer community in land and desert succession.

III) Division: - Pteridophyta

The Pteridophytes include **Horsetails** and **Ferns**. The Pteridophytes are **used for medicinal purpose** and as **soil-binders**. They are also frequently **grown as ornamentals**.

General Characteristics

- The Pteridophytes are found in cool, damp, shady places.
- Evolutionary the Pteridophytes are the first terrestrial plants to possess vascular tissues - xylem and phloem
- In Pteridophytes **the dominant phase in the life cycle is sporophytic plant body**.
- They have well-differentiated true stem, leaves, and roots
- Leaves may be small (**microphylls**) as in *Selaginella* or large (**macrophylls**) as in ferns.
- Sporophytes bear **sporangia**, which develop in association with leaves called **sporophylls**.
- In some Pteridophytes, sporophylls form distinct, compact structures called **strobili** or **cones** (*Selaginella*, *lycopodium*, *Equisetum*).

- Sporangia produce spores by meiosis in spore mother cells.
- Spores germinate to form small, multicellular, free-living photosynthetic thalloid gametophyte called **prothallus**.

Prothallus (Gametophyte)

- Require cool, damp, shady places to grow
- Also require water for sexual reproduction
- The gametophytes bears male and female sex organ called **Antheridia** and **Archegonia**, respectively
- Fusion of male gamete (**Antherozoid**) with the egg present in the Archegonium results in the formation of **Zygote**
- Zygote produces well-differentiated, multicellular **sporophyte** (dominant phase of the Pteridophytes.)


Sporophyte

- In majority of the Pteridophytes all the spores are of similar kinds; such plants are called **Homosporous**.
- Genera like *Selaginella* and *Salvinia* which produce two kinds of spores, large (**Macrospore**) and small (**Microspore**) are called **Hetresporous**.
- The megaspores and microspores germinate and give rise to female and a male gametophyte, respectively.
- The female gametophytes in these plants are retained on the parent sporophyte for variable periods.
- The development of the zygotes into the young embryos takes place within the female gametophytes. This step is a precursor to the **seed habit** considered an important step in evolution.
- The Pteridophytes are classified into four classes
 1. **Psilopsida** (whisk fern) Ex - *Psilotum*
 2. **Lycopsida** (club mosses) Ex - *Selaginella*, *Lycopodium*
 3. **Sphenopsida** (Horse tail) Ex - *Equisetum*
 4. **Pteropsida** (ferns) Ex - *Dryopteris*, *Pteris*, *Adiantum*, *Salvinia*, *Marsilea*.

IV) Division: - Gymnospermae

- Word gymnosperms, ***gymnos*** – naked, ***sperma*** – seeds
- Ovules not enclosed by any ovary wall
- Seeds formed after fertilization are not covered (i.e., naked).
- Include medium-sized trees, shrubs, and tall trees
- Contains the world's largest plant *Sequoia* - the giant redwood
- Plants have **tap roots**. Roots in some genera show **symbiotic associations**.
 1. **Mycorrhiza** shows association of fungi with ***Pinus*** roots.
 2. Coralloid roots of ***Cycas*** show association with **N₂-fixing Cyanobacteria**.
- The stems could be branched (*Pinus*, *Cedrus*) or unbranched (*Cycas*)
- The leaves may be simple (*Pinus*) or compound (*Cycas*)
- The leaves in gymnosperms are well-adapted to withstand extreme temperature, humidity and wind. In conifers, needle-like leaves with thick cuticle and sunken stomata reduce surface area and help to reduce water loss.
- Gymnosperms are **Hetresporous**.
- They produce haploid **microspores** and **megaspore** within **sporangia** that are borne on **sporophylls** which are arranged spirally along the axis to form lax or compact **Strobilli** or **cone**.

- The strobili bearing only microsporophylls and microsporangia are called **Microsporangiate** or **male strobili**.
- The microspores in the microsporangia develop into a **male gametophyte** generation which is reduced and is confined to only a limited number of cells. This reduced gametophyte is called a **pollen grain** and it develops within the microsporangia.
- The strobili or cones bearing megasporophylls with ovules or megasporangia are called **Macrosporangiate** or **female strobili**.
- Male and female strobili may be borne on same tree (***Pinus***) or on different trees (***Cycas***).
- Megaspore mother cell divides meiotically to form four megaspores.
- Megaspore mother cell is a differentiated cell of nucellus. Nucellus protected by envelopes is known as an **ovule**.
- One of the megaspore enclosed within the megasporangium(nucleus) develops into a multicellular female gametophyte that bears two or more female sex organ or **Archegonia** which remains within megasporangium
- Male and female gametophytes do not have independent existence, hence remain within sporangia.
- Steps in fertilization:



Gymnosperms are classified into three classes.

1. **Cycadopsida** (cycadophyta). Ex – *Cycas*
2. **Coniferopsida** (conifers). Ex – *Pinus*, *Sequoia*, *Ginkgo* (Living fossil)
3. **Gnetopsida**. Ex – Gnetum.

V) Division: - Angiosperms (Flowering plant)

- Unlike the gymnosperms where the ovules are naked, in the angiosperms or flowering plants, the pollen grains and ovules are developed in specialized structures called flowers.
- In angiosperms, the seeds are enclosed in fruits.
- The angiosperms are an exceptionally large group of plants occurring in wide range of habitats.
- They range in size from the smallest *Wolffia* to tall trees of *Eucalyptus* (over 100 metres).
- They provide us with food, fodder, fuel, medicines and several other commercially important products.
- They are divided into two classes: the dicotyledons and the monocotyledons.