

# Worksheet — Fruits and Seeds

Student Name: \_\_\_\_\_ Date: \_\_\_\_\_

Section: \_\_\_\_\_ Class/Roll No: \_\_\_\_\_

## A. Fill in the Blanks

1. A fruit developing without fertilization is called \_\_\_\_\_.
2. The outer layer of the pericarp is called the \_\_\_\_\_.
3. The middle layer of the pericarp is called the \_\_\_\_\_.
4. The single cotyledon in monocot seeds is called the \_\_\_\_\_.

## B. One Word Answer

1. Name the fruit type of mango and coconut. Ans: \_\_\_\_\_
2. Name the outer layer of the seed coat. Ans: \_\_\_\_\_
3. Name the sheath protecting the plumule in monocot seeds. Ans: \_\_\_\_\_
4. Name the sheath protecting the radicle in monocot seeds. Ans: \_\_\_\_\_

## C. Match the Following

| Column A  | Column B          |
|-----------|-------------------|
| Epicarp   | Outer layer       |
| Mesocarp  | Middle layer      |
| Endocarp  | Inner layer       |
| Scutellum | Monocot cotyledon |

## D. True or False

1. Dicot seeds usually have two cotyledons. (True/False)
2. Monocot seeds usually lack endosperm. (True/False)
3. Mango is classified as a berry. (True/False)
4. Parthenocarpic fruits contain seeds. (True/False)

***E. Short Answer Questions***

1. Define parthenocarpic fruit with an example. **[2 Marks]**
2. Differentiate between testa and tegmen. **[2 Marks]**

***F. Short Essay Questions***

1. Describe the three layers of the pericarp with examples. **[3 Marks]**
2. Differentiate between dicot and monocot seeds. **[3 Marks]**

***G. Long Answer / Essay Questions***

1. Describe in detail the structure of a monocot seed, including the roles of the scutellum, coleoptile, and coleorhiza. **[5 Marks]**