



Microalgae for microplastic removal from water and wastewater: a review

Paulo M. S. Sousa^{1,2} · Kerry A. Kinney³ · Cátia A. Sousa^{1,2,4,5,6} · Manuel Simões^{1,2,6}

Received: 17 April 2024 / Accepted: 23 December 2024 / Published online: 5 January 2025
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2025

Abstract

The worldwide microplastic pollution in waters requires efficient removal techniques, yet actual wastewater treatment methods are limited. Here we review the use of microalgae for microplastic removal, with focus on microplastics in aquatic systems and wastewaters, legislation and regulations, common removal techniques, and microalgae for microplastic removal. We describe the mechanisms involved in microalgae-microplastics aggregation. We also present the criteria for selecting adequate microalgae for microplastics removal from wastewater.

Keywords Aquatic contamination · Microplastic pollution · Microalgae-based system · Microalgae-microplastics interaction · Removal mechanism · Wastewater treatment

Introduction

Microplastics are emerging contaminants of worldwide concern due to their negative impacts on aquatic ecosystems and human health [1]. Microplastics are defined as tiny solid plastic particles between 1 µm and 5 mm in diameter that

have a potentially negative impact on aquatic environments and are harmful to all living organisms [1, 2]. The annual global production of plastics is enormous with 400.3 million tons generated in 2022 [3]. According to data collected in 2019 by the Organization for Economic Cooperation and Development (OECD), a massive 353 million tons of plastic waste was generated worldwide, with only 9% being recycled, and nearly 50% ending up in landfills [4]. Moreover, in 2019, of the global plastic waste generated, an estimated 6.1 million tons was discharged into water environments, and 1.7 million tons flowed into the oceans [4]. This poses a threat to the quality and availability of water on our planet. It is estimated that there are 30 million tons of plastic waste in seas and oceans (Fig. 1), with rivers being the primary locations for plastic waste in aquatic environments, around 109 million tons [4]. Recently, Eriksen et al. [5] monitored the rapid increase of floating plastic waste estimating that there are about 82 to 358 trillion plastic particles weighing between 1.1 to 4.9 million tons.

Plastic waste introduced into the environment undergoes mechanical, biological, and chemical degradation processes, changing its physical and chemical properties and leading to fragmentation and the formation of microplastics [6, 7]. As a result, microplastics can reach the aquatic environment from a primary or a secondary source [8]. Primary sources directly release small particles into the environment from plastic pellets, personal care products containing microbeads, paint, household laundry wastewater, and vehicle tire

✉ Cátia A. Sousa
csousa@fe.up.pt

✉ Manuel Simões
mvs@fe.up.pt

¹ LEPABE-Laboratory for Process Engineering, Environment, Biotechnology and Energy, Faculty of Engineering, University of Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal

² ALiCE-Associate Laboratory in Chemical Engineering, Faculty of Engineering, University of Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal


³ Architectural, and Environmental Engineering, Department of Civil, The University of Texas at Austin, Austin, TX, USA

⁴ School of Engineering, ISEP/P.PORTO, Polytechnic of Porto, Rua Dr. António Bernardino de Almeida, 431, 4249-015 Porto, Portugal

⁵ School of Engineering, CIETI, Polytechnic of Porto, Porto, Portugal

⁶ Department of Chemical and Biological Engineering, Faculty of Engineering, ALiCE/LEPABE, University of Porto, Rua Dr. Roberto Frias, s/n, 4200-465 Porto, Portugal

Fig. 1 Global pollution of plastic and microplastics. Since 2019, an estimated 30 million tons of plastic waste currently pollutes the oceans, while rivers are the main spot for plastic waste in aquatic environments, containing approximately 109 million tons [Eriksen et al. [15], GESAMP [6], OECD [4], and Eriksen et al. [5]]. The primary and major source of plastics and microplastics contamination in aquatic environments is the ineffective wastewater treatment by current processes. As a result, a novel and sustainable approach using microalgae-based systems is gaining attention within the scientific community for removing these pollutants



wear, among others [6, 9–11]. Secondary sources of microplastics are generated by the degradation of larger plastic objects into smaller plastic fragments after exposure to specific conditions in aquatic ecosystems, such as plastic bags and bottles, fishing wastes, farming film, and other large-size plastic wastes [9].

Several recent studies have reviewed microplastics pollution in soils [12], oceans, freshwater, and drinking water [13–16]. However, wastewater represents a significant source of microplastics released into aquatic systems [17, 18] and warrants greater attention, particularly since current wastewater treatment processes are unable to remove microplastics efficiently. This work reviews the growing problem of microplastics pollution in aquatic environments and assesses the ineffectiveness of conventional wastewater treatment processes for its removal (Fig. 1). This work is further strongly focused on the high potential of biological systems for bioremediation processes, applying microalgae-based systems in particular. The feasible mechanisms, as well as the advantages of applying these microorganisms to remove microplastics from wastewater, were also reviewed.

Categories of microplastics

From the million tons of plastic waste generated, all kinds of products are found with different polymers and chemical compositions. Plastics are beneficial in daily lives due to their wide applicability in many fields or industrial sectors, as well as their physical and mechanical characteristics, *e.g.*, elasticity, insulation, density, durability, waterproof, and corrosion resistance, among others, and their cost-effective production [17, 19–21]. They are synthesized from

low-molecular organic units (monomers) [20] and the most common organic units found in different kinds of plastics are ethylene, propylene, styrene, phenol, formaldehyde, vinyl chloride, and acetonitrile. These monomers submitted to a polymerization process are responsible for the production of different classes of plastics. For example, propylene is the base constituent for polypropylene (PP) plastics, styrene for polystyrene (PS) polymers and vinyl chloride is a monomer for polyvinyl chloride (PVC), products widely used in our daily lives. However, other polymers require more complex syntheses, involving chemical reactions with alcohols, *e.g.*, ethylene glycol, acids, *e.g.*, terephthalic acid, or methyl methacrylate to develop complex polymers (Table 1).

The most common types of microplastics that are of worldwide environmental concern are the following, as summarized by Plastics Europe [22] and Plastic Oceans [23] (Table 1): i) polypropylene (PP); ii) polyethylene (PE) divided into two subtypes — iii) high-density polyethylene (HDPE) and iv) low-density polyethylene (LDPE); v) polyvinyl chloride (PVC); vi) polyethylene terephthalate (PET); vii) polystyrene (PS); and viii) polyurethane (PUR). Despite these eight different types of polymers, other types of microplastics can be found dumped into ecosystems, particularly polymethyl methacrylate (PMMA) and polyacrylonitrile (PAN) [24–26].

Also, microplastics derived from polypropylene (PP), polyethylene (PE) (high-density polyethylene and low-density polyethylene), and polyethylene terephthalate (PET) comprise approximately 78% of the estimated waste generated worldwide (Table 1). Moreover, these plastics have a long estimated degradation time, particularly high-density polyethylene (HDPE) products, which can require up to 1200 years to biodegrade [27]. Therefore, considering

Table 1 The most polluting plastic waste, sources, and estimated degradation time

Microplastics	Density (g/cm ³)	Chemical formula	The most common waste debris	Estimated degradation time (years)	Estimated waste world-wide generated (Million tons)	References
PP	0.90–0.91	[CH ₂ –CH(CH ₃)] _n	Microwave dishes, ice cream tubs, bottle caps, single-use face masks	Up to 450	62 ^a	[4, 9, 28, 29]
HDPE	0.95–0.97	(CH ₂ –CH ₂) _n	Shampoo bottles, milk bottles, freezer bags, ice cream containers	Up to 1200	44.7 ^a	[4, 9, 27, 30, 31]
LDPE	0.92–0.93	(CH ₂ –CH ₂) _n	Bags, trays, containers, food packaging film	Up to 1000	49.3 ^a	[4, 9, 27, 30–32]
PVC	1.3–1.6	(CH ₂ –CHCl) _n	Water pipes, packaging, wires and cables, blood bags	Up to 50	21.2 ^a	[4, 9, 30, 33, 34]
PET	1.3–1.4	(C ₁₀ H ₈ O ₄) _n	Water bottles, dispensing containers, biscuit trays	approximately 450	24.9 ^a	[4, 9, 27, 30, 35]
PS	1.0–1.1	[CH ₂ –CH(C ₆ H ₅)] _n	Cutlery, plates, cups	Between 500–900	15.2 ^a	[4, 27, 30, 36, 37]
PUR	1.05	[C ₃ H ₈ N ₂ O] _n	Footwear, thermal isolation	20–30	11.3 ^a	[4, 9, 30, 38, 39]
PMMA	1.2	[CH ₂ –C(CH ₃)CO ₂ CH ₃] _n	Shatterproof windows, skylights, illuminated signs, and optical fibers	Not mentioned	≈1	[40, 41]
PAN	1.18	(CH ₂ –CHCN) _n	Fibers in the hot gas filtration system, outdoor awnings, sail for yachts	Not determined	Not determined	[42]

PP-polypropylene; HDPE-high-density polyethylene; LDPE-low-density polyethylene; PVC-polyvinyl chloride; PET-polyethylene terephthalate; PS-polystyrene; PUR-polyurethanes; PMMA-polymethyl methacrylate; PAN-polyacrylonitrile

^athe values listed in the literature are relative to the year 2019

the characteristics of microplastics and their very extensive biodegradation timescale (Table 1), the scientific community has initiated research to understand the phenomena of microplastics accumulation in the environments, particularly in aquatic ecosystems.

Microplastics in aquatic ecosystems

Microplastics have been increasingly accumulating in aquatic environments through anthropogenic action, prompting an emergent research area of “White Pollution”. Over the last decades, numerous studies have been published investigating the presence of microplastics in oceans, revealing concerning numbers [6, 8, 15, 43, 44]. For example, Eriksen et al. [5] analyzed data on floating ocean plastics from various stations worldwide (approximately 12,000 stations) over 40 years, estimating that the global abundance of plastic particles floating in the oceans is approximately 82 to 358 trillion, with a weight of 4.9 million tons. The particles enter the marine environment via different primary or secondary pathways, through land-based human activity, or in

the marine environment, *e.g.*, fishing [5]. Subsequently, they are dispersed and transported across the oceans by large-scale abiotic factors such as sea currents, waves, wind, geostrophic circulation, and weather events [6, 8, 15, 44, 45]. Other important variables are the density, shape, and size of microplastics (Table 1), which also affect the transportation and dispersion patterns [45].

Even though the majority of plastic debris accumulates in areas of the northern hemisphere (56%) [15], probably due to a higher density of human activity and population, microplastics are ubiquitous in global ocean regions, including in the frozen Arctic and Antarctic waters, in seabed sediments, beaches, in coastal waters, and throughout the water column [8, 15, 44, 46–55]. According to the International Maritime Organization (IMO) report [6], the polymers most frequently found in microplastics recovered from marine environments are polyethylene (PE) (79%), polypropylene (PP) (64%), and polystyrene (PS) (40%). The IMO report offers a comprehensive study, gathering information on the industrial synthesis of the different types of polymers, including some ordinary applications, sources (primary or secondary) of plastic waste

in the aquatic environment, environmental exposure to degradation or fragmentation into microplastics, and the factors influencing microplastics dispersion and accumulation in worldwide seas and oceans [6].

Microplastics pollution extends to freshwater environments, such as the dynamic natural ecosystems, *e.g.*, rivers, lakes, estuaries, ice and glaciers, among others, and terrestrial systems, *e.g.*, urban waters and wastewater [45, 56]. This pollution rapidly spreads, leading to numerous persistent plastic particles and fibers in worldwide freshwater ecosystems (Table 2), endangering their crucial chemical and microbiological quality for human subsistence [45, 56].

The abundance of microplastics in freshwaters depends on natural environments (*i.e.* dynamic or static ecosystems) and anthropogenic factors. In fact, in an open and dynamic ecosystem, like rivers, the abundance of microplastics can be conditioned by the river course, ending in the marine environment causing the dispersion of the particles. On the other side, static and stagnant environments, such as lakes, stimulate the accumulation or deposition of microplastics in the ecosystem [45, 118]. Moreover, microplastics enter into freshwaters through different sources: i) surface runoff (resulting from weather conditions), or wastewater effluents (both treated and untreated); ii) sewage overflows, industrial effluents (primary source); and iii) degraded plastic waste or atmospheric deposition (Fig. 2). Surface runoff and wastewater effluents are recognized as the main sources of pollution for freshwater systems [119]. According to Liu et al. [120], an average of 7.2 billion microplastic particles per day enter rivers from wastewater treatment plants (Fig. 2).

Based on data from different studies (Table 2), microplastics are omnipresent across worldwide water and their surroundings, stimulating the accumulation or deposition of microplastics in water ecosystems. Moreover, no pattern directly correlates the highest abundance of microplastics with a particular aquatic environment or geographical country. The concentrations of the different polymers are heterogeneous even in the same type of environment or country.

Regulation and legislation on microplastics

One of the biggest gaps concerning microplastics is the lack of standard international regulations regarding microplastics discharge and monitoring in water, particularly wastewater [121]. As such, urgent actions are needed to establish policies and guidelines to minimize, or even ban, the production of certain plastics and mitigate the presence of microplastics in wastewater and not only their consumption. Although there are worldwide regulation gaps concerning plastic production, limited progress has been made in some countries to ban specific plastic products from being manufactured and marketed [122–125]. For instance, in Europe, at the request of the European Commission, the European Chemicals

Agency proposed in 2019 a wide restriction on microplastics in products on the market in the European Union and European Economic Area to prevent or reduce their release into the marine environment, *e.g.*, single-use plastics (Directive 2019/904) [126]. More recently, in August 2022, the European Commission released a draft proposal to restrict intentionally added microplastics in daily-use products. The restriction would ban the use of synthetic polymer microparticles in a diverse range of consumer products, for example, in cosmetics, cleaning products, pesticides, and sports fields, among others-banning particles smaller than 5 mm and fiber-like particles smaller than 15 mm [122, 127].

Furthermore, the European Commission also revised and improved the Directive to handle microparticle pollution in wastewater treatment, providing a strong starting point for the implementation of measures to minimize the dispersion of microplastics in the environment. In the Urban Waste Water Treatment Directive (UWWTD) (Council Directive 91/271/EEC), new standards and goals were added to mitigate wastewater pollution, with the main focus on: i) reducing pollution and energy used in wastewater treatment plants (more technological development and recycling); ii) improving water quality; iii) making industry pay for the treatment of micro-pollutants; and iv) leading more to a circular sector [127].

Across the European regulatory landscape, different governmental regulations and laws are drafted in each country, in parallel with the European Commission Directives [123, 124, 126, 130, 131]. For instance, in 2019, the Portuguese government put into force the law No. 76/2019 to ban the use in commercial establishments or the commercial sale of "plastic tableware" and other materials made of polystyrene (PS), *e.g.*, cotton swabs, to reduce consumption by 90% before December 2030 [132]. More restrictive measures have been applied in France, Italy, Germany, and Belgium, among others, with complete bans on plastic bags and selling other single-use products made of different types of plastic [124, 133].

Great efforts have also been made beyond Europe as well to regulate the manufacture and sale of plastic products [130]. The United Nations has developed guidelines for controlling and preventing the generation of plastic waste, and there are also relevant studies outlining policies applied by several countries around the world (with different enforcement efforts) to reduce or ban plastic waste, avoiding reaching aquatic ecosystems [124, 130, 133].

Regulatory efforts to ban the manufacture or introduction of plastic microbeads in cosmetic and hygiene products (primary microplastics in waters) have been implemented by many countries such as the United States since December 2015 ("Microbead-Free Waters Act"), in Canada since 2016, in England since 2019, joining the European Union in this initiative [122, 123, 130]. But, looking at the US regulation,

Table 2 Pollution by microplastics of water environments, with the respective concentrations and sizes of microplastics

Country or Ocean region	Aquatic Environment	Most types of microplastics (shape)	The abundance of the microplastics	Size (mm)	References
Portugal	River–freshwater	PP (29%), PE (29.4%), PS (8.8%) and PET (8.8%)–foam, film and fibers	≤ 1265 particles/m ³	≥ 0.055	[57]
	Ocean coast–marine (surface water)	Predominantly pellets, fragments, and polymeric foams	1908 ± 454 items/m ²	4.0–5.0	[58]
China	Ocean–marine	PP, PE, PS–film, fiber, granule, pellet	545 ± 282 particles/m ³	< 0.50	[59]
	Urban Waters	Not determined	1597 to $12,611$ particles/m ³	< 1.00	[60]
	Urban Waters	PP (34%), PE (29%), PS (23%)–Fibers	911.57 ± 199.73 to 3395.27 ± 707.22 particles/m ³	0.50 – 1.00	[61]
	Lake–freshwater (surface water)	50–91% fibers, 5.67–33.33% beads, 2.63–20.00% fragments	0.62 to 4.31 particles/m ²	0.9 – 0.333	[62]
	River–freshwater	PP (35.7%), PE (28.6%) and PET (28.6%)–fibers (80.9%), fragments (18.9%) and films (2.2%)	379 to 7924 items/m ³	0.02 – 1 mm (48%) and 1 – 2 mm (36.5%)	[63]
	River–freshwater	Fragment, fiber, film, and granules	0.0158 ± 0.0098 items/ m ³	0.2 – 0.5 (56.4%)	[64]
	Ocean–marine	Fibrous rayon (67%) and PET (23%)–fibers (88 to 100%)	0.0046 to 0.064 items/m ³	0.010 – 4.556	[65]
	River–freshwater	Fibers (71%) and fragments (17%)	5×10^{-8} to 32×10^{-5} particles/m ³	0.36 – 0.99 (72%) and 1.0 – 4.75 (26%)	[66]
	River–freshwater	Not determined	Average particles 2×10^{-5} to 6×10^{-5} /m ³		[67]
	River–freshwater	Not determined	Average particles 1×10^{-6} /m ³	0.3 – 2.0	[68]
Indonesia	Ocean–marine	Fibers	5.0 to 18.4 particles/m ³	< 0.10	[69]
	Lake–freshwater	Not determined	Average 2.7×10^{-6} particles/m ³	0.355 – 0.999 (81%) and 1.000 – 4.749 (17%)	[70]
	Sea–marine	PE (53%) and PP (43%), and also PS, PA, and PVC–fragments (86%)	≤ 4.12 microplastics/m ³ (mean 0.34 ± 0.80 microplastics/m ³)	0.5 – 5.0	[71]
	Sea–marine	PP	7×10^{-5} to 1.11×10^{-3} particles/m ³	0.270 – 1.279	[72]
Mongolia	River–freshwater	Fibers, fragments, microbeads, and films	5.77 ± 1.25 items/m ³	1.5	[73]
	Lake–freshwater	Fragments (40%), films (38%) and fibers (20%)	Average 1.2×10^{-6} particles/m ³	0.333 – 5.0	[74]
India	River–freshwater	Rayon (54%), PMMA (24%), PET (8%), PVC (6%) and PES (5%)–fiber (91%) and fragments (9%)	3.8×10^{-5} items/m ³	2.459 ± 0.209	[75]
	Ocean–marine	PET, PE, rayon, PA and PVC	2.3 ± 2.1 items/m ³	0.108 – 4.703	[76]

Table 2 (continued)

Country or Ocean region	Aquatic Environment	Most types of microplastics (shape)	The abundance of the microplastics	Size (mm)	References
Italy	Ocean-marine	PE (23.89%), PP (15.93%), alkyd (13.29%), PS (8.85%)—fibers, fragments, films, and pellets	1.25 ± 0.88 particles/m ³	0.30–4.75	[77]
	Lake-freshwater	LDPE, HDPE, PET, and PP—fragments, films and fibers	0.0088 ± 0.0003 particles/m ³	0.02–5.00	[78]
	River-freshwater	PET and PA (38%), PE (26%), PP (18%), PVC (10%), PS (5%)—filaments, fragments and films	Average 206 ± 31.9 particles/m ³	0.10–2.00	[79]
	Lake-freshwater (surface water)	Fragments and fibers	Average of 3.0 particles/m ² on Lake Chiusi surface water and an average of 2.5 particles/m ² on Lake Bolsena surface water	0.5–1.0	[80]
France	River-freshwater	Fibers (48%)	Average 0.0003 particles/m ³ in plankton net	0.1–5.0	[81]
	River-freshwater	2-D fragments and 3-D spherules	Average 0.00035 particles/m ³ in manta trawl		
Switzerland	River-freshwater	PE (62%), PP (15%) and PS (12%)—films, fragments and foams	Average 7 × 10 ⁻⁸ particles/m ³	Not determined	[82]
Belgium	River-freshwater	PP (49%), PS (49%), and PVC (2%)	Up to 0.0048 particles/m ³	0.025–1.000	[83]
Austria	River-freshwater	Pellets (17%), flakes (10.4%) and spherules (3.6%)	Average 0.0032 particles/m ³	Not determined	[84]
Spain	Glacial	Fragments (59.7%), fibers (38.8%), and spheres (1.5%)	0.0213 particles/m ³	< 0.045	[85]
Germany	Sea-marine	HDPE (64.7%), PP (27%), and PS (0.2%) – fragments (77.2%), tar (15.7%), and films (3.9%)	From 0.27–136.70 items/m ³	0.5–1.0 (49.7%) and > 1.0 (33.6%)	[86]
	Sea-marine	Granulates, fibers, and black carbon (BC) particles	1770 particles/m ³	Not determined	[55]
	River-freshwater	PE (20%), PP (16%), PVC, PET and polyester, polychloroprene, and PA—fragments	28–134 particles/m ³	0.3–1 mm	[87]
Japan	Sea-marine	Spherical particle, microbead	0.03 to 0.075 particles/m ³	< 2.00	[88]
Colombia	Lake-freshwater	Polyester (57.9%), PS (47.0%), and PET (35.3%) – fragments (20.8%), foams (0.2%), and pellets (0.1%)	Average 0.0019 particles/m ³	0.35	[89]
South Africa	Sea-marine	Fibers (90%)	257.9 ± 53.36 to 1215 ± 276.7 particles/m ³	0.08–5.00	[90]
Brazil	Sea-coastline	Fibers, fragments, styrofoam and pellets	12 to 1300 particles/m ²	Not determined	[91]

Table 2 (continued)


Country or Ocean region	Aquatic Environment	Most types of microplastics (shape)	The abundance of the microplastics	Size (mm)	References
Finland	Sea-marine	PE (81.7%), PE (16.20%) – fragments, films	1.4 to 21.3 particles/m ³	0.30–1.00	[92]
	River-freshwater	PE (29%), polyester (26%), PA (19%), ethylene vinyl alcohol (9%), nylon (9%), and HDPE (8%) – fibers	0.38 ± 0.14 – 0.71 ± 0.25 particles/m ³	Not determined	[93]
Scotland	Lake-freshwater	PE, PP, PET, PMMA, and PAN-fibers (64%) and fragments (36%)	Average concentrations 0.27 ± 0.18 microplastics/m ³	0.020–0.300	[94]
	Sea-marine	PP (26%), PS (17%), PVC (12%), PE (10%)—fragments, fibers and beads	9.113 × 10 ⁻⁸ microplastics/m ²	0.30–5.00	[95]
Denmark	Sea-marine	Fragment, fiber	0.07 ± 0.02 particles/m ³	0.30–0.63	[96]
	Sea-marine	PMMA, PA, PP, PE, and polyesters – fragments (54%) and fibers	103 ± 86 items/m ³	< 0.1	[97]
Australia	Freshwater	PE, rayon-fibers	0.004 ± 0.0027 items/m ³	0.036–4.668	[98]
	Ocean-marine	Fragment, fiber	0.031 particles/m ³	< 2.00	[88]
Iceland	Glacial	PUR, PVC, PA, and acrylonitrile butadiene styrene	Not determined	≤ 3.00	[56]
Norway	Sea-Artic	Polyester (15%), PA (15%), PE (5%), PMMA (10%), PVC (5%), cellulose-fibers (95%), fragment, films	0.34 ± 0.31 particles/m ³	1.93 ± 1.22	[99]
Sweden Malaysia	Sea-marine	Not determined	≤ 102 000 particles/m ³	≥ 0.30	[100]
	River-freshwater	PE, PS, polycarbonate, and PET-fibers (68.4%), film, and fragment	0.060 to 0.128 items/m ³	0.10–5.00	[101]
South Korea	River-freshwater	PE, PS (16.9%), silicone (23.1%), PTFE, polyester (13.8%), PP (15.4%)—fragments (73%), and fibers	Surface waters – 42.9 particles/m ³ and water column ≤ 180.0 particles/m ³	mostly < 1.00	[102]
	Jungnang stream-freshwater	PE, PP, and PET-fragments and fibers	0.0098 ± 0.008 particles/m ³	≤ 0.2	[103]
Vietnam	River-freshwater	Polyesters (70%), PET (9%), PE (7%), PP (4%) and rayon (4%)—fibers (92%)	Between 172 000 items/m ³ to 519 000 items/m ³	0.050–0.450	[104]
Iran	River-freshwater	PE (36.6%), PA, and PS-fibers (53.5%), fragments and foams	1 to 43 items/m ³ (mean 12.8 ± 10.5)	0.10–5.00	[105]
Pakistan	Lake-freshwater	PP, PE, Polyesters, PET, PVC-fibers (38.6%), sheets (2.5%), and foams (2.2%)	2074 ± 3651 particles/m ³	≤ 1.00	[106]
UK	Sea-marine (surface waters)	PP, rayon, polyester, PA, PET, and PMMA-fibers, fragments, and pellets	0.26 to 0.68 items/m ³	0.053–3.00	[107]

Table 2 (continued)

Country or Ocean region	Aquatic Environment	Most types of microplastics (shape)	The abundance of the microplastics	Size (mm)	References
Arctic	Sea-marine	Polyester, PAN, PA, and PVC-fibers (90%) and fragments	0.7 particles/m ³	1.00–2.00 (62%) and 2.00–5.00 (38%)	[108]
	Chukchi Sea, Western Ocean	PE (34.9%), PP (18.6%), PUR (16.3%), silicone resin (7.0%), PET and PVC (4.7%)	≤ 0.0188 pieces/m ²	1.0–2.0 (37.2%)	[109]
Switzerland, France, Germany, and Netherlands	River-freshwater	PS (29.7%), PP (16.9%), other types (13.6%), acrylate (9.3%), polyester (5.1%), and PVC (1.7%)—spherules (45.2%), fragments (37.5%), fibers (2.5%)	Average 5×10^{-5} particles/m ³	0.3–1.0	[110]
Open Ocean	North Eastern Mediterranean	Fibers (70%)	42 ± 46.7 particles/m ³	0.1–2.5	[111]
	Northwest Europe	Fragments (63%), film (14%), pellets (10%), foam (8%)	≤ 1.5 particles/m ³	1.00–2.79	[112]
	North East Pacific Ocean	Essentially fibers and fragments	9 180 particles/m ³	0.60 ± 0.22	[113]
	Southern Caspian Sea (surface water)	PP, PE, PS—fragments (38%), styro-foam (31%), film (20%)	3.449×10^{-8} particles/m ²	1.00–4.75	[114]
	North and Central Adriatic Sea (surface waters)	PE (80.7%), PP (9.3%), PA (2.6%), PS (1.9%) and PET (1.4%)—fragments, filaments, films, pellets and granulates	1.78×10^{-8} to 1.55×10^{-6} items/m ²	Not determined	[115]
	Arabian Sea, Indian Ocean	PE, PP, PA, and PVC-fiber (77.14%), fragment (21.78%), and pellet (1.07%)	Average 0.013 ± 0.002 particles/m ³	0.864 ± 0.074	[116]
	Northwestern Pacific Ocean, Indonesian seas	Nylon (15%), polyester (13%), polyvinyl alcohol (10%), PS (8%), PE (8%), PP (4%),	< 0.018 particles/m ³	0.20–0.35	[117]

PE—polyethylene; PP—polypropylene; PLA—polylactide; PA—polyamide; PS—polystyrene; PUR—polyurethanes; PVC—polyvinyl chloride; PET—polyethylene terephthalate; PAN—polyacrylonitrile; PMMA—polymethyl methacrylate; PTFE—polytetrafluoroethylene

Fig. 2 Pathways allowing microplastics to enter aquatic environments include industrial activities, the economic sector, and urban areas. These sources contribute to the contamination of aquatic environments with up to 125 trillion microplastic particles globally [4, 15, 120, 128, 129]. Aquatic ecosystems, such as freshwater bodies and oceans, are heavily impacted by microplastic-based pollution, primarily due to the inefficient wastewater treatment processes



Wang et al. [134] point to the concern for the lack of federal legislation (mostly state or city laws being enforced) on plastic products such as plastic bags or food containers, the lack of public investment for increased sustainability,

and the gap in environmental and societal education on this issue. Still, some states such as California, Maine, Hawaii, and cities like New York have laws that ban the use of plastic bags, highlighting California as the only state with more

stringent regulation for the control of microplastics in wastewater and drinking water [134]. Moreover, Sorensen et al. [135] warn that although there are government policies and laws controlling the abundance of microplastics in the environment and food products, they are ambiguous and mostly unenforceable.

According to Costa et al. [130], almost 150 countries around the world already have some policies that reduce plastic usage. On the Asian continent, Bangladesh was one of the first countries worldwide to ban low-density polyethylene (LDPE) single-use items from the market (since 2002), India with the same guidelines (essentially for low-density polyethylene (LDPE) bags) in 2016, and more recently China banned certain products made from plastic microbeads and plastic bags [124, 130]. Also, African, and South and Central American countries, *e.g.*, Rwanda, Chile, and Costa Rica, have government laws to ban the consumption of single-use products made from plastic [124, 130, 133].

Overall, it is urgent to reinforce the regulations and guidelines currently in force to enhance the monitoring of microplastics in wastewater. Despite all legislative efforts to regulate the production and usage of plastics worldwide, there is still a need to deal with a considerable number of other sources or pathways of microplastics entering wastewater that are not controlled, *e.g.*, domestic laundry water or run-off.

Microplastics in wastewaters

The wastewater environment is a significant source responsible for releasing microplastics into aquatic ecosystems, primarily through sludge, road runoff, and effluent discharges [17, 136, 137]. Microplastics stem from household items such as toothbrushes, cosmetics, cleaning products, personal care items, and synthetic fibers, among others [17, 138]. These particles vary in morphological and chemical properties and are consistently discharged in substantial quantities into receiving waters [17]. Reddy et al. [17] reported that the influx of microplastics into wastewater is influenced not only by household sources but also by pollutants from the industrial sector, human activities, demographic factors, and local conditions (Fig. 2). Plastic particles derived from construction materials, *e.g.*, insulation, medical equipment, tire wear, pipelines, automobiles, and the textile industry also contribute significantly to the considerable abundance of microplastics in wastewater. This is exacerbated by factors such as population density, societal habits, cultural practices, economic activities, and daily and seasonal weather fluctuations. The billions of plastic particles released daily into wastewater and other aquatic environments, eventually reach wastewater treatment plants (Table 3), depending not

only on anthropogenic and abiotic factors but also on the annual wastewater inflow (m^3/year) to the wastewater treatment plants [137].

Microplastics in wastewater have diverse shapes and sizes that are influenced by the wastewater influent sources, the material degradation processes, and wastewater discharge sources. Synthetic fibers, *e.g.*, polyesters (PES), polyethylene (PE), or polyamide (PA), dominate in wastewater treatment plants due to household laundry and textile production. However, particles in various forms such as pellets, fragments, foams, flakes, and films are also identified [120, 137, 139, 140]. Several studies estimate that synthetic fibers constitute 40% to 89% of the microplastics found in wastewater, highlighting household washing as a major contributor [120, 137, 138, 140, 141]. These particles are typically smaller than 1 mm, often ranging between 0.15 and 0.33 mm [120, 137, 138, 140, 141]. Wastewater hosts a wide array of plastic polymers, including polypropylene (PP) ($\approx 21\%$), polyethylene (PE) ($\approx 23\%$), polyethylene terephthalate (PET) ($\approx 7\%$), polyamide (PA) ($\approx 2\%$), polystyrene (PS) ($\approx 13\%$), polyvinyl chloride (PVC) ($\approx 7\%$), polyurethanes (PUR) ($\approx 5\%$), polyvinyl alcohol (PVAL) ($\approx 3\%$), polyesters (PES) ($\approx 2.5\%$), and acrylic ($\approx 10\%$), frequently detected in both influent and effluent samples across wastewater treatment plant facilities [137, 139–141]. These polymers exhibit varying physicochemical characteristics independent of geographical location, wastewater treatment plant types and capacities, biotic and abiotic transport processes, and discharge sources (Table 3).

Common techniques for microplastic removal in wastewater treatment plants

The presence of high amounts of microplastics in wastewater poses substantial environmental risks, not only because of the microplastics themselves but also due to their potential to adsorb various other pollutants [142, 143]. Owing to their hydrophobic surface properties, microplastics can interact with and adsorb heavy metals, *e.g.*, copper, zinc, lead, iron, chromium, cadmium, arsenic, persistent pollutants, and pharmaceuticals and personal care products (PPCPs). This includes harmful contaminant additives like polybrominated diphenyl ethers, phthalate esters, or polychlorinated biphenyls, originating from their chemical synthesis, raising concerns about the effective removal of these microparticles from aquatic environments [141–148]. Recent studies have also investigated the interaction between microplastics and oil in aquatic ecosystems, revealing evidence of hetero-agglomeration between plastic particles and oil seeps or slicks on ocean water surfaces [149–151].

Table 3 Microplastics wastewater samples, with the respective microplastics concentrations, particle size, and their removal efficiency

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Germany	WWTPs	Primary, Secondary, and Tertiary treatments	PE, PP, PS, PA, PVC, PUR, PET, ethylene vinyl acetate, polyvinyl alcohol, acrylonitrile butadiene styrene, polylactide, and paint	Not mentioned	> 500 µm: 0 to 50 microplastics/m ³ ; < 500 µm: 10 to (9 × 10 ³) microplastics/m ³	97.0	< 0.5 or ≥ 0.5	[204]
	WWTPs	Primary, Secondary, and Tertiary treatments – consist of a mechanical treatment stage, a biological stage with denitrification and nitrification, and chemical phosphorus precipitation with aluminum-based precipitants	PE (39.7%), PP (28.9%), PET (21.6%), and PS (9.8%)	8.8 × 10 ² ± 2.7 × 10 ² microplastics/m ³	7.0 ± 2.2 microplastics/m ³	99.2 ± 0.29	< 5.0	[168]
Thailand	WWTPs	A conventional WWTPs and an underground WWTPs coupled with ultrafiltration	PET, PE, and PP (60% was fibers shape)	conventional – 0.01655 ± 0.00992 microplastics/m ³ and with UF – 0.0077 ± 0.00721 microplastics/m ³	conventional – 0.00352 ± 0.00143 microplastics/m ³ and with UF – 0.01067 ± 0.003.51 microplastics/m ³	Ultrafiltration–96.97% and Conventional–86.14%	0.05–0.5	[177]
	DWTP	Primary and Secondary treatments – (with chlorination)	PP (20%), PE (30–55%), PET (15–40%) – fragments (90%), fibers (10%)	7.5 × 10 ⁴ particles/m ³	6.5 × 10 ⁴ particles/m ³	27.7	0.05–1.0	[205]
	WWTPs	WWTP–A is a cyclic activated sludge system; WWTP–B is an oxidation ditch; and WWTP–C is a CAS system, which undergoes continuous operation	PET (18%), PE (13%), polyacrylate (12%), PP (9%) – fibers (57–61%), also fragments and sheet	0.0122 pieces/m ³	0.0020 pieces/m ³	84.0	Not mentioned	[115]
Italy	Conventional municipal WWTPs	Primary, secondary, and pilot-scale tertiary treatment facility, with ultrafiltration with a pore size of 0.1 µm	PET (40%), PE (29%), PP (10%), PMMA (6%) and PS (3%)–Fibers, films and fragments	0.077 ± 0.007 particles/m ³	0.0023 ± 0.0015 particles/m ³	Conventional treatment–86.14%; and ultrafiltration–96.97%	0.05–0.5	[206]
	Municipal WWTPs	Primary, secondary, and tertiary treatments (full-scale plant with CAS system versus UASB + anaerobic membrane bioreactor	PE (43–53%), PP (10–13%), polyesters (9–10%), PUR (9%), PS (4%), PA (2%), and PVC (2%)–fragments (27–36%), lines (9%) and films (55–73%)	3.6 × 10 ³ microplastics/m ³ –CAS and 0.00172 microplastics/m ³ –UASB + AnMBR	5.2 × 10 ⁴ microplastics/m ³ –CAS and 0.0002 microplastics/m ³ –UASB + AnMBR	Conventional process–86%; UASB + AnMBR–94%	0.1–0.5	[183]
	WWTPs	Primary, secondary, and tertiary treatments–articulated in screening, grit and grease removal stages, biological treatment, sedimentation (with recycled activated sludge), sand filter treatment, and disinfection	Polyesters (35%) and PA (17%)–films (73%), fragments (21%), and lines (6%)	2.5 × 10 ⁴ ± 3.0 × 10 ⁴ microplastics/m ³	4.0 × 10 ⁴ ± 1.0 × 10 ⁴ microplastics/m ³	84.0	< 1.0	[178]

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Scotland	WWTPs	Primary, Secondary, and Tertiary treatments	PP (most abundant synthetic polymer–23%) in (fibers shape (67%))	8.1×10^8 particles/day	2.2×10^7 particles/day	97.0	0.060 and 2.8	[205]
	Secondary WWTPs	Four stages of treatment (grit and grease effluent, primary effluent, and the final effluent)	Alkyds (28.7%), polystyrene–acrylic (19.1%), polyester (10.8%), PUR (8.9%), and PMMA (8.3%) (flakes (67.3%), fibers (18.5%), film (9.9%), beads (3.0%))	$0.0157 (\pm 0.00523)$ microplastics/ m^3	$2.5 \times 10^{-4} (\pm 4.0 \times 10^{-5})$ microplastics/ m^3	98.41	average 0.598 (± 0.089)	[136]
France	WWTPs	Primary, Secondary, and Tertiary treatments	PS (37.5%) and PE (39.7%), PP, PVC, PET, PA, PUR (3–17%)—fibers and fragments	0.244 items/ m^3	0.00284 items/ m^3	98.83	0.2–0.5	[207]
	Four WWTPs with ultrafiltration as tertiary treatment	WWTP–A—the main process is MBR ($0.02 \mu m$ UF); WWTP–B–MBR ($0.1 \mu m$ UF); WWTP–C—Physico-chemical with biofiltration; WWTP–D—activated sludge	PE, PA	$4.203\text{--}42.00$ microplastics/ m^3	Not mentioned	WWTP–A and B–88–99.8% WWTP–C–98–99.9% WWTP–D–72%	Not mentioned	[208]
WTTP		Primary, Secondary, and Tertiary treatments (with screening, then grit and oil removal, settled in a primary settling tank before undergoing biological treatment that consists of biofilters)	Fibers ($\geq 90\%$)	$260\text{--}320 \times 10^3$ particles/ m^3	$14\text{--}50 \times 10^3$ particles/ m^3	84.38–94.62	0.1–5.0	[81]
USA	Tertiary WWTPs	Primary, Secondary, and Tertiary treatments	Not mentioned	0.001 particle/ m^3	8.8×10^{-7} microplastics/ m^3	99.9	0.1–0.6	[209]
	WWTPs	Primary and Secondary treatments	Essentially PE and PMMA ($\approx 70\%$)—microfibers,	$0.1\text{--}0.25$ Microplastic / m^3	<0.027 Microplastic / m^3	75 to 99	$0.060\text{--}0.418$	210
WWTPs		Primary, Secondary, and Tertiary treatments (development of AnMBR)	Fibers (55–62%), fragments (22–26%), and microbeads (11–16%)	0.133 ± 0.0356 microplastics/ m^3	<0.005 microplastics/ m^3	AnMBR–99.4%, while in the other two WWTPs–61–73%	0.58	[182]
Municipal WWTPs		Different types of WWTPs facilities, populations, and advanced filtration types	Fibers (59%), followed by fragments (33%), films (5%), foams (2%), and pellets (1%)	Not mentioned	4.0×10^{-6} to 1.95×10^{-1} microplastics/ m^3	Not determined	$0.125\text{--}0.355$ (57%)	[211]
DWTPs		Primary, Secondary, and Tertiary treatments (Coagulation through aluminum and iron—salts, sedimentation, and ultrafiltration)	PE	Not mentioned—synthetic solution	Not mentioned	45.34 ± 3.93	<0.5	[212]
Conventional WWTPs		Primary and Secondary treatments (Sedimentation and Aerated grit chamber)	PA (54.8%), PE (9.0%), PP (9.6%), PVC (2.5%), and polycarbonate (1.0%)—fibers (34–57%) and fragment (31–46%)	0.0799 microplastics/ m^3	0.0284 microplastics/ m^3	64.4	$0.02\text{--}0.3$ (45–80%)	[213]

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Municipal WWTPs	WWTPs	Three stages of treatments (with anaerobic–anoxic–oxic and MBR or cyclic activated sludge technology and fiber rotary filter)	PE (34.95–46.26%), PP (31.07%), PET (22.45%), phenolic resin (18.45%), and PS (15.53%) – fragments (34–38%) and granulates (28–29%)	$1.03\text{--}2.205 \times 10^{-2}$ microplastics/ m^3	$2.4\text{--}3.4 \times 10^{-4}$ microplastics/ m^3	97.67–98.46	<0.7	[188]
	Municipal WWTPs	Primary, Secondary, and Tertiary treatments (oxidation ditch versus MBR)	PET (47%), PS (20%), PE (18%), and PP (15%)–fragments (65%), fibers (21%), which mainly were PET, with limited films (12%) and foams (2%)	$2.8 \times 10^{-4} \pm 2 \times 10^{-5}$ microplastics/ m^3	$5 \times 10^{-5} \pm 1 \times 10^{-5}$ microplastics/ m^3	99.5% in the membrane system versus 97% in oxidation ditch	0.0625–0.5	[181]
	Urban WWTP	Primary treatment includes a course, thin, cyclone, and aeration grit; Secondary treatment includes the sequencing batch reactor activated sludge process, and CAS; and tertiary treatment includes high-efficiency sedimentation tanks, activated sand filters, denitrification filters, and disinfection tanks	PP, PE, PMMA, PET, and PVC–fragment, fiber, and thin film	0.0325 ± 0.001 items/ m^3	$5.0 \times 10^{-3} \pm 4.0 \times 10^{-4}$ items/ m^3	84.6	Mainly 0.03–0.10	[214]
Municipal sewage treatment plants	Until Tertiary treatment (include an aerated grit chamber, primary sedimentation tank, and secondary sedimentation tank following treatments (anaerobic, anoxic, and aerobic), and denitrification, ultra-filtration, ozonation, and ultraviolet)		PET (42.25%), PES (19.09%), and PP (13.05%) – more than 70% of microplastics	0.1203 ± 0.0013 items/ m^3	$5.9 \pm 2.2 \times 10^{-4}$ items/ m^3	95.16 ± 1.57	Average 0.682 ± 0.529	[187]
Wastewater	Electrocoagulation–anode material made of aluminum and iron metal	PE, PMMA, cellulose acetate, and PP (granulates)	Synthetic solutions using controlled concentration of microplastics–0.05; 0.10; 0.20; 0.50; 0.80; 1.00 g/L			91–99.9	0.0063–0.287	[173]
Freshwater	Tertiary treatments (a novel mechanism based on electrostatic–force–induced aggregation and flotation)	PE, PS, PVC, and fiber mixture	0.026 \pm 0.006 microplastics/ m^3	Not mentioned	90.2–95.7	<1.0		[174]
WWTPs (only filtration system)	Dynamic membrane filtration and transmembrane pressure	Not determined	Synthetic solutions using controlled concentration of microplastics–0.1; 0.20; 0.40; 0.60; 0.80; 1.00 g/L			Not determined	<0.09	[215]
Advanced DWTPs	Coagulation and flocculation combined with sedimentation, sand filtration, and filter combined with ozonation	PET (55.4–63.1%), PE (15.1–23.8%), PP (8.4–18.2%), and polyacrylamide–fiber, spheres, and fragments	6.614 ± 11.32 microplastics/ m^3	$\approx 0.930 \pm 72$ microplastics/ m^3	82.1–88.6	0.001–0.010 (85%)		[175]

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Spain	WWTPs	Four stages of treatment	LDPE (52.4%), PP (11.3%), HDPE (9.0%) (fragments (46.9%), films (34.0%), beads (11.5%), fibers (7.4%), and foam (0.2%))	3.20×10^{-3} ($\pm 6.7 \times 10^{-4}$) microplastics/m ³	3.1 (± 0.6) $\times 10^{-4}$ microplastics/m ³	90.3	0.4–0.6	[166]
	DWTPs	Primary, and secondary treatments in urban and industrial waste	PVC, HDPE, PE	0.64503 ± 0.18224 microplastics/m ³ to 1.56749 ± 0.41318 microplastics/m ³	0.01640 ± 0.00785 microplastics/m ³ to 0.13135 ± 0.09536 microplastics/m ³	91.0–97.0	0.100–0.355	[216]
	Urban WWTP	Constituted by a pre-treatment (sieving, grit remover, degreaser), primary treatment, and secondary treatment (biological reactor, gravity thickener, floatation, anaerobic stabilization, dehydration, and co-generation)	PE (8–79%), synthetic cellulose (0–26%), PP (0–39%), PVC (0–17%), synthetic polymeric resin (0–91%), PA (0–21%), PAN (0–13%) and polyester (0–12%)–fragments (42%), fibers (44%) and films (10%)	0.365 to 1.058 microplastics/m ³	0.032 to 0.178 microplastics/m ³	96–99	0.02–2.0	[217]
Municipal WWTP	Primary and secondary treatments		Polyester fibers, PE, dyed cotton, PP and cellophane fibers, PET (51% were fragments and 49% fibers)	0.171 ± 0.042 particles/m ³	0.0107 ± 0.0052 particles/m ³	94.0	0.025 – 0.104 (54%)	[178]
Municipal WWTP	Primary, secondary (one with oxidation ditch) and the other with integrated fixed film activated sludge, and two tertiary treatments		PE (30%), PET (29%), PP (11%), and PS (7%)–fragment, foam, film, fiber, and granulate	0.0155 ± 0.0035 particles/m ³ and 0.0385 ± 0.0025 particles/m ³	$2.0 \times 10^{-4} \pm 1.0 \times 10^{-4}$ particles/m ³ and $1.0 \times 10^{-4} \pm 1.0 \times 10^{-4}$ particles/m ³	With oxidation – 90% and with integrated fixed film activated–97%	0.05 – 0.2 (41%) and 0.2 – 0.5 (27%)	[218]
DWTPs	Primary, and Secondary treatments (with chlorine dioxide and carbon dioxide are added as initial disinfection, followed by the coagulation and floatation step, and ozonation followed by granular activated carbon filtering. The advanced membrane technology		PE and PP – fibers (45–60%) and fragments (40–55%)	$9.6 \times 10^{-4} \pm 4.6 \times 10^{-4}$ microplastics/m ³	$6 \times 10^{-5} \pm 4 \times 10^{-5}$ microplastics/m ³	93 ± 5	0.02 – 0.50	[184]
UK	WWTP	Primary, secondary, and tertiary treatment processes (primary settlement, activated biological anoxic treatment, and activated biological aerobic treatment)	PE (50%), PVC (42%), nylon (4%), and PP (4%)	0.034 microplastics/m ³	0.0015 microplastics/m ³	76.9	0.392 (± 0.027)	[219]
Synthetic wastewater	Electrocoagulation treatment–electrode made of aluminum		PE	Synthetic solution using a controlled concentration of PE– 0.10 g/L		99.24	0.3 – 0.355	[172]
Treatment Plants	A wide range of applied techniques, until tertiary processes		PS and acrylonitrile butadiene styrene	0.0049 microplastics/m ³	1.0×10^{-6} microplastics/m ³	99.99	≥ 0.025	[220]

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Denmark	Conventional WWTP	First, Secondary treatments (applying several levels of filtration through biofilters)	PE dominated the particle numbers (34%), followed by PP, PVC, PS, PA, and PMMA. PUR was rare and below 1%	917 items/m ³	≈ 200 items/m ³	79.0	< 0.3	[221]
	WWTPs	First, Secondary and Tertiary treatments	Acrylic (27%), PE (27%), PP (39%)	7,216 microplastics/m ³	0.054 items/m ³	98.0	0.010–0.50	[222]
	DWTPs	First, Secondary and Tertiary treatments	PA, PVC, PS, PE, PU, PP, PES—fibers (19%)	Detected 174 ± 405 microplastics/m ³			0.008–0.374	[223]
Hong Kong	WWTPs	Until tertiary treatments, including chemical—enhanced primary treatment	Fibers 55–71%; fragments 25–26% and sheet 19%	1.01–2.06 × 10 ³ microplastics/m ³	2.7–4.0 × 10 ⁴ microplastics/m ³	60.4–86.9	1.0–5.0 (≈ 60–64% of the particle)	[224]
Netherlands	Tap Water	Primary and secondary treatments (by conventional or membrane reactor)	Fibers (98.7%), film (2.2%)	Quantification – An average of 2.18 × 10 ³ ± 1.65 × 10 ⁴ particles/m ³			0.05–4.83	[225]
	Wastewater		Fibers, spheres, and foils	0.068–0.910 particles/m ³	0.009–0.091 particles/m ³ (average of 0.052 particles/m ³)	72.0	0.010–5.0	[180]
Finland	Municipal WWTPs	Primary and secondary treatments	PE and PET (79.1%), PE (11.4%), PA (3.7%), and PP (< 0.1%)	0.0576 (± 0.0124) microplastics/m ³	0.0004 (± 0.0001) microplastics/m ³	99.4	< 1.0	[167]
	Secondary WWTPs	Primary and secondary treatments (include coagulation and flocculation (using ferric chloride, polyaluminum chloride, and cationic polyamine)	PS—spheres	5.0 × 10 ² microplastics/m ³ –182 × 10 ³ microplastics/m ³	0.5 × 10 ³ –13.2 × 10 ³ microplastics/m ³	for 0.001 mm–95.0%; and for 0.0063 mm–76.0%	0.001–0.0063	[171]
	Wastewater	Membrane bioreactor; rapid sand filter; dissolved air flotation; disc filter	PE (≈ 60%), followed by PE (≈ 14%), polyacrylates (av. 7%), PVC (≈ 5%), PS (≈ 4%), and PP (≈ 3%)	6.9 × 10 ³ (± 1.0 × 10 ³) microplastics/m ³	5.0 × 10 ⁶ (± 4.0 × 10 ⁶) microplastics/m ³	MBR—99.9%; RSF—97%; DAF—95%; DF—98.5%	0.020–0.100	[169]
Israel	WWTPs	Primary, Secondary, and Tertiary treatments (consist of bar screens, grit removal, primary clarifiers, biological nutrient removal, activated sludge reactors, final clarifiers, effluent filtration, and disinfection)	PE (13.9%), PVC (5.9%), PP (2.0%), polycarbonate (1.5%), PFTE (1.0%), polyolefin elastomer (0.5%), PS (0.5%), PUR (0.5%) and nylon 66 (0.5%)—most of the fragments are fibers (74%)	0.065 microplastics/m ³	0.00197 microplastics/m ³	97.0	0.2 to 0.45 (with a peak abundance at 0.35)	[226]
Australia	Municipal WWTPs	Primary and advanced secondary treatments, which include screening (mesh size of 3 mm) and sedimentation, biological treatment, flocculation, disinfection and de-chlorination processes, ultrafiltration, reverse osmosis, and de-carbonation	PE (10%), PET (80%), PP (10%)	WWTP A—0.092 microplastics/m ³ , WWTP B—0.098 microplastics/m ³ and WWTP C—0.055 microplastics/m ³	WWTP A—0.00096 microplastics/m ³ , WWTP B—0.00098 microplastics/m ³ and WWTP C—0.00091 microplastics/m ³	WWTP A—97.6%, WWTP B—90.2%, and WWTP C—80.4%	0.10–0.19	[185]

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Republic of Korea	WWTPs	Tertiary treatments—used inorganic coagulant followed by ozone (WWTP-A); membrane disc-filter (WWTP-B) and rapid sand filtration (WWTP-C)	Microbeads, fibers, fragments, and sheet	4.2–31.4 microplastics/m ³	0.033–0.297 microplastics/m ³	WWTP-A–99.2%; WWTP-B–99.1%; WWTP-C–98.9%	Not mentioned	170
	Municipal WWTP	Primary, Secondary, and Tertiary treatments, <i>e.g.</i> , most of them apply anaerobic–anoxic–aerobic or modified treatment processes	PP (39.6%), PE (25.6%), PET (21.3%), also PS, PMMA, PA, PUR, and polyether–fragments (68.2%), fibers (31.8%)	0.010 to 0.470 items/m ³	4 × 10 ^{−6} to 5.1 × 10 ^{−4} items/m ³	98.7 to 99.99 (in 31 WWTPs)	0.045–5.00	186
	WWTPs	Until Tertiary treatments, with first physical processes (bar screening, grit removal, and primary clarifier), biological processes, and finally advanced treatment (coagulation and disc filter)	Microbeads, fibers, and fragments	2.223–10.044 microplastics/m ³	0.029–0.447 microplastics/m ³	98–99	Not mentioned	227
Sweden	WWTPs	Primary, Secondary, and Tertiary treatments – mechanically, chemically, and biologically (conventional process)	Fibers, fragments, and flakes	15.1 ± 0.89 × 10 ³ particles/m ³	8.25 ± 0.85 particles/m ³	≈ 99.9	≥ 0.3	100
Canada	Secondary WWTPs	Primary and Secondary treatments (applying oil extraction followed by digestion and vacuum filtration)–Sedimentation	89.2% of synthetic fibers—made of PS, PES, and PA—fibers (65.6%), fragments (28.1%), and pellets (5.4%), while a small abundance of foam, granules, and sheets (0.22%, 0.45% and 0.20%, respectively)	0.032 ± 0.007 microplastics/m ³	5.0 × 10 ^{−4} ± 2.0 × 10 ^{−4} microplastics/m ³	98.3	< 1.0	228
Czech Republic	Freshwater	Magnetic extraction with coated iron–nanoparticles	PE, PS, PUR, PVC, and PP	Synthetic solution using controlled concentrations of microplastics—0.005, 0.010 particles/L		84.0	< 0.2–1.0	229
	DWTPs	Primary, Secondary, and Tertiary treatments—coagulation and flocculation, flotation, sand filtration, and granular activated carbon filtration	PS (≤ 68%), PP (16–26%), PE (24%), PAN, polybutylacrylate, PVC, and PMMA (≤ 5%)	1.473 ± 0.034 to 3.605 ± 0.497 particles/m ³	0.338 ± 0.076 to 0.628 ± 0.028 particles/m ³	70.0	0.001–0.01	230
India	DWTPs	Primary and Secondary treatments—chlorination, coagulation, pulse clarification, and sand filtration	PE (33%), PET (56%), PP (4%), PS (3%)—fibers (52–59%), films/fragments (41–48%)	0.0179 items/m ³	0.0028 items/m ³	85.39	0.025–0.10	176

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Iran	WWTP	Primary and Secondary treatment processes, including screening (mesh size of 8 mm), grit chamber, and primary sedimentation, followed by aeration, secondary sedimentation, and chlorination process	fibers (73%), film (9%), and granule (17.6%)	9.2×10^{-4} microplastics/m ³	8.4×10^{-4} microplastics/m ³	90.9	< 0.84	²³¹
	Conventional WWTPs	Primary and Secondary treatments—its main units are a bar screen, an aerated grit chamber, a flow metering, a primary settling tank, an anoxic tank, an aeration basin, a clarifier, and a disinfection unit	PET and polyester (73–91%), PMMA, PA, PE, and PP (10%)—essentially fibers	12,667 microplastics/m ³	423 ± 44.9 microplastics/m ³	96.7	< 0.5	²³²
	Municipal WWTP	Not mentioned	Not mentioned	Average of 0.0545 ± 0.0377 microplastics/m ³	$1.8 \times 10^{-3} \pm 2.0 \times 10^{-3}$ microplastics/m ³	96.7	0.025–0.3	²³³
Poland	Municipal WWTP	Consists of a mechanical part, a biological part, and a sludge treatment line with a biogas installation	PET (44%), PE (14.15%), PP (8.56%), PS (2.93%), and PUR (2.69%)—fragments (33%), film (12.3%), foam (8.31%) and beads (3.42%)	4.09×10^{-3} particles/m ³	2.5×10^{-4} ($\pm 4.0 \times 10^{-5}$) particles/m ³	> 90	< 1.0 (50.12%) and 0.1–0.2 (30.81%)	²³⁴
Slovenia	Municipal Lab-scale WWTP	Primary (mechanical and biological)—sequencing batch reactor	Microbeads from cosmetics	Synthetic solution with an Microplastic concentration of 5×10^{-4} g/m ³	21 particles/m ³	52.0	≈ 0.1	²³⁵
Turkey	WWTPs	Primary and Secondary treatments	PP (13.8%), PET (50%), PE (29%)—fibers (55–88%), film (2.5–18%), fragment (10–27%)	Seyhan WWTP (4 825 697 ± 575 630 microplastics per day); Yüreğir WWTPs (351 019 ± 26 WWTP (2 040 639 ± 343 859 microplastics/day)	Seyhan WWTPs (1 249 102 ± 146 373 microplastics per day); Yüreğir WWTPs (351 019 ± 26 800 microplastics/day)	Yüreğir WWTPs (79.0%) and Seyhan WWTPs (73.0%)	< 5.0	²³⁶
	Municipal WWTP	Primary and Secondary treatments—including screens, aerated grit chambers, and conventional activated sludge	PE (54.5%), EVA (37.84%)—essentially fibers (41.78–60.77%)	4.80 particles/m ³	0.14 particles/m ³	97.08	0.1–0.5 (58.57–80.07%)	²³⁷
	WWTPs	Primary, Secondary and Tertiary treatments—including anaerobic oxidation and membrane bioreactor	PE (28%) and PET (28%), PA (21%), PP (9%), PVC (8%) and PES (6%)—fiber, fragment and fiber	0.040 to 0.079 particles/m ³	0.0004 to 0.0056 particles/m ³	Membrane bioreactor (99%); trickling filter (98%) and activated sludge process (91%)	0.5–1.0	²³⁸
Vietnam	Four Municipal WWTPs	Primary, Secondary, and Tertiary treatments (including CAS treatment, a sequencing batch reactor coupled with a UV channel, and trickling filters followed by aerated lagoons and maturation ponds)	Fragments and fibers	1860 items/m ³ to 125 000 items/m ³	140 items/m ³ and 813 items/m ³	69–99	Average 1.0	²³⁹

Table 3 (continued)

Country	Sampling	WWTPs treatments	Most types of microplastics (shape)	Abundance in influent	Abundance in effluent	Efficient removal (%)	Size (mm)	References
Indonesia	Municipal WWTP	Primary and Secondary treatments-including Moving Bed Biofilm Reactor and a High Rate Clarifier	fibers (68–70.17%), fragments (23.68–26.24%), films (0.71–2.9%), microbeads (0.4–1.4%) and foams (1.4–2.8%)	0.017 (± 0.006) items/m ³	0.0014 (± 0.0001) items/m ³	Setiabudi WWTP–91.29	0.1–5.0	²⁴⁰
Bangladesh	3 DWTPs	Primary, Secondary, and Tertiary treatments	PP (48%), PE (35%), PET (11%), and PS (6%)–fragments were (90%), fibers (10%)	DWTP–A and DWTP–B– $2.6 \times 10^2 \pm 1.0 \times 10^{-2}$ items/m ³ and DWTP–C– $6.2 \times 10^3 \pm 1.6 \times 10^{-3}$ items/m ³	DWTP–A and DWTP–B– $3.0 \times 10^4 \pm 3.0 \times 10^5$ and DWTP–C– $4.0 \times 10^4 \pm 1.0 \times 10^5$ – $5.0 \times 10^4 \pm 2.0 \times 10^4$ items/m ³	Up to 99%	0.02–5.0	²⁴¹
Lithuania	Secondary WWTP	Primary and Secondary treatments	PP, PE, and PS, being essentially fibers and fragments	0.08 to 0.34 particles/m ³	0.012–0.025 particles/m ³	55.4 \pm 3.9	0.02–1.0	²⁴²
Fiji	Two municipal WWTPs–Kinoya and Natabua	Primary and Secondary treatments (including simple secondary clarifier, trickling filter, and Anaerobic pond)	Cellophane, PP, polyester, and polyvinyl acetate–ethylene–fibers and fragments (both 44%)	Kinoya– $3.45 \pm 0.3 \times 10^{-3}$ particles/m ³ Natabua– $2.9 \pm 1.1 \times 10^{-3}$ particles/m ³	Kinoya – $0.3 \pm 0.26 \times 10^{-3}$ particles/m ³ Natabua– $0.53 \pm 0.42 \times 10^{-3}$ particles/m ³	Kinoya–91% Natabua–81% (33%)	0.25 (67%) and 2.0 (33%)	²⁴³
Marocco	Urban and Industrial WWTPs	One WWTP, utilized lamellar decantation and a submarine emissary, and the other used activated sludge	PE (29–20%), PET (21–24%), PP (10–8%), PVC (11–9%), PS (9–12%), PA (10–6%), ethylene–vinyl acetate (8–13%), and HDPE (2–8%)–fibers, fragments, films, and pellets	1.114 \pm 0.090 particles/m ³	0.607 \pm 0.101 particles/m ³	46	0.1–0.5	²⁴⁴
Kazakhstan	Municipal WWTP	Constituted with mechanical treatment, primary and secondary sedimentation, biological treatment (Oxidation process), subsequent coagulation, floatation, granular filtration, and UV disinfection	PE, PET, PS, PVC, PA, PP, polyester, and ethylene–acrylic acid copolymer–fibers, fragments, and films	0.0471 \pm 0.0376 microplastics/ m ³ to 0.0694 \pm 0.0400 microplastics/ m ³	0.0041 \pm 0.0031 microplastics/ m ³ to 0.0054 \pm 0.0035 microplastics/ m ³	Between 88.6–93.0	0.1–0.5 (66.45–75.13%)	²⁴⁵
Oman	Three Municipal WWTPs	Until tertiary treatments–WWTP–A (with CAS); WWTP–B (Sequencing Batch Reactor); WWTP–C (with MBR)	PP, LDPE, PUR, PET, and PVC–fiber (60.7%)	WWTP–A– 3.6 – 8.5×10^{-3} particles/m ³ , WWTP–B – 4.6 – 8.7×10^{-3} particles/m ³ , and WWTP–C– 3.6 – 7.1×10^{-3} particles/m ³	WWTP–A– 1.0×10^{-3} particles/m ³ , WWTP–B– 1.4×10^{-3} particles/m ³ , and WWTP–C– 1.0×10^{-3} particles/m ³	WWTP–A (82.5%), WWTP–B (77.4%), and WWTP–C (79.2%),	Average 0.045–0.425	²⁴⁶
Nepal	WWTPs	Primary and Secondary treatments (including secondary aeration tank)	Essentially PE and PS–fiber, fragments, foam, and pellets	0.0312 \pm 0.0173 particles/m ³	$8.50 \times 10^{-3} \pm 5.60 \times 10^{-3}$ particles/m ³	72.5	Most common 0.15–5.0	²⁴⁷
Kuwait	Three municipal WWTPs	Primary, Secondary, and Tertiary treatments (including reverse osmosis, ultrafiltration membranes, an oxidation ditch system for secondary treatment, sand filtration, UV, and chlorination as tertiary treatment)	PA (\approx 55%), PP (\approx 30%), and PE (\approx 15%)–fiber (78–84%), fragment (13–20%), and films (2–4%)	0.119 to 0.230 particles/m ³	0.001 to 0.012 particles/m ³	81.3%	Essentially 0.063–0.333	²⁴⁸

Regarding the potential adsorption of microplastics with toxic heavy metals and other emerging contaminants, recent research indicates an increased risk to human health, as these particles may enter the food and water supply chain [13, 142, 146, 147, 152, 153]. The aggregation of microplastics in various environmental media (freshwater, soils, and so on) highlights their role as vectors for the persistence of other emerging contaminants [144]. Moreover, the prevalence of microplastics in agricultural drainage water, in groundwater supplies, and in drinking and tap water [12, 13, 119, 154–159] directly impacts human health. Several polymers have been detected in human tissues or blood [30, 160, 161]. Certain microplastics, *e.g.*, polypropylene (PP), polyethylene (PE), polyethylene terephthalate (PET), and polyamide (PA) have also been reported in bottled water, whether in plastic, glass, or cardboard bottles [13, 154, 162–164]. Hence, the efficient removal of microplastics from wastewater is crucial, despite the lack of high-efficiency treatments in wastewater treatment plants or drinking water treatment plants (DWTPs) (Table 3).

In some conventional or municipal wastewater treatment plants, performing a primary, secondary, and often advanced tertiary treatment is necessary to remove microplastics efficiently [137, 165]. Pretreatment processes can remove between 35 and 59% of the plastic particles in conventional wastewater treatment plants, but, when combined with primary treatment, are responsible for the removal of the majority of microplastics (estimated between 50 and 98%), with only the smallest particles remaining in the wastewater [137, 165]. Secondary treatment yields an additional removal of 0.2–14%. Finally, tertiary treatments can contribute to a high-efficiency elimination of the smallest particles [137, 165].

In recent years, numerous techniques have been developed or improved in wastewater treatment plants to remove microplastics, depending on the design of the wastewater treatment plant and its location or population density (Table 3). These techniques include grit chamber and primary sedimentation [136, 166, 167], precipitation [168], dissolved air flotation [169], coagulation [170, 171], electrocoagulation [172, 173], electrostatic forces [174], sand filtration [175], rapid (gravity) sand filter (RSF) [170, 176], ultrafiltration [177], activated sludge process [178, 179], membrane bioreactor (MBR) [167, 169, 180–183], reverse osmosis [184, 185], anaerobic-anoxic-aerobic (A²O) treatments [186–188], and ozonation [175]. The efficiency of wastewater microplastics removal (Table 3) is very uneven, with lower rates obtained in conventional wastewater treatment plants [189], contrasting with high-efficiency removal rates of 99.9% obtained in tertiary wastewater treatment plants that have implemented new methodologies, *e.g.*, implemented MBR [169, 186]. Chemical precipitation, *e.g.*, with aluminum, ultrafiltration, and A²O treatments can

also eliminate the microplastics most commonly found in wastewater, with removal percentages of 99.2%, 97%, and 98.5–99.9%, respectively. It should also be noted that electrocoagulation treatment can be another alternative method (91–99.24%), but the studies performed so far have been on synthetic solutions with controlled microplastics concentrations [172, 173, 190].

On the other hand, in terms of biological treatments, MBR is considered one of the methods that can lead to high microplastics removal yields (between 94 and 99.9%). For instance, Talvite et al. [169] achieved 99.9% efficiency of the treatment compared to other methodologies applied in wastewater treatment plants, such as RSF (97%), dissolved air flotation (DAF, 95%), and disc filter (98.5%). Moreover, by comparing the performance of MBR against two conventional wastewater treatment plants, a 99.4% yield was obtained [182]. A cost-effectiveness analysis by Vuori et al. [190] concluded that treating wastewater by MBR to mitigate microplastics pollution provides higher cost savings in both private and social costs, compared to the implementation of conventional activated sludge (CAS) or RSF, which have a lower removal efficiency (88% to 97.1%, respectively) and a higher environmental impact [190]. Even so, despite proving to be an efficient method for lower-capacity wastewater treatment plants (with tertiary treatments implemented), the incorporation of MBR into tertiary treatment in wastewater treatment plants entails prohibitive energy costs and difficulties in controlling fouling. Moreover, there are gaps related to the low cleaning efficiency of the membranes, the failure to obtain ecological status, and, above all, the absence of specificity for microplastics removal [191–193].

Overall, the methods currently used in wastewater treatment plants have several limitations, *e.g.*, process costs, modest removal efficiency, lack of environment-friendly status, and the absence of specificity regarding the microplastics problem, boosting the necessity of developing new technologies and approaches. As such, efforts are underway to enhance existing water treatment processes and develop new, cost-effective, and efficient technologies for sustainable water treatment. These efforts focus on using advanced materials and technologies such as protein nanofibrils, metal–organic frameworks (MOFs), covalent organic frameworks, nanocomposites, fluorinated oligoamide nanorings, and bio-based materials (such as agricultural residues, algae, bacteria, and fungi) [194, 195].

Given the necessity for sustainable and cost-effective methods to remove microplastics from aquatic environments, the use of microorganisms has gained attention in the scientific community for biodegradation and retention of microplastics present in wastewater, considering their abundance in wastewater [196, 197]. Biostimulation and bioaugmentation, mainly involving fungi and bacterial species, are being explored as bioremediation strategies for microplastics.

These bioremediation approaches are influenced by a wide range of biotic (the species of microorganisms and the number of living cells, the interactions in microbial consortia, cell hydrophobicity, and enzymatic diversity) and abiotic factors (UV light, pH, temperature, and oxygen (O_2) concentration, among others) [198, 199].

Some microorganisms can biodegrade microplastics in aquatic environments by breaking down the complex polymerized structure of plastics into smaller molecules (into their monomers), through the secretion of intracellular or extracellular enzymes, such as polyhydroxybutyrate (PHB), poly(3-hydroxybutyrate-co-3-hydroxyvalerate) (PHBV), or poly(3-hydroxyalkanoate) (PHA), among others [200]. This biodegradation process involves the first stage where biofilms form on the plastic fragment, followed by biofragmentation which involves the secretion of enzymes by the microorganisms surrounding the plastic particles [201, 202]. The third step is the assimilation (polymer degradation to monomers) and culminates in the mineralization process, with the synthesis of secondary metabolites such as carbon dioxide (CO_2), water (H_2O), and methane (CH_4) [198–200, 203]. For instance, Miloloža et al. [199] underlined that biodegradation of microplastics by microorganisms depends on many factors, such as the metabolic versatility of microorganisms, environmental factors, and the physicochemical characteristics of the polymers that constitute the plastics (Table 1).

The concentration of microplastics entering municipal wastewater treatment plants ranges from very low to extremely high concentrations—from 2.8×10^{-4} microplastics per m^{-3} to very high concentrations of 55×10^4 microplastics per m^3 (Fig. 2). The abundance of plastic particles in the influent is influenced by the type and capacity of wastewater treatment plant, but more importantly, by the population density (Table 3). In the effluent, the concentrations are much lower, ranging from 1.0×10^{-6} microplastics per m^{-3} to $14\text{--}50 \times 10^3$ microplastics per m^{-3} , depending on the treatment stages (pretreatment, primary, secondary, and tertiary processes) that yield the overall efficiency of the method. The morphology (shape and size) as well as PE–polyethylene; PP–polypropylene; PLA–polylactide; PA–polyamide; PS–polystyrene; PUR–polyurethanes; PVC–polyvinyl chloride; PET–polyethylene terephthalate; PAN–polyacrylonitrile; PMMA–polymethyl methacrylate; PTFE–polytetrafluoroethylene; LDPE–low-density polyethylene; HDPE–high-density polyethylene; EVA–Ethylene–vinyl acetate; WWTPs–wastewater treatment plants; DWTPs–drinking water treatment plants; CAS–conventional activated sludge; AnMBR–anaerobic membrane bioreactor; MBR–membrane bioreactor; UF–ultrafiltration; UV–ultraviolet; UASB–pilot-scale up flow granular sludge blanket; WWTPs–A, B, C or D–different WWTPs in the same city or country the type of plastic polymer also affects the

performance of the remediation methods. The fibers are the most common shape detected in wastewater treatment plant systems, and the sizes of microplastics are on average less than 0.5 mm, being polyethylene (PE), polypropylene (PP), polystyrene (PS), polyesters (PES), and polyethylene terephthalate (PET) as the most commonly detected polymers (Table 3). As such, the currently implemented wastewater treatment processes are not properly designed to remove microplastics, showing a lack of specificity for microplastics structures and shapes [137, 249]. These lack specificity for microscopic-sized particles, making them environmentally unfriendly and economically unsustainable due to their reliance on complex multi-step processes and chemical treatments, as reviewed by several authors [17, 176, 249, 250]. Moreover, the energy used in conventional wastewater treatment operations is remarkable (represents an estimation of 2% to 4% of a country's electricity consumption) [196, 251]. Given the current limitations in removing microplastics from wastewater, alternative processes, such as the use of microalgae-based systems, have gained attention as more sustainable and efficient options [252–254].

Microalgae potential for microplastic removal

While numerous well-explored studies focus on the advantages of using microorganisms for the bioremediation of microplastics in soils and aquatic environments, predominantly marine, they primarily concentrate on employing various species of bacteria, some fungi, and other larger eukaryotic organisms for the biodegradation of plastic particles [198, 200, 255]. Several studies have emerged exploring the potential of these microorganisms for biological treatment within wastewater treatment plants, targeting heavy metals, organic pollutants, and, more recently, microplastics [165, 196, 256, 257]. For instance, Nyika et al. [196] discussed the role of specific microorganisms, mainly bacteria and fungi, in wastewater bioremediation, to eliminate heavy metals, dyes, or persistent organic pollutants. The authors covered general factors that influence these treatments, such as the physicochemical bioavailability of contaminants, the nature of the contaminant (chemical structure, concentration, solubility, and toxicity), environmental and microbial properties (enzyme induction, level of mutation), and co-metabolism versus growth substrate (associated to predation, succession, and competition during microbial interaction). In the same line, Tang and Hadibarata [202] discussed the potential and limitations of applying bioremediation as a secondary or tertiary treatment in wastewater treatment plants, but with the main purpose of microplastics removal. That study focused on bacteria, e.g., *Vibrio*, *Bacillus*, *Acetobacteriodes*, *Saccharomonospora*, *Rhodococcus*, and *Pseudomonas*, and

fungi, *e.g.*, *Penicillium*, *Aspergillus*, and *Fusarium*, with the potential to degrade polymers, exhibiting efficiencies ranging between 8.46% and 36.4%, depending on the type of plastic, the conditions employed, and microbial species [202]. Though, these processes are time-consuming, more than 40 days are necessary before any degradation or removal of these polymers occurs [202].

In addition to bacterial and fungal species, microalgae possess a high potential for the removal and biodegradation of microplastics in wastewater [258]. When compared to bioremediation systems relying on bacteria, microalgae offer several advantages in wastewater treatment. Microalgae exhibit remarkable adaptability to their environment, efficiently removing nutrients, displaying high seasonal tolerance, and showcasing superior metabolic performance concerning nutrients commonly found in wastewater [202, 251, 256]. Furthermore, microalgae generate less sludge and demonstrate significantly better tolerance to other emerging contaminants, such as pharmaceuticals and personal care products [125, 251, 259]. The exploration of microalgae-based systems for wastewater biological treatment is novel and has the potential to yield environmentally friendly and sustainable processes [254]. These systems yield high-value end products and use relatively straightforward microorganisms to integrate into existing wastewater systems (Fig. 3) [251].

Microalgae are a diverse category of photosynthetic microorganisms, either eukaryotic or prokaryotic, which can use the sunlight that illuminates the surface of wastewater as an energy source, coupled with the capture of atmospheric carbon dioxide (CO₂) for their basic metabolism and growth [254, 261, 262]. Active green (*Chlorophyta*), red (*Rhodophyta*), and brown (*Phaeophyta*) algae are commonly identified as belonging to the category of photosynthetic microorganisms, as well as cyanobacteria [263]. These microorganisms are metabolically flexible with photoautotrophic, mixotrophic, or heterotrophic metabolism and can adapt to a variety of water environments, even those with extreme growth conditions (*i.e.* highly saline water) [258, 264]. Furthermore, the biomass derived from microalgae in wastewater holds substantial potential for diverse applications in sustainable product development. It can serve as a source for extracting high-value compounds like proteins, polysaccharides, fatty acids, and vitamins, among others, catering to various biotechnological uses [254, 263, 265]. For instance, in the green industry, this biomass can contribute to the production of biofuel or biogas. Additionally, it finds utility as feed for fish and animals, biofertilizers, food ingredients, bioplastics, cosmetics, and biopharmaceutical drugs, and aids in carbon dioxide (CO₂) mitigation, promoting industrial sustainability and embracing the concept of a circular economy with its “zero-waste” concept (Fig. 3) [254, 263, 265, 266]. Hence, microalgae not only extract


nutrients (such as nitrogen and phosphorus) from wastewater for their growth and metabolism but also exhibit remarkable resilience to highly toxic wastewater conditions. This versatility highlights microalgae as efficient biological systems capable of effectively treating various types of effluents within wastewater treatment plants.

Moreover, microalgae have a high potential to remove different types of emerging contaminants found in wastewater [267], including heavy metals (by adsorption and uptake), persistent organic pollutants (by bioaccumulation, biosorption, and metabolization), and controlling pathogen growth (by increasing pH and temperature, and excreting growth inhibitory metabolites) [254] and more recently microplastics [200, 268]. As such, microalgae-based biological treatment systems offer lower capital and operational costs, short growth cycles, ease of operation, and environmentally friendly processes (Fig. 3) [258, 269, 270].

More recently, a novel biotechnological approach has emerged, focusing on studying microalgae capacity to stimulate the synthesis of specific enzymes, *e.g.*, polyethylene terephthalate (PET) hydrolase (PETase), superoxide dismutase, mono(2-hydroxyethyl) terephthalic acid hydrolase (MHETase), cutinases, esterases, and lipases, among others, for plastic decomposition and disposal. These enzymes can be further genetically modified for that purpose [201, 271–276]. Synthetic enzymes represent a promising application in the bioremediation of microplastics, potentially playing a pivotal role in the removal or elimination of microplastics from wastewater [201].

On the other hand, microalgae, with their uncomplicated genetic structure, are amenable to genetic manipulation to produce and secrete enzymes responsible for breaking down or eliminating microplastics [201]. This genomic and post-genomic approach, known as Algomics, is highly important for understanding algal cellular mechanisms [201, 277]. *C. reinhardtii* and *Phaeodactylum tricornutum* are microalgae that, despite inhabiting different aquatic environments, can express specific enzymes to degrade microplastics by being genetically manipulated [273, 275]. Kim et al. [273] demonstrated that *C. reinhardtii* strain CC-124 was able to synthesize a PETase, which is responsible for hydrolyzing polyethylene terephthalate (PET). After a period of 4 weeks of catalytic activity of PETase, through the interaction between *C. reinhardtii* and the microplastic, it was possible to detect the release of terephthalic acid, one of the metabolites of polyethylene terephthalate (PET) polymer degradation, a process which was corroborated by scanning electron microscopy (SEM) inspections that confirmed morphological changes on the surface of the microplastic. This was the first report showing the success of a microalga-based PETase in microplastic degradation. Moreover, in Moog et al. [275], *P. tricornutum* was used as a microbial cell factory to produce and secrete engineered PETase under

Fig. 3 “Zero waste” concept where microalgae-based bioremediation is evolving as a promising strategy for removing emerging contaminants, including microplastics, from wastewater. This approach offers advantages over traditional approaches like low costs, fast growth, and eco-friendly processes. Microalgae use the contaminants as a carbon source necessary for microalgal growth, resulting in high-value biomass or compounds (*i.e.* vitamins and lipids). By incorporating these systems into wastewater treatment, the industry can move closer to the zero-waste concept, promoting both sustainability and improving water quality [adapted from Srimongkol et al. [260]]



saltwater mesophilic conditions. The approach consisted of adapting the gene sequence encoding PETase^{R280A} (previously sequenced in the bacteria *Ideonella sakaiensis* [278]) and transferring it to *P. tricornutum* (PETase^{R280A}-FLAG) to study its biodegradation of polyethylene terephthalate (PET) and the copolymer polyethylene terephthalate glycol. After 5 weeks of incubation of the microplastic fragments with the marine diatom, it was observed that the enzyme was actively biodegrading the polymer, and products from the polyethylene terephthalate (PET) substrate degradation process, TPA and mono(2-hydroxyethyl) terephthalic acid, were detected. This study proposed that microalgae can be an alternative model for the synthetic biology of polyethylene terephthalate (PET) degradation enzymes compared to bacteria from aquatic environments [275].

The metabolic engineering of microalgae to produce specific enzymes presents a promising approach to develop proficient microorganisms for microplastics removal from aquatic environments. Other studies assessing the potential of microalgae to transform microplastics include the use of *Spirulina* sp. which was able to synthesize extracellular enzymes to degrade or adsorb polyethylene terephthalate (PET) and polypropylene (PP) microparticles [276, 279], while *C. vulgaris* was found to be able to degrade bisphenol-A, an additive compound present in polycarbonate,

polyacrylates, and polyesters [280, 281], without estrogenic activity [271, 276].

Selecting microalgae species for wastewater treatment

The selection of suitable microalgae species for wastewater treatment involves crucial criteria, particularly their ability to grow in a diversity of process conditions and to form harmless algal blooms [282]. Wastewater is a complex mixture of various substances that can vary widely in composition and typically includes organic matter, inorganic compounds (such as metals), nutrients (namely nitrogen and phosphorus), and other pollutants, *e.g.*, personal care and pharmaceutical products, and plastics [254, 283]. As such, the wastewater complex composition can be toxic to microalgae and/or unsuitable for their growth [260, 263].

Out of the thousands of known microalgae strains (approximately 300,000), the initial screening process should be focused on selecting strains that predominantly thrive in freshwater environments and can adapt to growth in the wastewater [284]. Also, microalgae require nutrients (namely nitrogen and phosphorus) and specific oligoelements, *e.g.*, iron, manganese, copper, and zinc, to replicate [254] and these are readily available in wastewater. However,

high concentrations of heavy metals can be present in wastewater which can inhibit microalgae growth [285]. For example, copper, when present at high concentrations in wastewater, can significantly decrease the biomass concentration, chlorophyll, and carotenoid content [286]. In fact, according to a toxicological study performed by Cid et al. [287], concentrations of copper above 0.1 mg/L in aquatic environments can be toxic to microalgae, reducing their growth by 50%, and at concentrations higher than 1.0 mg/L, it can inhibit completely their growth. Also, other factors characteristic of wastewater have to be considered since different microalgae have specific process requirements: pH levels (with ideal values ranging between 6 and 9, but some species survive in an acidic pH), organic matter availability, irradiation duration, and intensity (directly related to photosynthesis), the concentration of atmospheric carbon dioxide (CO₂), turbulence, and the water temperature [254, 263].

An essential criterion in choosing strains for phytoremediation processes is avoiding the formation of harmful algal blooms (HABs) [254, 288]. According to the Environmental Protection Agency (EPA) [289], the excessive abundance of nitrogen and phosphorus (the main nutrients) in the receiving water can lead to eutrophication, causing a rapid overgrowth of microalgae in a short time. This overgrowth consumes significant amounts of oxygen (O₂) in the environment, making it inhospitable for other life forms to survive. In some cases, there is a production of chemicals, metabolites, or toxic compounds that are harmful to aquatic life and human health (through contaminating the drinking water), causing significant negative impacts on the environmental health, ecosystems, and economy [290]. Therefore, microalgae species must be selected based on their harmless to the surrounding environment character and incapability to form HABs [254, 288]. Considering these criteria, microalgae typically tested for wastewater bioremediation belong to the genera *Chlorella*, *Scenedesmus*, *Desmodesmus*, *Neochloris*, *Nitzschia*, *Cosmarium*, and *Chlamydomonas* [254].

Toxicological profile

While microalgae can endure harsh conditions in wastewater [254, 265, 291], it is crucial to consider the influence of microplastics on their growth and metabolic activity. A wide variety of studies analyzing the microplastics toxicity and ecotoxicity have been done involving different microalgae species, including freshwater strains (as environmental model organisms), with different types and concentrations of microplastics [142, 274, 292–301]. According to Prata et al. [298], microplastics have negative impacts on microalgae metabolism, photosynthesis, and morphology. In fact, microplastics can decrease the growth of some microalgae, potentially inhibiting growth when high concentrations of microplastics or nanosized particles are present [296].

For instance, the presence of polystyrene (PS) decreased the growth of *Scenedesmus obliquus* (about 2.5% at 1000 mg/L), *Pseudokirchneriella subcapitata* (concentrations between 1 and 100 mg/L), *C. reinhardtii* (an EC₅₀ of 0.54 mg/L), *Chlorella pyrenoidosa* (until 22 days) [298], and *Chlorella vulgaris* (26.9% at 35 particles/L) [301]. According to Rani-Borges et al. (2021), 78 days after contact with these emerging contaminants (PS, polypropylene (PP), polyvinyl chloride (PVC), polyethylene (PE), polyethylene terephthalate (PET), and polyamide (PA)), growth was reduced in most microalgae species, e.g., *P. subcapitata*, *C. reinhardtii*, *Chlamydomonas pyrenoidosa*, and *Scenedesmus obliquus*. Their interaction can also lead to a subsequent decrease in chlorophyll content and photosynthetic activity (affecting the photosynthetic efficiency), either by acting as inhibitors of microalgae growth, or by the shadow effect they can cause in the aquatic environment (by decreasing the amount of light falling on the microalgae), or by the potential to decrease the expression of photosynthetic genes [201]. It is worth noting the capacity of polystyrene (PS) to decrease the polysaccharide content of extracellular polymeric substances (EPS) per cell of *C. vulgaris*, its total CYP450 content, and total CYP450 activity, as well as completely downregulating the expression on key genes related to algal cell metabolic activity, e.g., *psbA*, *psaB*, and *rbc* [301]. Also, microplastics can exert adverse effects on different stages of the photosynthesis process in microalgae—from the electron donor site to the electron transport chain, leading to oxidative stress. At the morphological level, severe consequences can occur, e.g., including unclear pyrenoids, plasma detached from the cell wall, deformed thylakoids, and cell wall thickening, and modulated energy metabolism [201, 296, 298]. Similarly, polystyrene microplastics were responsible for a 40% decrease in chlorophyll content in *S. obliquus* (at a concentration ≥ 100 mg/L [302]) and *C. pyrenoidosa* (up to 10 days, at a concentration of 50 mg/L) [292]. Other microplastics have been studied for their ecotoxicological impact on microalgae, including polypropylene (PP), high-density polyethylene (HDPE), polyvinyl chloride (PVC), polyethylene terephthalate (PET), polyethylene (PE), and polyacrylonitrile (PAN) [296, 299, 303, 304] which were found to inhibit growth, reduce photosynthetic activity, affect enzymatic activity, and alter cell morphology [296, 299, 303, 304].

The toxicological effects of microplastics on microalgae depend on a diversity of factors such as the size and shape of the plastic particles, polymer type, exposure time, and even the surrounding environment [298, 299]. These toxicological studies show that smaller particles tend to be more toxic to microalgae, due to the capacity for adsorption to the cell surface [296]. It is noteworthy that the toxicological impacts on microalgae may be temporary, with an initial high vulnerability followed by adaptive responses to the adverse

conditions [296, 305]. The hypothetical mechanisms for their detoxification can be via homo-aggregation or hetero-aggregation to microplastics as well as the development of biofilms on the surface of the microplastics to enhance nutrient use in the aquatic environment [296, 298, 305].

Therefore, choosing the most suitable microalgae for specific bioremediation treatments to eliminate microplastics from wastewater demands a thorough consideration of this toxicological profile and the aforementioned selection criteria. This selection process requires an understanding of how these microorganisms interact with the surfaces of microplastics, along with a keen awareness of the toxic effects these pollutants may have on the microalgae species selected.

Microalgae-microplastics interaction


Regarding the removal of microplastics, an important characteristic of these microorganisms when interacting with the surface of microplastics (Fig. 4) is the ability to form hetero-aggregates, by secreting extracellular polymeric substances, as part of the biofouling formation process [296, 306]. Manzi et al. [201] found that the most common mechanisms to remove microplastics using microalgae entail colonization, aggregation, and adsorption processes. Aggregation is the most plausible mechanism for application to wastewater treatment plants since it entails reduced adverse effects on the microalgae [201]. These authors further verified the ability of microalgae to aggregate on microplastics to form hetero-aggregates and the ability to colonize the surface of the microplastics inducing changes in the chemical structure of the plastic and initiating the removal process by aggregation [201]. Moreover, Cunha et al. [307] reported the interaction between freshwater microalgae (*Microcystis paniformis* and *Scenedesmus* sp.) and microplastics, and found that microalgae were able to colonize microplastics of polystyrene (PS) and polymethyl methacrylate (PMMA), leading to the formation of hetero-aggregates. *C. reinhardtii* was able to form aggregates on polypropylene (PP) microparticles after 20 days of exposure [303]. However, no hetero-aggregate formation occurred on high-density polyethylene (HDPE) surfaces. These authors hypothesized that microalgae can produce different types of extracellular polymeric substances depending on the type of polymer, resulting in higher affinity and cohesion for certain microplastics [303]. Other studies also support the high aggregation ability of freshwater microalgae to the surface of diverse microplastics, such as low-density polyethylene (LDPE) [308], polystyrene (PS) [268, 309], polymethyl methacrylate (PMMA) [309], and polylactide (PLA) [309].

Regarding the removal of plastics from wastewater, Kumar et al. [310] and Sarmah et al. [311] investigated the effectiveness of different species of photosynthetic

microorganisms (such as *Scenedesmus dimorphus*-a green microalga, *Anabaena spiroides*-a blue-green alga, and *Navicula pupula*-a diatom) to degrade low-density polyethylene (LDPE) and high-density polyethylene (HDPE). These three photosynthetic microorganisms were more prone to form hetero-aggregates on the surface of low-density polyethylene (LDPE) fragments than on the high-density polyethylene (HDPE) surface, with *A. spiroides* being the most prone of the three species to colonize low-density polyethylene (LDPE) [310]. Through SEM analysis and mass balance of the polyethylene (PE) fragments, the highest biodegradation efficiency was achieved by *A. spiroides* ($8.18 \pm 0.07\%$), followed by *N. pupula* contributing ($4.44 \pm 0.82\%$) and *S. dimorphus* ($3.74 \pm 0.26\%$). However, a higher biodegradation performance of low-density polyethylene (LDPE) was obtained by the two cyanobacteria, *Phormidium lucidum* and *Oscillatoria subbrevis* [311]. After 42 days of exposure, a 30% weight loss of low-density polyethylene (LDPE) was estimated by Fourier-transform infrared spectroscopy (FTIR) analysis, demonstrating a decrease in intensity of some characteristic peaks of polyethylene (PE), e.g., 2364 cm^{-1} , after being incubated with these two cyanobacteria. With complementary thermogravimetric-differential scanning calorimetry (TGA–DSC) analysis and carbon/hydrogen/nitrogen elemental analysis, it was possible to verify that in the particles degraded the total carbon percentage was reduced (3%–4%) and the melting point of the polymer decreased [310]. This could mean that the microplastics are used as a carbon source for their growth and metabolic activity, leading to the biodegradation of the polymers. *Chlorella* sp. in a consortium with a cyanobacterium also showed the potential to degrade low-density polyethylene (LDPE) bag particles [308]. The microalgae can colonize, aggregate, and adhere to the surface of low-density polyethylene (LDPE) microparticles, with support from *Cyanobacteria* sp. through the production of extracellular polymeric substances that facilitates the adhesion of *Chlorella* sp. Through field-emission scanning electron microscopy-energy dispersive X-ray (FESEM-EDX) analysis, complemented with FTIR, carbon analysis, and differential scanning calorimetry (DSC), Govindan et al. [308] estimated that approximately 38% of the amount of carbon decreased after hetero-aggregation of the consortium of these two microorganisms.

For the same type of microplastic source, the green microalgae *Uronema africanum* Borge also demonstrated to be of interest to biodegrade these polymers in a laboratory-scale context [313]. Although the biodegradation efficiency of microplastics in the 30-day trial (by carbon proportion) was not estimated, atomic force microscopy (AFM) and SEM analyses, along with gas chromatography-mass spectrometry (GC–MS) and FTIR, revealed that the microalgae can initiate the colonization and degradation of low-density polyethylene (LDPE) bags. In overview, this aggregation

Fig. 4 Three mechanisms used by microalgae to remove microplastics: bioadsorption, biodegradation, and bioaccumulation. Through microalgae metabolism, several metabolites can be formed by microplastic degradation (*i.e.* microplastic monomers, including propylene, phenol, and ethylene) along with high-value products such as lipids, vitamins, proteins, and polysaccharides [adapted from Rempel et al. [312]]



process enables the microalgae to act as a flocculant, causing precipitation or sedimentation of microplastics in the water system [306, 314]. So, from a treatment point of view, this strategy may hold the potential to be applied in wastewater treatment plants to remove microplastics, causing the microplastic-algal flocks to sediment and subsequently be removed through simple physical techniques, *e.g.*, settling.

Other removal mechanisms of emerging contaminants from wastewater by microalgae-based systems include bioaccumulation, bioadsorption, biodegradation, photolysis, hydrolysis, and volatilization [265, 269, 270, 282, 315]. Concerning bioremediation, three process pathways are feasible: bioadsorption, biodegradation, and bioaccumulation (Fig. 4). Bioadsorption, occurs when the contaminant is adsorbed to the cell wall or other substances excreted by the cell via interaction with the surface of the contaminant

(a physical and passive process) [265, 269, 270]. For this biological process, the chemical structure of the contaminant is a key factor, but other aspects like temperature, solubility, and pH can influence microalgae interaction [315].

Nolte et al. [316] studied the adsorption of microplastics to *P. subcapitata*, where atomic force microscopy (AFM) and depletion measurements demonstrated that positively or neutrally charged polystyrene (PS) microplastics were adsorbed more efficiently than negatively charged microplastics, mainly due to particle–particle (inter-particles) and particle–cell membrane interactions. In addition, both the physicochemical properties of the microplastics and environmental conditions (the cation concentration in the freshwater media) were found to be crucial factors involved in the absorption of the microalgae to the microplastic [316]. Another work was also developed to study the adsorption of

polystyrene (PS) microplastics to *Fucus vesiculosus* [317]. The adsorption of polystyrene (PS) microplastics was highly efficient (circa 95% of the particles remained adsorbed). *F. vesiculosus* has phaeophycean hairs, rich in alginate, that allow an easy adhesion of polystyrene (PS) microplastics [317]. Peller et al. [318] used *Cladophora* sp. to remove synthetic textile microfibers (MFs) and other microplastics (namely polyethylene (PE), polyethylene terephthalate (PET), and polyacrylonitrile (PAN)) from freshwater (in Laurentian Great Lakes), observing high rates of MFs adsorption to the alga [318]. This phenomenon was attributed to the chemical composition of the cell wall and the high surface area. More recently, Kiki et al. [319] reported the adsorption of polyethylene (PE), polyvinyl chloride (PVC), and polyamide (PA) microplastics to *C. vulgaris* over a 30-days exposure period, observing that the microalga colonized the microplastics and induced structural damage, fractures, and aging. A decrease in intensity of characteristic peaks was observed, e.g., 3300 cm^{-1} , 1641 cm^{-1} , and 686 cm^{-1} for polyamide (PA); 1245 cm^{-1} and 1093 cm^{-1} for polyvinyl chloride (PVC); 2920 cm^{-1} and 1463 cm^{-1} for polyethylene (PE) [319]. That study proposes that the aging of microplastics can alter the physicochemical properties of the polymers, resulting in decreased hydrophobicity, which enhances the adsorption of organic micropollutants from microplastics to the cell wall of *C. vulgaris* [319]. Compounds such as bisphenol-A, bisphenol-S, bisphenol-F, and parabens (propyl-paraben, PPBen, and benzyl-paraben, BPBen), among others, form hydrogen bonds to strengthen the microalgae multicellular structure, particularly the extracellular polymeric substances matrix [319]. It is noteworthy that when the experiments were submitted to a river freshwater environment, biofouling by algae considerably increased the adsorption capacity of polyethylene (PE) and polyvinyl chloride (PVC) and weakened the adsorption of the amide group on polyamide (PA) [319].

Moreover, a combination of mechanisms can also occur. According to Cheng et al. [309], microplastics removal by microalgae can be stimulated by two mechanistic pathways: i) hetero-aggregation between *S. abundans* and microplastics, promoting the formation of extracellular polymeric substances that stimulates mutual aggregation; ii) microplastics adsorption by the cell wall of microalgae. Comparing the two mechanisms, hetero-aggregation coupled with prolonged exposure to microplastics (enhancing the synthesis of binding extracellular polymeric substances) in the medium, stimulates the removal of microplastics compared to the adsorption approach, a finding that was corroborated by SEM analysis. However, in the same study, the PLA micro-particles were only efficiently removed in the water system by short-period adsorption to the microalgae cell wall in the

presence of extracellular polymeric substances. This means that the ability of microalgae to remove microplastics, as well as the associated mechanisms (including the production of specific extracellular polymeric substances), depends on the type of plastic the microalgae will try to adapt to and the period of exposure to the contaminant [309].

On the other hand, bioaccumulation, unlike bioadsorption, is an active process and consists of the transport of the emerging contaminants into exclusively living cells, crossing the cell wall, and binding to intracellular proteins or other cell substances [265, 270]. Maia et al. [269] argue that bioadsorption and bioaccumulation processes are not sustainable solutions for emerging contaminants removal, because the contaminants remain inside the living microalgae cells, transforming them into possible toxic biomass, hindering the valorization into high-value products.

Lastly, biodegradation is considered the most promising pathway because it allows the fragmentation of emerging contaminants macromolecules into simple non-toxic, or less hazardous breakdown molecules via microalgae catalytic activity for their growth [265, 269, 270, 315]. Furthermore, it is a process that can occur intracellularly and also extracellularly (or a combination of both) through the secretion of ligninolytic enzymes or extracellular polymeric substances. In biodegradation, the emerging contaminants can be used as a carbon source or an electron donor or acceptor for algal metabolic activity, or for co-metabolism, in which the emerging contaminants are degraded by enzymes [265, 320] (Abdelfattah et al. 2023; Anand et al. 2023). It is worth noting that the metabolites resulting from biodegradation must be identified and quantified, as they may prevail in the environment even though the contaminant has been degraded [269].

Regarding the removal of microplastics by biodegradation, it was found in freshwater environments that the microalga *Spirulina* sp. can also biodegrade microplastics [321]. Hadiyanto et al. [321] despite the initial aim of evaluating the toxicological effects of polyethylene (PE) and polypropylene (PP) microparticles on *Spirulina* sp., found by FTIR and energy dispersive X-ray analysis that the total carbon content after 30 days of experimentation had decreased by 1.62% and 1.08% in the polyethylene (PE) and polypropylene (PP) particles, respectively. There was a decrease in the intensity of some peaks in the spectrum of the microplastics of polyethylene (PE) and polypropylene (PP) that were in contact with the microalgae (the most evident at 3281 cm^{-1}). By complementary SEM analysis, it was also visualized that cracks have been formed on the surface of the microplastics by interaction with the microalgae [321]. *Spirulina* sp. was also evaluated in the biodegradation efficiency of polyethylene terephthalate (PET) and polypropylene (PP)

microparticles after 112 days [279]. After the long period of interaction between the microplastics and the microalgae, it was observed by EDX analysis that the carbon content in the polyethylene terephthalate (PET) and polypropylene (PP) microplastics decreased by 48.61% and 36.7%, respectively. The biodegradation of these polymers by *Spirulina* sp. was also corroborated by FTIR analysis. The infrared spectra of the polyethylene terephthalate (PET) particles after contact with the microalgae were characterized by the decrease in intensity of the peaks in the band located between 1613 and 1505 cm^{-1} , associated with carbon (C=C) bonds. In polypropylene (PP) particles, new peaks appeared essentially between 1599–1534 cm^{-1} and 1731 cm^{-1} , which indicates the presence of by-products (carbonyl groups in esters and ketones) resulting from the polypropylene (PP) oxidation process. Morphological analysis was also carried out by Khoironi et al. [279] using SEM and tensile strength. Throughout the 112 days, the structure of the microplastics was losing its hardness and elasticity, observing the appearance of cavities and erosion on the surface of microplastics. *Scenedesmus abundans* was also used for efficiently removing polystyrene (PS), polymethyl methacrylate (PMMA), and poly(lactic acid) (PLA) microparticles, obtaining an average removal rate of 84% [309]. Among the three types of polymers, *S. abundans* showed the highest removal efficiency for polymethyl methacrylate (PMMA) microparticles (with a removal efficiency of 98%), while for polystyrene (PS) and PLA microplastics, the removal was 84% and 87%, respectively [309].

Indirect photolysis, hydrolysis, and volatilization are less-studied processes applied by microalgae in microplastics bioremediation treatments [304, 315]. Indirect photolysis can be applied to more resistant pollutants in the medium, through the production of reactive oxygen species (ROS) by the microalgae [315]. This process depends on the photosynthetic capacity of the microalgae, and the process pH and temperature [251]. Hydrolysis and volatilization, on the other hand, are much more application-specific, for example, in organic pollutants with ester or amide groups [322].

Overall, all these potential bioremediation mechanisms that can be applied to different types of water contaminants, as described above, are equally applicable to microplastics [267]. As such, microalgae can be implemented as a tertiary treatment, for wastewater polishing, since the water may contain less toxic chemicals and heavy metals at this stage, which does not threaten the microalgae metabolic activity in the wastewater environment and does not impact microplastics removal [201]. However, it will be necessary to further explore these microalgae-microplastics interactions in a wastewater context and understand the influence that biotic and abiotic factors, e.g., temperature, pH, concentration of organic matter, and nutrients, have on the biodegradability of microplastics.

Conclusion

Bioremediation processes are starting to be recognized as promising strategies to be applied to remove these emerging contaminants, specifically the use of microalgae-based systems, bridging the current gaps, e.g., low-efficiency rates, in municipal wastewater treatment plants. Microalgae demonstrate the ability to form hetero-aggregates with plastic microparticles. They possess mechanisms capable of degrading the polymeric molecules into simpler and safer compounds through enzymatic activity or extracellular polymeric substances secretion. This process results in biomass with high-value, contributing to the development of more environmentally friendly approaches, which will also result in a more sustainable industry. In fact, by applying microalgae-based systems in wastewater bioremediation, there is an opportunity to advance the concept of zero-waste, thereby improving the sustainability of ecosystems and water quality. However, additional in-depth studies will be necessary to identify the mechanisms of microplastics removal from wastewater by microalgae, whether through bioflocculation or biodegradation processes, taking into account the limitation of current scientific studies highlighting the removal capacity of these micropollutants in a wastewater context. It is crucial to consider the specific parameters characterizing wastewater, as they influence the metabolic function and growth of performance microalgae, as well as the microbiome present in the aquatic environment, which can facilitate hetero-aggregation and consortia between different microbial species. This comprehensive understanding of the interactions and potential mechanisms associated with the removal of microplastics from wastewater will contribute to the development of more effective and environmentally sustainable strategies for mitigating microplastics pollution.

Author contributions P.S: Conceptualization, writing—review and editing. K.A.K: review and editing; C.A.S: Conceptualization, Writing—review and editing. M.S: review, editing, and funding acquisition. All authors had revised the manuscript and approved the final version.

Funding This work was supported by national funds through FCT/MCTES (PIDDAC): LEPABE, UIDB/00511/2020 (DOI: 10.54499/UIDB/00511/20200) and UIDP/00511/2020 (DOI:10.54499/UIDP/00511/2020) and ALiCE, LA/P/0045/2020 (DOI: 10.54499/LA/P/0045/2020); and the grant attributed by Portuguese Foundation for Science and Technology–FCT—to Paulo Sousa (2022.14159.BD).

Data availability Not applicable.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Ethic approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

Code availability Not applicable.

References

- Ojha A, Mishra A, Tiwary D (2022) Microplastic in the aquatic ecosystem and Human health implications. In: plastic and microplastic in the environment, pp 49–62. <https://doi.org/10.1002/9781119800897.ch4>
- Frias JPGL, Nash R (2019) Microplastics: finding a consensus on the definition. *Marine Pollut Bull* 138:145–147. <https://doi.org/10.1016/j.marpolbul.2018.11.022>
- Virginia Janssens, Meysmans Anne (2023) Plastics the fast facts 2023 : a review on plastic contributions to European economy. <https://plasticseurope.org/de/knowledge-hub/plastics-the-fast-facts-2023/>
- OECD. Plastic waste in (2019) (2022). <https://doi.org/10.1787/a92f5ea3-en>
- Eriksen M, Cowger W, Erdle LM, Coffin S, Villarrubia-Gómez P, Moore CJ et al (2023) A growing plastic smog, now estimated to be over 170 trillion plastic particles afloat in the world's oceans—urgent solutions required. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0281596>
- GESAMP (2015) Sources, fate and effects of microplastics in the marine environment: a global assessment. London
- Zhang K, Hamidian AH, Tubić A, Zhang Y, Fang JKH, Wu C et al (2021) Understanding plastic degradation and microplastic formation in the environment: a review. *Environ Pollut* 274:116554. <https://doi.org/10.1016/j.envpol.2021.116554>
- Auta HS, Emenike CU, Fauziah SH (2017) Distribution and importance of microplastics in the marine environment: a review of the sources, fate, effects, and potential solutions. *Environ Int* 102:165–176. <https://doi.org/10.1016/j.envint.2017.02.013>
- An L, Liu Q, Deng Y, Wu W, Gao Y, Ling W (2020) Sources of microplastic in the environment. *Handb Environ Chem* 95:143–159. https://doi.org/10.1007/978-1-4939-9949-9_449
- Friot D, Boucher J (2017) Primary microplastics in the oceans | IUCN Library System. <https://portals.iucn.org/library/node/46622>
- Kavya A, Sundarajan S, Ramakrishna S (2020) Identification and characterization of micro-plastics in the marine environment: a mini review. *Mar Pollut Bull* 160:111704. <https://doi.org/10.1016/j.marpolbul.2020.111704>
- Samandra S, Johnston JM, Jaeger JE, Symons B, Xie S, Currell M et al (2022) Microplastic contamination of an unconfined groundwater aquifer in Victoria, Australia *Sci Total Environ* 802:149727. <https://doi.org/10.1016/j.scitotenv.2021.149727>
- Singh S, Trushna T, Kalyanasundaram M, Tamhankar AJ, Diwan V (2022) Microplastics in drinking water: a macro issue. *Water Supply* 22:5650–5674. <https://doi.org/10.2166/ws.2022.189>
- Cole M, Lindeque P, Halsband C, Galloway TS (2011) Microplastics as contaminants in the marine environment: a review. *Mar Pollut Bull* 62:2588–2597. <https://doi.org/10.1016/j.marpolbul.2011.09.025>
- Eriksen M, Lebreton LCM, Carson HS, Thiel M, Moore CJ, Borerro JC et al (2014) Plastic pollution in the world's oceans: more than 5 trillion plastic pieces weighing over 250,000 tons afloat at sea. *PLoS ONE* 9:1–15. <https://doi.org/10.1371/journal.pone.0111913>
- Eerkes-Medrano D, Thompson RC, Aldridge DC (2015) Microplastics in freshwater systems: a review of the emerging threats, identification of knowledge gaps and prioritisation of research needs. *Water Res* 75:63–82. <https://doi.org/10.1016/j.watres.2015.02.012>
- Reddy AS, Nair AT (2022) The fate of microplastics in wastewater treatment plants: an overview of source and remediation technologies. *Environ Technol Innov* 28:102815. <https://doi.org/10.1016/j.eti.2022.102815>
- Wu X, Zhao X, Chen R, Liu P, Liang W, Wang J et al (2022) Wastewater treatment plants act as essential sources of microplastic formation in aquatic environments: a critical review. *Water Res* 221:118825. <https://doi.org/10.1016/j.watres.2022.118825>
- Andrady AL, Neal MA (2009) Applications and societal benefits of plastics. *Philos Trans R Soc London Ser B, Biol Sci* 364:1977–1984. <https://doi.org/10.1098/rstb.2008.0304>
- Baur E, Osswald TA (2019) Rudolph, N. 2 - Introduction, 13–43
- Wiesinger H, Wang Z, Hellweg S (2021) Deep dive into plastic monomers, additives, and processing aids. *Environ Sci Technol* 55:9339–9351. <https://doi.org/10.1021/acs.est.1c00976>
- Plastics (2022) - the facts 2021 - An analysis of European plastics production, demand and waste data, 11 <https://plasticseurope.org/wp-content/uploads/2021/12/Plastics-the-Facts-2021-web-final.pdf>
- Hardin T (2021) Plastic: it's not all the same. *Plast. Ocean* 1. <https://plasticoceans.org/7-types-of-plastic/>
- Charmondusit K, Seeluangsawat L (2009) Recycling of poly(methyl methacrylate) scrap in the styrene-methyl methacrylate copolymer cast sheet process. *Resour Conserv Recycl* 54:97–103. <https://doi.org/10.1016/j.resconrec.2009.07.005>
- Selvaranjan K, Navaratnam S, Rajeev P, Ravintherakumaran N (2021) Environmental challenges induced by extensive use of face masks during COVID-19, a review and potential solutions. *Environ Chall*. <https://doi.org/10.1016/j.envc.2021.100039>
- De Tommaso J, Dubois JL (2021) Risk analysis on pmma recycling economics. *Polymers (Basel)*. <https://doi.org/10.3390/polym13162724>
- Chamas A, Moon H, Zheng J, Qiu Y, Tabassum T, Jang JH et al (2020) Degradation rates of plastics in the environment. *ACS Sustain Chem Eng* 8:3494–3511. <https://doi.org/10.1021/acssuschemeng.9b06635>
- For Biotechnology Information NC (2022) Polypropylene. PubChem. <https://pubchem.ncbi.nlm.nih.gov/compound/POLYPROPYLENE>
- Nghiem LD, Iqbal HMN, Zdarta J (2021) The shadow pandemic of single use personal protective equipment plastic waste: a blue print for suppression and eradication. *Case Stud Chem Environ Eng* 4:100125. <https://doi.org/10.1016/j.cscee.2021.100125>
- Geyer R, Jambeck JR, Law KL (2022) Production, use, and fate of all plastics ever made. *Sci Adv* 3:e1700782. <https://doi.org/10.1126/sciadv.1700782>
- HDPE and LDPE (2022) PubChem Subst Rec SID 319064333. <https://pubchem.ncbi.nlm.nih.gov/substance/57309676>
- Mallisetty R, Veluru S, Talib Hamzah H, Hamem MNK, Tukar-ambai M, Rao Poiba V et al (2023) Biodegradation of low density polyethylene (LDPE) by *Paenibacillus* Sp. and *Serratia* Sp. isolated from marine soil sample. *Mater Today Proc*. <https://doi.org/10.1016/j.matpr.2023.05.639>
- National Center for Biotechnology Information. Polyvinyl Chloride (2022) PubChem Compd. Summ. Poly(vinyl chloride). https://pubchem.ncbi.nlm.nih.gov/compound/Poly_vinyl_chloride
- Shashoua Y (2001) Degradation of plasticized PVC inhibiting the deterioration of plasticized poly(vinyl chloride) – a museum

- perspective contents (Ph.D. thesis). Denmark's Technical University, 136
35. Guo S-L, Chen B-L, Durrani SA (2020) Chapter 3 - Solid-state nuclear track detectors. Handbook of radioactivity analysis, 4th edn. Academic Press, pp 307–407
 36. Polystyrene (2015) Polym database 1. <https://pubchem.ncbi.nlm.nih.gov/compound/Polystyrene>
 37. Shah S, Ahmad I, Ishaq M (2017) Degradation study of used polystyrene with UV irradiation. Adv Mater Sci 2:1–6. <https://doi.org/10.15761/AMS.1000130>
 38. Bhuvaneswari G H (2018) 3 - Degradability of polymers. Recycling of polyurethane foams. Elsevier Inc. <https://doi.org/10.1016/b978-0-323-51133-9.00003-6>
 39. National Center for Biotechnology Information (2022) Ethylurea PubChem Compd Summ CID 12254, 1. <https://pubchem.ncbi.nlm.nih.gov/compound/Ethylurea>
 40. European Commission (2019) Innovative acrylic(PMMA- Polymethyl methacrylate) recycling technology complying with regulations. <https://cordis.europa.eu/project/id/856103>
 41. Methyl methacrylate (2022) PubChem Compd Summ CID 6658, 1. <https://pubchem.ncbi.nlm.nih.gov/patent/US-4108923-A>
 42. Poly(Acrylonitrile) (2016) Polym database, 1. <https://polymerdatabase.com/polymers/polyacrylonitrile.html>
 43. Andrady AL (2011) Microplastics in the marine environment. Mar Pollut Bull 62:1596–1605. <https://doi.org/10.1016/j.marpolbul.2011.05.030>
 44. Jiang J-Q (2018) Occurrence of microplastics and its pollution in the environment: a review. Sustain Prod Consum 13:16–23. <https://doi.org/10.1016/j.spc.2017.11.003>
 45. Li J, Liu H, Paul Chen J (2018) Microplastics in freshwater systems: a review on occurrence, environmental effects, and methods for microplastics detection. Water Res 137:362–374. <https://doi.org/10.1016/j.watres.2017.12.056>
 46. Woodall LC, Sanchez-Vidal A, Canals M, Paterson GLJ, Coppock R, Sleight V et al (2022) The deep sea is a major sink for microplastic debris. R Soc Open Sci 1:140317. <https://doi.org/10.1098/rsos.140317>
 47. Barnes DKA, Walters A, Gonçalves L (2010) Macroplastics at sea around Antarctica. Mar Environ Res 70:250–252. <https://doi.org/10.1016/j.marenvres.2010.05.006>
 48. Lusher AL, Tirelli V, O'Connor I, Officer R (2015) Microplastics in Arctic polar waters: the first reported values of particles in surface and sub-surface samples. Sci Rep 5(1):1–9. <https://doi.org/10.1038/srep14947>
 49. Obbard RW, Sadri S, Wong YQ, Khitun AA, Baker I, Thompson RC (2014) Global warming releases microplastic legacy frozen in Arctic sea ice. Earth's Future 2:315–320. <https://doi.org/10.1002/2014EF000240>
 50. Gallagher A, Rees A, Rowe R, Stevens J, Wright P (2016) Microplastics in the solent estuarine complex, UK: an initial assessment. Mar Pollut Bull 102:243–249. <https://doi.org/10.1016/j.marpolbul.2015.04.002>
 51. Gauquie J, Devriese L, Robbens J, De Witte B (2015) A qualitative screening and quantitative measurement of organic contaminants on different types of marine plastic debris. Chemosphere 138:348–356. <https://doi.org/10.1016/j.chemosphere.2015.06.029>
 52. Thompson RC, Olsen Y, Mitchell RP, Davis A, Rowland SJ, John AWG et al (2004) Lost at sea: where is all the plastic? Science 80(304):838. <https://doi.org/10.1126/science.1094559>
 53. Alomar C, Estarellas F, Deudero S (2016) Microplastics in the Mediterranean Sea: deposition in coastal shallow sediments, spatial variation and preferential grain size. Mar Environ Res 115:1–10. <https://doi.org/10.1016/j.marenvres.2016.01.005>
 54. Liebezeit G, Dubaish F (2012) Microplastics in beaches of the east Frisian Islands Spiekeroog and Kachelotplate. Bull Environ Contam Toxicol 89:213–217. <https://doi.org/10.1007/s00128-012-0642-7>
 55. Dubaish F, Liebezeit G (2013) Suspended microplastics and black carbon particles in the Jade system, Southern North Sea. Water Air Soil Pollut 224:1352. <https://doi.org/10.1007/s11270-012-1352-9>
 56. Stefánsson H, Peternell M, Konrad-Schmolke M, Hannesdóttir H, Ásbjörnsson EJ, Sturkell E (2021) Microplastics in glaciers: first results from the Vatnajökull ice cap. Sustain 13:4183. <https://doi.org/10.3390/su13084183>
 57. Rodrigues MO, Abrantes N, Gonçalves FJM, Nogueira H, Marques JC, Gonçalves AMM (2018) Spatial and temporal distribution of microplastics in water and sediments of a freshwater system (Antuã River, Portugal). Sci Total Environ 633:1549–1559. <https://doi.org/10.1016/j.scitotenv.2018.03.233>
 58. Antunes J, Frias J, Sobral P (2018) Microplastics on the Portuguese coast. Mar Pollut Bull 131:294–302. <https://doi.org/10.1016/j.marpolbul.2018.04.025>
 59. Zhu L, Bai H, Chen B, Sun X, Qu K, Xia B (2018) Microplastic pollution in North Yellow Sea, China: observations on occurrence, distribution and identification. Sci Total Environ 636:20–29. <https://doi.org/10.1016/j.scitotenv.2018.04.182>
 60. Di M, Wang J (2018) Microplastics in surface waters and sediments of the three Gorges reservoir. China Sci Total Environ 616–617:1620–1627. <https://doi.org/10.1016/j.scitotenv.2017.10.150>
 61. Zhou G, Wang Q, Zhang J, Li Q, Wang Y, Wang M et al (2020) Distribution and characteristics of microplastics in urban waters of seven cities in the Tuojiang River basin. China Environ Res 189:109893. <https://doi.org/10.1016/j.envres.2020.109893>
 62. Hu D, Zhang Y, Shen M (2020) Investigation on microplastic pollution of Dongting Lake and its affiliated rivers. Mar Pollut Bull 160:111555. <https://doi.org/10.1016/j.marpolbul.2020.111555>
 63. Lin L, Zuo LZ, Peng JP, Cai LQ, Fok L, Yan Y et al (2018) Occurrence and distribution of microplastics in an urban river: a case study in the Pearl River along Guangzhou City. China Sci Total Environ 644:375–381. <https://doi.org/10.1016/j.scitotenv.2018.06.327>
 64. Wang D, Jiang SY, Fan C, Fu L, Ruan HD (2023) Occurrence and correlation of microplastics and dibutyl phthalate in rivers from Pearl River Delta. China Mar Pollut Bull 197:115759. <https://doi.org/10.1016/j.marpolbul.2023.115759>
 65. Ding J, Sun C, He C, Zheng L, Dai D, Li F (2022) Atmospheric microplastics in the Northwestern Pacific Ocean: distribution, source, and deposition. Sci Total Environ 829:154337. <https://doi.org/10.1016/j.scitotenv.2022.154337>
 66. Baldwin AK, Corsi SR, Mason SA (2016) Plastic debris in 29 great lakes tributaries: relations to watershed attributes and hydrology. Environ Sci Technol 50:10377–10385. <https://doi.org/10.1021/acs.est.6b02917>
 67. McCormick A, Hoellein TJ, Mason SA, Schluep J, Kelly JJ (2014) Microplastic is an abundant and distinct microbial habitat in an urban river. Environ Sci Technol 48:11863–11871. <https://doi.org/10.1021/es503610r>
 68. Yonkos LT, Friedel EA, Perez-Reyes AC, Ghosal S, Arthur CD (2014) Microplastics in four estuarine rivers in the Chesapeake bay, USA. Environ Sci Technol 48:14195–14202. <https://doi.org/10.1021/es5036317>
 69. Di Mauro R, Kupchik MJ, Benfield MC (2017) Abundant plankton-sized microplastic particles in shelf waters of the northern Gulf of Mexico. Environ Pollut 230:798–809. <https://doi.org/10.1016/j.envpol.2017.07.030>
 70. Eriksen M, Mason S, Wilson S, Box C, Zellers A, Edwards W et al (2013) Microplastic pollution in the surface waters of the

- Laurentian Great Lakes. *Mar Pollut Bull* 77:177–182. <https://doi.org/10.1016/j.envpol.2021.116695>
71. Nitzberg EJ, Parmar S, Arbuckle-Keil G, Saba GK, Chant RJ, Fahrenfeld NL (2024) Microplastic concentration, characterization, and size distribution in the Delaware Bay estuary. *Chemosphere* 361:142523. <https://doi.org/10.1016/j.chemosphere.2024.142523>
 72. Khoironi A, Hadiyanto H, Anggoro S, Sudarno S (2020) Evaluation of polypropylene plastic degradation and microplastic identification in sediments at Tambak Lorok coastal area, Semarang, Indonesia *Mar Pollut Bull* 151:110868. <https://doi.org/10.1016/j.marpolbul.2019.110868>
 73. Mangarengi NAP, Zakaria R, Zubair A, Langka P, Riswanto NA (2023) Distribution of microplastic abundance and composition in surface water around anthropogenic areas (case study: Jeneberang River, south Sulawesi, Indonesia). In: IOP conference series earth environment science, 1134. <https://doi.org/10.1088/1755-1315/1134/1/012039>
 74. Free CM, Jensen OP, Mason SA, Eriksen M, Williamson NJ, Boldgiv B (2014) High-levels of microplastic pollution in a large, remote, mountain lake. *Mar Pollut Bull* 85:156–163. <https://doi.org/10.1016/j.marpolbul.2014.06.001>
 75. Napper IE, Baroth A, Barrett AC, Bhola S, Chowdhury GW, Davies BFR et al (2021) The abundance and characteristics of microplastics in surface water in the transboundary Ganges River. *Environ Pollut* 274:116348. <https://doi.org/10.1016/j.envpol.2020.116348>
 76. Li J, Gao F, Zhang D, Cao W, Zhao C (2022) Zonal distribution characteristics of microplastics in the southern Indian ocean and the influence of ocean current. *J Mar Sci Eng* 10:290. <https://doi.org/10.3390/jmse10020290>
 77. Robin RS, Karthik R, Purvaja R, Ganguly D, Anandavelu I, Mugilarasan M et al (2020) Holistic assessment of microplastics in various coastal environmental matrices, southwest coast of India. *Sci Total Environ* 703:134947. <https://doi.org/10.1016/j.scitotenv.2019.134947>
 78. Ephsy D, Raja S (2023) Characterization of microplastics and its pollution load index in freshwater Kumaraswamy Lake of Coimbatore. *India Environ Toxicol Pharmacol* 101:104207. <https://doi.org/10.1016/j.etap.2023.104207>
 79. Rajan K, Khudsar FA, Kumar R (2023) Urbanization and population resources affect microplastic concentration in surface water of the River Ganga. *J Hazard Mater Adv* 11:100342. <https://doi.org/10.1016/j.hazadv.2023.100342>
 80. Fischer EK, Paglialonga L, Czech E, Tamminga M (2016) Microplastic pollution in lakes and lake shoreline sediments – a case study on Lake Bolsena and Lake Chiusi (central Italy). *Environ Pollut* 213:648–657. <https://doi.org/10.1016/j.envpol.2016.03.012>
 81. Dris R, Gasperi J, Rocher V, Saad M, Renault N, Tassin B et al (2015) Microplastic contamination in an urban area: a case study in greater Paris. *Environ Chem* 12:592–599. <https://doi.org/10.1071/EN14167>
 82. Faure F, Demars C, Wieser O, Kunz M, de Alencastro LF, Faure F et al (2015) Plastic pollution in Swiss surface waters: nature and concentrations, interaction with pollutants. *Environ Chem* 12:582–591. <https://doi.org/10.1071/EN14218>
 83. Semmouri I, Vercauteren M, Van Acker E, Pequeur E, Asselman J, Janssen C (2023) Distribution of microplastics in freshwater systems in an urbanized region: a case study in Flanders (Belgium). *Sci Total Environ* 872:162192. <https://doi.org/10.1016/j.scitotenv.2023.162192>
 84. Lechner A, Keckeis H, Lumesberger-Loisl F, Zens B, Krusch R, Tritthart M et al (2014) The Danube so colourful: a potpourri of plastic litter outnumbers fish larvae in Europe's second largest river. *Environ Pollut* 188:177–181. <https://doi.org/10.1016/j.envpol.2014.02.006>
 85. Godoy V, Calero M, González-Olalla JM, Martín-Lara MA, Olea N, Ruiz-Gutierrez A et al (2022) The human connection, first evidence of microplastics in remote high mountain lakes of Sierra Nevada, Spain *Environ Pollut* 311:119922. <https://doi.org/10.1016/j.envpol.2022.119922>
 86. Campillo A, Almeda R, Vianello A, Gómez M, Martínez I, Navarro A et al (2023) Searching for hotspots of neustonic microplastics in the Canary Islands. *Mar Pollut Bull*. <https://doi.org/10.1016/j.marpolbul.2023.115057>
 87. Büngener L, Schäffer SM, Schwarz A, Schwalb A (2024) Microplastics in a small river: occurrence and influencing factors along the river Oker, Northern Germany. *J Contam Hydrol*. <https://doi.org/10.1016/j.jconhyd.2024.104366>
 88. Isobe A, Uchida K, Tokai T, Iwasaki S (2015) East Asian seas: a hot spot of pelagic microplastics. *Mar Pollut Bull* 101:618–623
 89. Rojas-Luna RA, Oquendo-Ruiz L, García-Alzate CA, Arana VA, García-Alzate R, Trilleras J (2023) Identification, abundance, and distribution of microplastics in surface water collected from Luruaco Lake, low basin Magdalena River, Colombia. *Water* (Switzerland). <https://doi.org/10.3390/w15020344>
 90. Nel HA, Froneman PW (2015) A quantitative analysis of microplastic pollution along the south-eastern coastline of South Africa. *Mar Pollut Bull* 101:274–279. <https://doi.org/10.1016/j.marpolbul.2015.09.043>
 91. de Carvalho DG, Baptista Neto JA (2016) Microplastic pollution of the beaches of Guanabara Bay, Southeast Brazil. *Ocean Coast Manag* 128:10–17. <https://doi.org/10.1016/j.ocecoaman.2016.04.009>
 92. Olivatto GP, Martins MCT, Montagner CC, Henry TB, Carreira RS (2019) Microplastic contamination in surface waters in Guanabara Bay, Rio de Janeiro. *Brazil Mar Pollut Bull* 139:157–162. <https://doi.org/10.1016/j.marpolbul.2018.12.042>
 93. da Costa ID, Costa LL, Zalmon IR (2023) Microplastics in water from the confluence of tropical rivers: overall review and a case study in Paraíba do Sul river basin. *Chemosphere* 338:139493. <https://doi.org/10.1016/j.chemosphere.2023.139493>
 94. Uurasjärvi E, Hartikainen S, Setälä O, Lehtiniemi M, Koistinen A (2020) Microplastic concentrations, size distribution, and polymer types in the surface waters of a northern European lake. *Water Environ Res* 92:149–156. <https://doi.org/10.1002/wer.1229>
 95. Russell M, Webster L (2021) Microplastics in sea surface waters around Scotland. *Mar Pollut Bull* 166:112210. <https://doi.org/10.1016/j.marpolbul.2021.112210>
 96. Tamminga M, Hengstmann E, Fischer EK (2018) Microplastic analysis in the South Funen Archipelago, Baltic Sea, implementing manta trawling and bulk sampling. *Mar Pollut Bull* 128:601–608. <https://doi.org/10.1016/j.marpolbul.2018.01.066>
 97. Liu Y, Lorenz C, Vianello A, Syberg K, Nielsen AH, Nielsen TG et al (2023) Exploration of occurrence and sources of microplastics (> 10 µm) in Danish marine waters. *Sci Total Environ* 865:161255. <https://doi.org/10.1016/j.scitotenv.2022.161255>
 98. Nan B, Su L, Kellar C, Craig NJ, Keough MJ, Pettigrove V (2019) Identification of microplastics in surface water and Australian freshwater shrimp *Paratya australiensis* in Victoria. *Australia Environ Pollut* 259:113865. <https://doi.org/10.1016/j.envpol.2019.113865>
 99. Lusher AL, O'Donnell C, Officer R, O'Connor I (2016) Microplastic interactions with North Atlantic mesopelagic fish. *ICES J Mar Sci* 73:1214–1225. <https://doi.org/10.1093/icesjms/fsv241>
 100. Magnusson K, Norén F (2014) Screening of microplastic particles in and down-stream a wastewater treatment plant (report). <https://www.diva-portal.org/smash/get/diva2:1549880/FULLTEXT01.pdf>

101. Karing DJ, Anggiani M, Cao LTT, El-shaammari M (2023) Occurrence of microplastics in Kemena River and Niah River of Sarawak, Malaysia. *Trop Environ Biol Technol* 1:1–13. <https://doi.org/10.53623/tebt.v1i1.220>
102. Park TJ, Lee SH, Lee MS, Lee JK, Lee SH, Zoh KD (2020) Occurrence of microplastics in the Han River and riverine fish in South Korea. *Sci Total Environ* 708:134535. <https://doi.org/10.1016/j.scitotenv.2019.134535>
103. Pham DT, Choi SH, Kwon JH (2024) Year-round spatial and temporal distribution of microplastics in water and sediments of an urban freshwater system (Jungnang Stream, Korea). *Environ Pollut* 357:124362. <https://doi.org/10.1016/j.envpol.2024.124362>
104. Lahens L, Strady E, Kieu-Le TC, Dris R, Boukerma K, Rinnert E et al (2018) Macroplastic and microplastic contamination assessment of a tropical river (Saigon River, Vietnam) transversed by a developing megacity. *Environ Pollut* 236:661–671. <https://doi.org/10.1016/j.envpol.2018.02.005>
105. Vayghan AH, Rasta M, Zakeri M, Kelly FJ (2022) Spatial distribution of microplastics pollution in sediments and surface waters of the Aras River and reservoir: an international river in North-western Iran. *Sci Total Environ* 843:156894. <https://doi.org/10.1016/j.scitotenv.2022.156894>
106. Irfan M, Qadir A, Mumtaz M, Ahmad SR (2020) An unintended challenge of microplastic pollution in the urban surface water system of Lahore. *Pakistan Environ Sci Pollut Res* 27:16718–16730. <https://doi.org/10.1007/s11356-020-08114-7>
107. Higgins C, Turner A (2023) Microplastics in surface coastal waters around Plymouth, UK, and the contribution of boating and shipping activities. *Sci Total Environ* 893:164695. <https://doi.org/10.1016/j.scitotenv.2023.164695>
108. Kanhai LDK, Gårdfeldt K, Lyashevskaya O, Hassellöv M, Thompson RC, O'Connor I (2018) Microplastics in sub-surface waters of the Arctic Central Basin. *Mar Pollut Bull* 130:8–18. <https://doi.org/10.1016/j.marpolbul.2018.03.011>
109. Ikenoue T, Nakajima R, Fujiwara A, Onodera J, Itoh M, Toyoshima J et al (2023) Horizontal distribution of surface microplastic concentrations and water-column microplastic inventories in the Chukchi Sea, western Arctic Ocean. *Sci Total Environ* 855:159564. <https://doi.org/10.1016/j.scitotenv.2022.159564>
110. Mani T, Hauk A, Walter U, Burkhardt-Holm P (2015) Microplastics profile along the Rhine River. *Sci Rep* 5(5):1–7. <https://doi.org/10.1038/srep17988>
111. Güven O, Gökdağ K, Jovanović B, Kideys AE (2017) Microplastic litter composition of the Turkish territorial waters of the Mediterranean Sea, and its occurrence in the gastrointestinal tract of fish. *Environ Pollut* 223:286–294. <https://doi.org/10.1016/j.envpol.2017.01.025>
112. Maes T, Van Der Meulen MD, Devriese LI, Leslie HA, Huvet A, Frère L et al (2017) Microplastics baseline surveys at the water surface and in sediments of the North-East Atlantic. *Front Mar Sci* 4:135. <https://doi.org/10.3389/fmars.2017.00135>
113. Desforges JPW, Galbraith M, Dangerfield N, Ross PS (2014) Widespread distribution of microplastics in subsurface seawater in the NE Pacific Ocean. *Mar Pollut Bull* 79:94–99. <https://doi.org/10.1016/j.marpolbul.2013.12.035>
114. Mataji A, Taleshi MS, Balimoghaddas E (2020) Distribution and characterization of microplastics in surface waters and the Southern Caspian Sea coastal sediments. *Arch Environ Contam Toxicol* 78:86–93. <https://doi.org/10.1007/s00244-019-00700-2>
115. Viršek MK, Lovšin MN, Koren Š, Kržan A, Peterlin M (2017) Microplastics as a vector for the transport of the bacterial fish pathogen species *Aeromonas salmonicida*. *Mar Pollut Bull* 125:301–309. <https://doi.org/10.1016/j.marpolbul.2017.08.024>
116. Goswami P, Selvakumar N, Verma P, Saha M, Suneel V, Vinithkumar NV et al (2023) Microplastic intrusion into the zooplankton, the base of the marine food chain: evidence from the Arabian Sea. *Indian Ocean Sci Total Environ* 864:160876. <https://doi.org/10.1016/j.scitotenv.2022.160876>
117. Yuan D, Corvianawatie C, Cordova MR, Surinati D, Li Y, Wang Z et al (2023) Microplastics in the tropical Northwestern Pacific Ocean and the Indonesian seas. *J Sea Res* 194:102406. <https://doi.org/10.1016/j.seares.2023.102406>
118. Lambert S, Wagner M (2018) Microplastics are contaminants of emerging concern in freshwater environments: an overview - freshwater microplastics: emerging environmental contaminants? Springer International Publishing, Cham, pp 1–23. https://doi.org/10.1007/978-3-319-61615-5_1
119. WHO (2019) Microplastics in drinking water
120. Liu W, Zhang J, Liu H, Guo X, Zhang X, Yao X et al (2021) A review of the removal of microplastics in global wastewater treatment plants: characteristics and mechanisms. *Environ Int* 146:106277. <https://doi.org/10.1016/j.envint.2020.106277>
121. Hettiarachchi H, Meegoda JN (2023) Microplastic pollution prevention: the need for robust policy interventions to close the loopholes in current waste management practices. *Int J Environ Res Public Health* 20:6434. <https://doi.org/10.3390/ijerph20146434>
122. Anagnosti L, Varvaresou A, Pavlou P, Protopapa E, Carayanni V (2021) Worldwide actions against plastic pollution from microbeads and microplastics in cosmetics focusing on European policies. Has the issue been handled effectively? *Mar Pollut Bull* 162:111883. <https://doi.org/10.1016/j.marpolbul.2020.111883>
123. Khan NA, Khan AH, López-Maldonado EA, Alam SS, López López JR, Méndez Herrera PF et al (2022) Microplastics: occurrences, treatment methods, regulations and foreseen environmental impacts. *Environ Res* 215:114224. <https://doi.org/10.1016/j.envres.2022.114224>
124. Knoblauch D, Mederake L (2021) Government policies combating plastic pollution. *Curr Opin Toxicol* 28:87–96. <https://doi.org/10.1016/j.cotox.2021.10.003>
125. Masiá P, Sol D, Ardura A, Laca A, Borrell YJ, Dopico E et al (2020) Bioremediation as a promising strategy for microplastics removal in wastewater treatment plants. *Mar Pollut Bull*. <https://doi.org/10.1016/j.marpolbul.2020.111252>
126. ECHA (2019) Annex XV Restriction Report - Microplastics. version 12, 22 August
127. European Commission (2022) Directive of the European Parliament and of the council concerning urban wastewater treatment (recast) (Text with EEA relevance) 345
128. Boucher J, Friot D (2017) International union for conservation of nature a global evaluation of sources primary microplastics in the oceans
129. Laura Parker (2020) Microplastics have moved into virtually every crevice on Earth. *Natl. Geogr. Mag*. <https://www.nationalgeographic.com/science/article/microplastics-in-virtually-every-crevice-on-earth>. Accessed 23 Feb 2023
130. da Costa JP, Mouneyrac C, Costa M, Duarte AC, Rocha-Santos T (2020) The role of legislation, regulatory initiatives and guidelines on the control of plastic pollution. *Front Environ Sci* 8:104. <https://doi.org/10.3389/fenvs.2020.00104>
131. Li Y (2022) legislation and policy on pollution prevention and the control of marine microplastics. *Water* (Switzerland). <https://doi.org/10.3390/w14182790>
132. Agência Portuguesa do Ambiente (2022) Alterada e Republicada pelo Decreto-Lei n. 78 / 2021 , de 24 de setembro De acordo com a alínea e) do artigo 2. da Lei n. 76 / 2019 , de 2 de setembro , conforme alterada pelo Decreto-Lei n. 78 / 2021 , de 24 de setembro , entende-se por “ L, 1–9
133. Ellen Macarthur Foundation (2001) Government insights - United Nations report. *Chem Eng News*, 79 35

134. Wang Q, Tweedy A, Wang HG (2022) Reducing plastic waste through legislative interventions in the United States: development, obstacles, potentials, and challenges. *Sustain Horizons* 2:100013. <https://doi.org/10.1016/j.horiz.2022.100013>
135. Sorensen RM, Kanwar RS, Jovanovi B (2023) Past, present, and possible future policies on plastic use in the US, particularly microplastics and nanoplastics: a review. *Integr Environ Assess Manag* 19:474–488. <https://doi.org/10.1002/ieam.4678>
136. Murphy F, Ewins C, Carbonnier F, Quinn B (2016) Wastewater treatment works (WwTW) as a source of microplastics in the aquatic environment. *Environ Sci Technol* 50:5800–5808. <https://doi.org/10.1021/acs.est.5b05416>
137. Sun J, Dai X, Wang Q, van Loosdrecht MCM, Ni B-J (2019) Microplastics in wastewater treatment plants: detection, occurrence and removal. *Water Res* 152:21–37. <https://doi.org/10.1016/j.watres.2018.12.050>
138. Bakaraki Turan N, Sari Erkan H, Onkal Engin G (2021) Microplastics in wastewater treatment plants: occurrence, fate and identification. *Process Saf Environ Prot* 146:77–84. <https://doi.org/10.1016/j.psep.2020.08.039>
139. Fältström E, Olesen KB, Anderberg S (2021) Microplastic types in the wastewater system—a comparison of material flow-based source estimates and the measurement-based load to a wastewater treatment plant. *Sustain* 13:5404. <https://doi.org/10.3390/su13105404>
140. Yaseen A, Assad I, Sofi MS, Hashmi MZ, Bhat SU (2022) A global review of microplastics in wastewater treatment plants: understanding their occurrence, fate and impact. *Environ Res* 212:113258. <https://doi.org/10.1016/j.envres.2022.113258>
141. Elkhatab D, Oyanedel-Craver V (2020) A critical review of extraction and identification methods of microplastics in wastewater and drinking water. *Environ Sci Technol* 54:7037–7049. <https://doi.org/10.1021/acs.est.9b06672>
142. Verla AW, Enyoh CE, Verla EN, Nwornorh KO (2019) Microplastic–toxic chemical interaction: a review study on quantified levels, mechanism and implication. *SN Appl Sci* 1:1–30. <https://doi.org/10.1007/s42452-019-1352-0>
143. Yuan Z, Nag R, Cummins E (2022) Human health concerns regarding microplastics in the aquatic environment - from marine to food systems. *Sci Total Environ* 823:153730. <https://doi.org/10.1016/j.scitotenv.2022.153730>
144. Alimi OS, Farner Budarz J, Hernandez LM, Tufenkji N (2018) Microplastics and nanoplastics in aquatic environments: aggregation, deposition, and enhanced contaminant transport. *Environ Sci Technol* 52:1704–1724. <https://doi.org/10.1021/acs.est.7b05559>
145. Cristaldi A, Fiore M, Zuccarello P, Conti GO, Grasso A, Nicolosi I et al (2020) Efficiency of wastewater treatment plants (Wwtps) for microplastic removal: a systematic review. *Int J Environ Res Public Health* 17:1–23. <https://doi.org/10.3390/ijerph17218014>
146. Liu S, Shi J, Wang J, Dai Y, Li H, Li J et al (2021) Interactions between microplastics and heavy metals in aquatic environments: a review. *Front Microbiol* 12:1–14. <https://doi.org/10.3389/fmicb.2021.652520>
147. Sun T, Wang S, Ji C, Li F, Wu H (2022) Microplastics aggravate the bioaccumulation and toxicity of coexisting contaminants in aquatic organisms: a synergistic health hazard. *J Hazard Mater* 424:127533. <https://doi.org/10.1016/j.jhazmat.2021.127533>
148. Zhang Z, Chen Y (2020) Effects of microplastics on wastewater and sewage sludge treatment and their removal: a review. *Chem Eng J* 382:122955. <https://doi.org/10.1016/j.cej.2019.122955>
149. Shan J, Wang J, Zhan J, Liu L, Wu F, Wang X (2020) Sorption behaviors of crude oil on polyethylene microplastics in seawater and digestive tract under simulated real-world conditions. *Chemosphere* 257:127225. <https://doi.org/10.1016/j.chemosphere.2020.127225>
150. Yang M, Zhang B, Xin X, Liu B, Zhu Z, Dong G et al (2022) Microplastic-oil-dispersant agglomerates in the marine environment: formation mechanism and impact on oil dispersion. *J Hazard Mater* 426:127825. <https://doi.org/10.1016/j.jhazmat.2021.127825>
151. Yang M, Zhang B, Xin X, Lee K, Chen B (2022) Microplastic and oil pollution in oceans: interactions and environmental impacts. *Sci Total Environ* 838:156142. <https://doi.org/10.1016/j.scitotenv.2022.156142>
152. Cao Y, Zhao M, Ma X, Song Y, Zuo S, Li H et al (2021) A critical review on the interactions of microplastics with heavy metals: mechanism and their combined effect on organisms and humans. *Sci Total Environ* 788:147620. <https://doi.org/10.1016/j.scitotenv.2021.147620>
153. Khalid N, Aqeel M, Noman A, Khan SM, Akhter N (2021) Interactions and effects of microplastics with heavy metals in aquatic and terrestrial environments. *Environ Pollut* 290:118104. <https://doi.org/10.1016/j.envpol.2021.118104>
154. Kirstein IV, Gomiero A, Vollertsen J (2021) Microplastic pollution in drinking water. *Curr Opin Toxicol* 28:70–75. <https://doi.org/10.1016/j.cotox.2021.09.003>
155. Zhang Q, Xu EG, Li J, Chen Q, Ma L, Zeng EY et al (2020) A review of microplastics in table salt, drinking water, and air: direct human exposure. *Environ Sci Technol* 54:3740–3751. <https://doi.org/10.1021/acs.est.9b04535>
156. Bigalke M, Fieber M, Foetisch A, Reynes J, Tollan P (2022) Microplastics in agricultural drainage water: a link between terrestrial and aquatic microplastic pollution. *Sci Total Environ* 806:150709. <https://doi.org/10.1016/j.scitotenv.2021.150709>
157. Lwanga EH, Beriot N, Corradini F, Silva V, Yang X, Baartman J et al (2022) Review of microplastic sources, transport pathways and correlations with other soil stressors: a journey from agricultural sites into the environment. *Chem Biol Technol Agric* 9:20. <https://doi.org/10.1186/s40538-021-00278-9>
158. Chia RW, Lee J-Y, Kim H, Jang J (2021) Microplastic pollution in soil and groundwater: a review. *Environ Chem Lett* 19:4211–4224. <https://doi.org/10.1007/s10311-021-01297-6>
159. Viaroli S, Lancia M, Re V (2022) Microplastics contamination of groundwater: current evidence and future perspectives, a review. *Sci Total Environ* 824:153851. <https://doi.org/10.1016/j.scitotenv.2022.153851>
160. Laura Parker (2022) Microplastics are in our bodies. How much do they harm us? *Environment*, 1–2. <https://www.nationalgeographic.com/environment/article/microplastics-are-in-our-bodies-how-much-do-they-harm-us>
161. Ragusa A, Svelato A, Santacroce C, Catalano P, Notarstefano V, Carnevali O et al (2021) Plasticenta: first evidence of microplastics in human placenta. *Environ Int* 146:106274. <https://doi.org/10.1016/j.envint.2020.106274>
162. Eerkes-Medrano D, Leslie HA, Quinn B (2019) Microplastics in drinking water: a review and assessment. *Curr Opin Environ Sci Heal* 7:69–75. <https://doi.org/10.1016/j.coesh.2018.12.001>
163. Gambino I, Bagordo F, Grassi T, Panico A, De Donno A (2022) Occurrence of microplastics in tap and bottled water: current knowledge. *Int J Environ Res Public Health* 19:5283. <https://doi.org/10.3390/ijerph19095283>
164. Mason SA, Welch VG, Neratko J (2018) Synthetic polymer contamination in bottled water. *Front Chem* 6:407. <https://doi.org/10.3389/fchem.2018.00407>
165. Krishnan RY, Manikandan S, Subbaiya R, Karmegam N, Kim W, Govarthan M (2023) Recent approaches and advanced wastewater treatment technologies for mitigating emerging microplastics contamination – a critical review. *Sci Total Environ* 858:159258. <https://doi.org/10.1016/j.scitotenv.2023.159258>

- Environ 858:159681. <https://doi.org/10.1016/j.scitotenv.2022.159681>
166. Bayo J, Olmos S, López-Castellanos J (2020) Microplastics in an urban wastewater treatment plant: the influence of physico-chemical parameters and environmental factors. *Chemosphere* 238:124593. <https://doi.org/10.1016/j.chemosphere.2019.124593>
 167. Lares M, Ncibi MC, Sillanpää M, Sillanpää M (2018) Occurrence, identification and removal of microplastic particles and fibers in conventional activated sludge process and advanced MBR technology. *Water Res* 133:236–246. <https://doi.org/10.1016/j.watres.2018.01.049>
 168. Wolff S, Weber F, Kerpen J, Winklhofer M, EngelhartBarkmann L (2021) Elimination of microplastics by downstream sand filters in wastewater treatment. *Water* 13:33. <https://doi.org/10.3390/w13010033>
 169. Talvitie J, Mikola A, Koistinen A, Setälä O (2017) Solutions to microplastic pollution – removal of microplastics from wastewater effluent with advanced wastewater treatment technologies. *Water Res* 123:401–407. <https://doi.org/10.1016/j.watres.2017.07.005>
 170. Hidayaturrehman H, Lee T-G (2019) A study on characteristics of microplastic in wastewater of South Korea: identification, quantification, and fate of microplastics during treatment process. *Mar Pollut Bull* 146:696–702. <https://doi.org/10.1016/j.marpolbul.2019.06.071>
 171. Rajala K, Grönfors O, Hesampour M, Mikola A (2020) Removal of microplastics from secondary wastewater treatment plant effluent by coagulation/flocculation with iron, aluminum and polyamine-based chemicals. *Water Res* 183:116045. <https://doi.org/10.1016/j.watres.2020.116045>
 172. Perren W, Wojtasik A, Cai Q (2018) Removal of microbeads from wastewater using electrocoagulation. *ACS Omega* 3:3357–3364. <https://doi.org/10.1021/acsomega.7b02037>
 173. Shen M, Zhang Y, Almatrafi E, Hu T, Zhou C, Song B et al (2022) Efficient removal of microplastics from wastewater by an electrocoagulation process. *Chem Eng J* 428:131161. <https://doi.org/10.1016/j.cej.2021.131161>
 174. Feilin H, Mingwei S (2022) Ecofriendly removing microplastics from rivers: a novel air flotation approach crafted with positively charged carrier. *Process Saf Environ Prot* 168:613–623. <https://doi.org/10.1016/j.psep.2022.09.060>
 175. Wang Z, Lin T, Chen W (2020) Occurrence and removal of microplastics in an advanced drinking water treatment plant (ADWTP). *Sci Total Environ* 700:134520. <https://doi.org/10.1016/j.scitotenv.2019.134520>
 176. Sol D, Laca A, Laca A, Díaz M (2021) Microplastics in wastewater and drinking water treatment plants: occurrence and removal of microfibrils. *Appl Sci* 11:10109. <https://doi.org/10.3390/app112110109>
 177. Tadsuwan K, Babel S (2022) Microplastic abundance and removal via an ultrafiltration system coupled to a conventional municipal wastewater treatment plant in Thailand. *J Environ Chem Eng* 10:107142. <https://doi.org/10.1016/j.jece.2022.107142>
 178. Edo C, González-Pleiter M, Leganés F, Fernández-Piñas F, Rosal R (2020) Fate of microplastics in wastewater treatment plants and their environmental dispersion with effluent and sludge. *Environ Pollut* 259:113837. <https://doi.org/10.1016/j.envpol.2019.113837>
 179. Magni S, Binelli A, Pittura L, Avio CG, Della Torre C, Parenti CC et al (2019) The fate of microplastics in an Italian wastewater treatment plant. *Sci Total Environ* 652:602–610. <https://doi.org/10.1016/j.scitotenv.2018.10.269>
 180. Leslie HA, Brandsma SH, van Velzen MJM, Vethaak AD (2017) Microplastics en route: field measurements in the Dutch river delta and Amsterdam canals, wastewater treatment plants, North Sea sediments and biota. *Environ Int* 101:133–142. <https://doi.org/10.1016/j.envint.2017.01.018>
 181. Lv X, Dong Q, Zuo Z, Liu Y, Huang X, Wu WM (2019) Microplastics in a municipal wastewater treatment plant: fate, dynamic distribution, removal efficiencies, and control strategies. *J Clean Prod* 225:579–586. <https://doi.org/10.1016/j.jclepro.2019.03.321>
 182. Michielssen MR, Michielssen ER, Ni J, Duhaime MB (2016) Fate of microplastics and other small anthropogenic litter (SAL) in wastewater treatment plants depends on unit processes employed. *Environ Sci Water Res Technol* 2:1064–1073. <https://doi.org/10.1039/C6EW00207B>
 183. Pittura L, Foglia A, Akyol Ç, Cipolletta G, Benedetti M, Regoli F et al (2021) Microplastics in real wastewater treatment schemes: comparative assessment and relevant inhibition effects on anaerobic processes. *Chemosphere* 262:128415. <https://doi.org/10.1016/j.chemosphere.2020.128415>
 184. Dalmau-Soler J, Ballesteros-Cano R, Boleda MR, Paraira M, Ferrer N, Lacorte S (2021) Microplastics from headwaters to tap water: occurrence and removal in a drinking water treatment plant in Barcelona Metropolitan area (Catalonia, NE Spain). *Environ Sci Pollut Res* 28:59462–59472. <https://doi.org/10.1007/s11356-021-13220-1>
 185. Ziajahromi S, Neale PA, Telles Silveira I, Chua A, Leusch FDL (2021) An audit of microplastic abundance throughout three Australian wastewater treatment plants. *Chemosphere* 263:128294. <https://doi.org/10.1016/j.chemosphere.2020.128294>
 186. Park H-J, Oh M-J, Kim P-G, Kim G, Jeong D-H, Ju B-K et al (2020) National reconnaissance survey of microplastics in municipal wastewater treatment plants in Korea. *Environ Sci Technol* 54:1503–1512. <https://doi.org/10.1021/acs.est.9b04929>
 187. Yang L, Li K, Cui S, Kang Y, An L, Lei K (2019) Removal of microplastics in municipal sewage from China's largest water reclamation plant. *Water Res* 155:175–181. <https://doi.org/10.1016/j.watres.2019.02.046>
 188. Yuan F, Zhao H, Sun H, Zhao J, Sun Y (2021) Abundance, morphology, and removal efficiency of microplastics in two wastewater treatment plants in Nanjing. *China Environ Sci Pollut Res* 28:9327–9337. <https://doi.org/10.1007/s11356-020-11411-w>
 189. Van DM, Le TXT, Vu ND, Dang TT (2022) Distribution and occurrence of microplastics in wastewater treatment plants. *Environ Technol Innov* 26:102286. <https://doi.org/10.1016/j.eti.2022.102286>
 190. Vuori L, Ollikainen M (2022) How to remove microplastics in wastewater? A cost-effectiveness analysis. *Ecol Econ* 192:107246. <https://doi.org/10.1016/j.ecolecon.2021.107246>
 191. van Haandel AC, van der Lubbe JGM (2012) Handbook of biological wastewater treatment: design and optimisation of activated sludge systems. IWA. <https://doi.org/10.2166/9781780400808>
 192. Krzeminski P, Leverette L, Malamis S, Katsou E (2017) Membrane bioreactors – a review on recent developments in energy reduction, fouling control, novel configurations, LCA and market prospects. *J Memb Sci* 527:207–227. <https://doi.org/10.1016/j.memsci.2016.12.010>
 193. Lesjean B, Tazi-Pain A, Thaure D, Moeslang H, Buisson HT (2011) persistent myths and the realities of membrane bioreactor technology for municipal applications. *Water Sci Technol* 63:32–39. <https://doi.org/10.2166/wst.2011.005>
 194. Naushad M, Sharma G, Alothman ZA (2019) Photodegradation of toxic dye using gum Arabic-crosslinked-poly(acrylamide)/Ni(OH)₂/FeOOH nanocomposites hydrogel. *J Clean Prod* 241:118263. <https://doi.org/10.1016/j.jclepro.2019.118263>
 195. Wu S, Shi W, Li K, Cai J, Chen L (2022) Recent advances on sustainable bio-based materials for water treatment: fabrication, modification and application. *J Environ Chem Eng* 10:108921. <https://doi.org/10.1016/j.jece.2022.108921>

196. Nyika J, Dinka MO (2022) A mini-review on wastewater treatment through bioremediation towards enhanced field applications of the technology. *AIMS Environ Sci* 9:403–431. <https://doi.org/10.3934/environsci.2022025>
197. Zhou Y, Kumar M, Sarsaiya S, Sirohi R, Awasthi SK, Sindhu R et al (2022) Challenges and opportunities in bioremediation of micro-nano plastics: a review. *Sci Total Environ* 802:149823. <https://doi.org/10.1016/j.scitotenv.2021.149823>
198. Aliko V, Multisanti CR, Turani B, Faggio C (2022) Get rid of marine pollution: bioremediation an innovative, attractive, and successful cleaning strategy. *Sustain* 14:11784. <https://doi.org/10.3390/su141811784>
199. Miloloža M, Cvetnić M, Kučić Grgić D, Očelić Bulatović V, Ukić Š, Rogošić M et al (2022) Biotreatment strategies for the removal of microplastics from freshwater systems, a review. *Environ Chem Lett* 20:1377–1402. <https://doi.org/10.1007/s10311-021-01370-0>
200. Lin Z, Jin T, Zou T, Xu L, Xi B, Xu D et al (2022) Current progress on plastic/microplastic degradation: fact influences and mechanism. *Environ Pollut* 304:119159. <https://doi.org/10.1016/j.envpol.2022.119159>
201. Manzi HP, Abou-Shanab RAI, Jeon BH, Wang J, Salama ES (2022) Algae: a frontline photosynthetic organism in the microplastic catastrophe. *Trends Plant Sci* 27:1159–1172. <https://doi.org/10.1016/j.tplants.2022.06.005>
202. Tang KHD, Hadibarata T (2022) The application of bioremediation in wastewater treatment plants for microplastics removal: a practical perspective. *Bioprocess Biosyst Eng* 45:1865–1878. <https://doi.org/10.1007/s00449-022-02793-x>
203. Hadian-ghazvini S, Saboor FH, Ardekani LS (2022) Microplastics pollution in aquatic media. Springer, Singapore. <https://doi.org/10.1007/978-981-16-8440-1>
204. Mintenig SM, Int-Veen I, Löder MGJ, Primpke S, Gerdt S G (2017) Identification of microplastic in effluents of waste water treatment plants using focal plane array-based micro-Fourier-transform infrared imaging. *Water Res* 108:365–372. <https://doi.org/10.1016/j.watres.2016.11.015>
205. Isobe A, Uchiyama-Matsumoto K, Uchida K, Tokai T (2017) Microplastics in the Southern Ocean. *Mar Pollut Bull* 114:623–626. <https://doi.org/10.1016/j.marpolbul.2016.09.037>
206. Tadsuwan K, Babel S (2022) Unraveling microplastics removal in wastewater treatment plant: a comparative study of two wastewater treatment plants in Thailand. *Chemosphere* 307:135733. <https://doi.org/10.1016/j.chemosphere.2022.135733>
207. Kazour M, Terki S, Rabhi K, Jemaa S, Khalaf G, Amara R (2019) Sources of microplastics pollution in the marine environment: importance of wastewater treatment plant and coastal landfill. *Mar Pollut Bull* 146:608–618. <https://doi.org/10.1016/j.marpolbul.2019.06.066>
208. Yang J, Monnot M, Sun Y, Asia L, Wong-Wah-Chung P, Doumenq P et al (2023) Microplastics in different water samples (seawater, freshwater, and wastewater): removal efficiency of membrane treatment processes. *Water Res* 232:119673. <https://doi.org/10.1016/j.watres.2023.119673>
209. Carr SA, Liu J, Tesoro AG (2016) Transport and fate of microplastic particles in wastewater treatment plants. *Water Res* 91:174–182. <https://doi.org/10.1016/j.watres.2016.01.002>
210. Conley K, Clum A, Deepe J, Lane H, Beckingham B (2019) Wastewater treatment plants as a source of microplastics to an urban estuary: removal efficiencies and loading per capita over one year. *Water Res X* 3:100030. <https://doi.org/10.1016/j.wroa.2019.100030>
211. Mason SA, Garneau D, Sutton R, Chu Y, Ehmann K, Barnes J et al (2016) Microplastic pollution is widely detected in US municipal wastewater treatment plant effluent. *Environ Pollut* 218:1045–1054. <https://doi.org/10.1016/j.envpol.2016.08.056>
212. Ma B, Xue W, Hu C, Liu H, Qu J, Li L (2019) Characteristics of microplastic removal via coagulation and ultrafiltration during drinking water treatment. *Chem Eng J* 359:159–167. <https://doi.org/10.1016/j.cej.2018.11.155>
213. Liu X, Yuan W, Di M, Li Z, Wang J (2019) Transfer and fate of microplastics during the conventional activated sludge process in one wastewater treatment plant of China. *Chem Eng J* 362:176–182. <https://doi.org/10.1016/j.cej.2019.01.033>
214. Zhang B, Wu Q, Gao S, Ruan Y, Qi G, Guo K et al (2023) Distribution and removal mechanism of microplastics in urban wastewater plants systems via different processes. *Environ Pollut* 320:121076. <https://doi.org/10.1016/j.envpol.2023.121076>
215. Li L, Xu G, Yu H, Xing J (2018) Dynamic membrane for micro-particle removal in wastewater treatment: performance and influencing factors. *Sci Total Environ* 627:332–340. <https://doi.org/10.1016/j.scitotenv.2018.01.239>
216. Franco AA, Arellano JM, Albendín G, Rodríguez-Barroso R, Quiroga JM, Coello MD (2021) Microplastic pollution in wastewater treatment plants in the city of Cádiz: abundance, removal efficiency and presence in receiving water body. *Sci Total Environ* 776:145795. <https://doi.org/10.1016/j.scitotenv.2021.145795>
217. Dronjak L, Exposito N, Sierra J, Schuhmacher M, Florencio K, Corzo B et al (2023) Tracing the fate of microplastic in wastewater treatment plant: a multi-stage analysis of treatment units and sludge. *Environ Pollut* 333:122072. <https://doi.org/10.1016/j.envpol.2023.122072>
218. Ma M, Huo M, Coulon F, Ali M, Tang Z, Liu X et al (2024) Understanding microplastic presence in different wastewater treatment processes: removal efficiency and source identification. *Sci Total Environ* 929:172680. <https://doi.org/10.1016/j.scitotenv.2024.172680>
219. Tagg AS, Sapp M, Harrison JP, Sinclair CJ, Bradley E, Ju-Nam Y et al (2020) Microplastic monitoring at different stages in a wastewater treatment plant using reflectance micro-FTIR imaging. *Front Environ Sci* 8:1–9. <https://doi.org/10.3389/fenvs.2020.00145>
220. Johnson AC, Ball H, Cross R, Horton AA, Jürgens MD, Read DS et al (2020) Identification and quantification of microplastics in potable water and their sources within water treatment works in England and Wales. *Environ Sci Technol* 54:12326–12334. <https://doi.org/10.1021/acs.est.0c03211>
221. Liu F, Nord NB, Bester K, Vollertsen J (2020) Microplastics removal from treated wastewater by a biofilter. *Water (Switzerland)* 12:1–11. <https://doi.org/10.3390/w12041085>
222. Simon M, van Alst N, Vollertsen J (2018) Quantification of microplastic mass and removal rates at wastewater treatment plants applying Focal Plane Array (FPA)-based Fourier Transform Infrared (FT-IR) imaging. *Water Res* 142:1–9. <https://doi.org/10.1016/j.watres.2018.05.019>
223. Kirstein IV, Hensel F, Gomiero A, Iordachescu L, Vianello A, Wittgren HB et al (2021) Drinking plastics? Quantification and qualification of microplastics in drinking water distribution systems by μ FTIR and Py-GC-MS. *Water Res* 188:116519. <https://doi.org/10.1016/j.watres.2020.116519>
224. Ruan Y, Zhang K, Wu C, Wu R, Lam PKS (2019) A preliminary screening of HBCD enantiomers transported by microplastics in wastewater treatment plants. *Sci Total Environ* 674:171–178. <https://doi.org/10.1016/j.scitotenv.2019.04.007>
225. Lam TWL, Ho HT, Ma ATH, Fok L (2020) Microplastic contamination of surfacewater-sourced tap water in hong kong. A preliminary study *Appl Sci* 10:3463. <https://doi.org/10.3390/app10103463>
226. Ben-David EA, Habibi M, Haddad E, Hasanin M, Angel DL, Booth AM et al (2021) Microplastic distributions in a domestic

- wastewater treatment plant: removal efficiency, seasonal variation and influence of sampling technique. *Sci Total Environ* 752:141880. <https://doi.org/10.1016/j.scitotenv.2020.141880>
227. Kwon HJ, Hidayaturrehman H, Peera SG, Lee TG (2022) Elimination of microplastics at different stages in wastewater treatment plants. *Water (Switzerland)* 14:2404. <https://doi.org/10.3390/w14152404>
 228. Gies EA, LeNoble JL, Noël M, Etemadifar A, Bishay F, Hall ER et al (2018) Retention of microplastics in a major secondary wastewater treatment plant in Vancouver. *Canada Mar Pollut Bull* 133:553–561. <https://doi.org/10.1016/j.marpolbul.2018.06.006>
 229. Grbic J, Nguyen B, Guo E, You JB, Sinton D, Rochman CM (2019) Magnetic extraction of microplastics from environmental samples. *Environ Sci Technol Lett* 6:68–72. <https://doi.org/10.1021/acs.estlett.8b00671>
 230. Pivokonsky M, Cermakova L, Novotna K, Peer P, Cajthaml T, Janda V (2018) Occurrence of microplastics in raw and treated drinking water. *Sci Total Environ* 643:1644–1651. <https://doi.org/10.1016/j.scitotenv.2018.08.102>
 231. Takdastan A, Niari MH, Babaei A, Dobaradaran S, Jorfi S, Ahmadi M (2021) Occurrence and distribution of microplastic particles and the concentration of di 2-ethyl hexyl phthalate (DEHP) in microplastics and wastewater in the wastewater treatment plant. *J Environ Manage* 280:111851. <https://doi.org/10.1016/j.jenvman.2020.111851>
 232. Alavian Petroody SS, Hashemi SH, van Gestel CAM (2021) Transport and accumulation of microplastics through wastewater treatment sludge processes. *Chemosphere* 278:130471. <https://doi.org/10.1016/j.chemosphere.2021.130471>
 233. Feizi F, Akhbarizadeh R, Hamidian AH (2024) Microplastics in urban water systems, Tehran metropolitan, Iran. *Environ Monit Assess* 196:643. <https://doi.org/10.1007/s10661-024-12815-8>
 234. Ormaniec P (2024) Occurrence and analysis of microplastics in municipal wastewater. *Poland Environ Sci Pollut Res* 31:49646–49655. <https://doi.org/10.1007/s11356-024-34488-z>
 235. Kalčíková G, Alič B, Skalar T, Bundschuh M, Gotvajn AŽ (2017) Wastewater treatment plant effluents as source of cosmetic polyethylene microbeads to freshwater. *Chemosphere* 188:25–31. <https://doi.org/10.1016/j.chemosphere.2017.08.131>
 236. Gündoğdu S, Çevik C, Güzel E, Kilercioğlu S (2018) Microplastics in municipal wastewater treatment plants in Turkey: a comparison of the influent and secondary effluent concentrations. *Environ Monit Assess* 190:626. <https://doi.org/10.1007/s10661-018-7010-y>
 237. Koyuncuoğlu P, Erden G (2023) Microplastics in municipal wastewater treatment plants: a case study of Denizli/Turkey. *Front Environ Sci Eng* 17:99. <https://doi.org/10.1007/s11783-023-1699-8>
 238. Kılıç E, Yücel N, Şahutoğlu SM (2023) Microplastic composition, load and removal efficiency from wastewater treatment plants discharging into Orontes River. *Int J Environ Res* 17:1–11. <https://doi.org/10.1007/s41742-023-00514-0>
 239. Le TMT, Truong TNS, Nguyen PD, Le QDT, Tran QV, Le TT et al (2023) Evaluation of microplastic removal efficiency of wastewater-treatment plants in a developing country. *Vietnam Environ Technol Innov* 29:102994. <https://doi.org/10.1016/j.eti.2022.102994>
 240. Setiadewi N, Henny C, Rohaningsih D, Waluyo A, Soewondo P (2023) Microplastic occurrence and characteristics in a municipal wastewater treatment plant in Jakarta. In: IOP conference series earth environment science, 1201, pp 0–9. <https://doi.org/10.1088/1755-1315/1201/1/012053>
 241. Islam MS, Islam Z, Ahmsim J, Momtaz N, Beauty SA (2023) Removal efficiencies of microplastics of the three largest drinking water treatment plants in Bangladesh. *Sci Total Environ* 895:165155. <https://doi.org/10.1016/j.scitotenv.2023.165155>
 242. Pleskytė S, Uogintė I, Burbulytė A, Byčėnienė S (2023) Characteristics and removal efficiency of microplastics at secondary wastewater treatment plant in Lithuania. *Water Environ Res* 95:e10958. <https://doi.org/10.1002/wer.10958>
 243. Rokomatu MN, Bhagwat-Russell G, Panneerselvam L, Raju S, Savou V, Waqainabete T et al (2023) Influence of different wastewater treatment processes on the rate and characteristics of MPs released from WWTPs in Fiji. *South Pacific Micropl and Nanopl* 3:23. <https://doi.org/10.1186/s43591-023-00068-0>
 244. Hajji S, Ben-Haddad M, Abelouah MR, Rangel-Buitrago N, Ait Alla A (2024) Microplastic characterization and assessment of removal efficiency in an urban and industrial wastewater treatment plant with submarine emission discharge. *Sci Total Environ* 945:174115. <https://doi.org/10.1016/j.scitotenv.2024.174115>
 245. Zhaxylykova D, Alibekov A, Lee W (2024) Seasonal variation and removal of microplastics in a central Asian urban wastewater treatment plant. *Mar Pollut Bull* 205:116597. <https://doi.org/10.1016/j.marpolbul.2024.116597>
 246. Al-Amri A, Yavari Z, Reza Nikoo M, Karimi M (2024) Microplastics removal efficiency and risk analysis of wastewater treatment plants in Oman. *Chemosphere* 359:142206. <https://doi.org/10.1016/j.chemosphere.2024.142206>
 247. Bastakoti S, Adhikari A, Thaiba BM, Neupane BB, Gautam BR, Dangi MB et al (2024) Characterization and removal of microplastics in the Guheshwori wastewater treatment plant. *Nepal Sci Total Environ* 935:173324. <https://doi.org/10.1016/j.scitoenv.2024.173324>
 248. Uddin S, Behbehani M, Habibi N, Faizuddin M, Al-Murad M, Martinez-Guijarro K et al (2022) Microplastics in Kuwait's wastewater streams. *Sustain*. <https://doi.org/10.3390/su142315817>
 249. Sadia M, Mahmood A, Ibrahim M, Irshad MK, Quddusi AHA, Bokhari A et al (2022) Microplastics pollution from wastewater treatment plants: a critical review on challenges, detection, sustainable removal techniques and circular economy. *Environ Technol Innov* 28:102946. <https://doi.org/10.1016/j.eti.2022.102946>
 250. Acarer S (2023) Microplastics in wastewater treatment plants: sources, properties, removal efficiency, removal mechanisms, and interactions with pollutants. *Water Sci Technol* 87:685–710. <https://doi.org/10.2166/wst.2023.022>
 251. Alazaiza MYD, Albahnasawi A, Ahmad Z, Bashir MJK, Al-Wahaibi T, Abujazar MSS et al (2022) Potential use of algae for the bioremediation of different types of wastewater and contaminants: production of bioproducts and biofuel for green circular economy. *J Environ Manage* 324:116415. <https://doi.org/10.1016/j.jenvman.2022.116415>
 252. Agüera A, Plaza-Bolaños P, Fernández FGA (2020) Removal of contaminants of emerging concern by microalgae-based wastewater treatments and related analytical techniques. *Curr Dev Biotechnol Bioeng*. <https://doi.org/10.1016/B978-0-12-819594-9.00020-6>
 253. Pacheco D, Rocha AC, Pereira L, Verdelhos T (2020) Microalgae water bioremediation: trends and hot topics. *Appl Sci* 10:1886. <https://doi.org/10.3390/app10051886>
 254. Sousa CA, Sousa H, Vale F, Simões M (2021) Microalgae-based bioremediation of wastewaters - influencing parameters and mathematical growth modelling. *Chem Eng J*. <https://doi.org/10.1016/j.cej.2021.131412>
 255. Miri S, Saini R, Davoodi SM, Pulicharla R, Brar SK, Magdoui S (2022) Biodegradation of microplastics: better late than never. *Chemosphere* 286:131670. <https://doi.org/10.1016/j.chemosphere.2021.131670>
 256. Maizatul AY, Radin Mohamed RMS, Al-Gheethi AA, Hashim MKA (2017) An overview of the utilisation of microalgae biomass derived from nutrient recycling of wet market wastewater

- and slaughterhouse wastewater. *Int Aquat Res* 9:177–193. <https://doi.org/10.1007/s40071-017-0168-z>
257. Saeed MU, Hussain N, Sumrin A, Shahbaz A, Noor S, Bilal M et al (2022) Microbial bioremediation strategies with wastewater treatment potentialities – a review. *Sci Total Environ* 818:151754. <https://doi.org/10.1016/j.scitotenv.2021.151754>
 258. Al-Jabri H, Das P, Khan S, Thafer M, Abdulquadir M (2021) Treatment of wastewaters by microalgae and the potential applications of the produced biomass—a review. *Water (Switzerland)* 13:27. <https://doi.org/10.3390/w13010027>
 259. Santos MJO, de Souza CO, Marcelino HR (2023) Blue technology for a sustainable pharmaceutical industry: microalgae for bioremediation and pharmaceutical production. *Algal Res* 69:102931. <https://doi.org/10.1016/j.algal.2022.102931>
 260. Srimongkol P, Sangtanoo P, Songserm P, Watsuntorn W, Karnchanatat A (2022) Microalgae-based wastewater treatment for developing economic and environmental sustainability: current status and future prospects. *Front Bioeng Biotechnol* 10:1–2. <https://doi.org/10.3389/fbioe.2022.904046>
 261. Vale F, Sousa CA, Sousa H, Santos L, Simões M (2022) Parabens as emerging contaminants: environmental persistence, current practices and treatment processes. *J Clean Prod* 347:131244. <https://doi.org/10.1016/j.jclepro.2022.131244>
 262. Wollmann F, Dietze S, Ackermann JU, Bley T, Walther T, Steingroewer J et al (2019) Microalgae wastewater treatment: biological and technological approaches. *Eng Life Sci* 19:860–871. <https://doi.org/10.1002/elsc.201900071>
 263. Khan MI, Shin JH, Kim JD (2018) The promising future of microalgae: current status, challenges, and optimization of a sustainable and renewable industry for biofuels, feed, and other products. *Microb Cell Fact* 17:1–21. <https://doi.org/10.1186/s12934-018-0879-x>
 264. Ahmad I, Abdullah N, Koji I, Yuzir A, Mohamad SE (2021) Potential of microalgae in bioremediation of wastewater. *Bull Chem React Eng Catal* 16:413–429. <https://doi.org/10.9767/bcrec.16.2.10616.413-429>
 265. Abdelfattah A, Ali SS, Ramadan H, El-Aswar EI, Eltawab R, Ho SH et al (2023) Microalgae-based wastewater treatment: mechanisms, challenges, recent advances, and future prospects. *Environ Sci Ecotechnology* 13:100205. <https://doi.org/10.1016/j.ese.2022.100205>
 266. Peter AP, Khoo KS, Chew KW, Ling TC, Ho SH, Chang JS et al (2021) Microalgae for biofuels, wastewater treatment and environmental monitoring. *Environ Chem Lett* 19:2891–2904. <https://doi.org/10.1007/s10311-021-01219-6>
 267. Satya ADM, Cheah WY, Yazdi SK, Cheng YS, Khoo KS, Vo DVN et al (2023) Progress on microalgae cultivation in wastewater for bioremediation and circular bioeconomy. *Environ Res* 218:114948. <https://doi.org/10.1016/j.envres.2022.114948>
 268. Cunha C, Silva L, Paulo J, Faria M, Nogueira N, Cordeiro N (2020) Microalgal-based biopolymer for nano- and microplastic removal: a possible biosolution for wastewater treatment. *Environ Pollut* 263:114385. <https://doi.org/10.1016/j.envpol.2020.114385>
 269. Maia C, Sousa CA, Sousa H, Vale F, Simões M (2023) Parabens removal from wastewaters by microalgae – ecotoxicity, metabolism and pathways. *Chem Eng J* 453:139631. <https://doi.org/10.1016/j.cej.2022.139631>
 270. Sutherland DL, Ralph PJ (2019) Microalgal bioremediation of emerging contaminants - opportunities and challenges. *Water Res* 164:114921. <https://doi.org/10.1016/j.watres.2019.114921>
 271. Chia WY, Ying Tang DY, Khoo KS, Kay Lup AN, Chew KW (2020) Nature's fight against plastic pollution: algae for plastic biodegradation and bioplastics production. *Environ Sci Ecotechnol* 4:100065. <https://doi.org/10.1016/j.ese.2020.100065>
 272. Danso D, Chow J, Streita WR (2019) Plastics: environmental and biotechnological perspectives on microbial degradation. *Appl Environ Microbiol* 85:e01095-e1119. <https://doi.org/10.1128/AEM.01095-19>
 273. Kim JW, Park SB, Tran QG, Cho DH, Choi DY, Lee YJ et al (2020) Functional expression of polyethylene terephthalate-degrading enzyme (PETase) in green microalgae. *Microb Cell Fact* 19:97. <https://doi.org/10.1186/s12934-020-01355-8>
 274. Larue C, Sarret G, Castillo-Michel H, Pradas del Real AE (2021) A critical review on the impacts of nanoplastics and microplastics on aquatic and terrestrial photosynthetic organisms. *Small* 17:1–28. <https://doi.org/10.1002/smll.202005834>
 275. Moog D, Schmitt J, Senger J, Zarzycki J, Rexer KH, Linne U et al (2019) Using a marine microalga as a chassis for polyethylene terephthalate (PET) degradation. *Microb Cell Fact* 18:171. <https://doi.org/10.1186/s12934-019-1220-z>
 276. Paladhi AG, Vallinayagam S, Rajendran S, Rathinam V, SharmaVK (2022) Chapter 3: Microalgae: a promising tool for plastic degradation. *Microbes Microb Biotechnol Green Remediat*, pp 575–587. <https://doi.org/10.1016/B978-0-323-90452-0.00049-9>
 277. Benmoussa M (2016) Alcomics for the development of a sustainable microalgae biorefinery. *Single Cell Biol* 5:1. <https://doi.org/10.4172/2168-9431.1000132>
 278. Yoshida S, Hiraga K, Takehana T, Taniguchi I, Yamaji H, Maeda Y et al (2016) A bacterium that degrades and assimilates poly(ethylene terephthalate). *Science* 351:1196–1199. <https://doi.org/10.1126/science.1246359>
 279. Khoironi A, Anggoro S, Sudarno (2019) Evaluation of the interaction among microalgae *Spirulina* sp, plastics polyethylene terephthalate and polypropylene in freshwater environment. *J Ecol Eng* 20:161–173. <https://doi.org/10.12911/22998993/108637>
 280. Gulnaz O, Dincer S (2009) Biodegradation of bisphenol A by *Chlorella vulgaris* and *Aeromonas hydrophila*. *J Appl Biol Sci* 3:79–84
 281. Hirooka T, Nagase H, Uchida K, Hiroshige Y, Ehara Y, Nishikawa JI et al (2005) Biodegradation of bisphenol A and disappearance of its estrogenic activity by the green alga *Chlorella fusca* var. *vacuolata*. *Environ Toxicol Chem* 24:1896–1901. <https://doi.org/10.1897/04-259R.1>
 282. Vale F, Sousa CA, Sousa H, Santos L, Simões M (2022) Impact of parabens on microalgae bioremediation of wastewaters: a mechanistic study. *Chem Eng J* 442:136374. <https://doi.org/10.1016/j.cej.2022.136374>
 283. Kaloudas D, Pavlova N, Penchovsky R (2021) Phycoremediation of wastewater by microalgae: a review. *Environ Chem Lett* 19:2905–2920. <https://doi.org/10.1007/s10311-021-01203-0>
 284. Duran SK, Kumar P, Sandhu SS (2021) A review on microalgae strains, cultivation, harvesting, biodiesel conversion and engine implementation. *Biofuels* 12:91–102. <https://doi.org/10.1080/17597269.2018.1457314>
 285. Tripathi M, Sahu JN, Ganesan P (2016) Effect of process parameters on production of biochar from biomass waste through pyrolysis: a review. *Renew Sustain Energy Rev* 55:467–481. <https://doi.org/10.1016/j.rser.2015.10.122>
 286. Cavalletti E, Romano G, Palma Esposito F, Barra L, Chiaiese P, Balzano S et al (2022) Copper effect on microalgae: toxicity and bioremediation strategies. *Toxics* 10:1–14. <https://doi.org/10.3390/toxics10090527>
 287. Cid A, Herrero C, Torres E, Abalde J (1995) Copper toxicity on the marine microalga *Phaeodactylum tricornutum*: effects on photosynthesis and related parameters. *Aquat Toxicol* 31:165–174. [https://doi.org/10.1016/0166-445X\(94\)00071-W](https://doi.org/10.1016/0166-445X(94)00071-W)
 288. Cai T, Park SY, Li Y (2013) Nutrient recovery from wastewater streams by microalgae: status and prospects. *Renew Sustain Energy Rev* 19:360–369. <https://doi.org/10.1016/j.rser.2012.11.030>

289. Environmental Protection Agency. (2023) The Effects: dead zones and harmful algal blooms. Nutr. Pollut, 1
290. Regmi C, Joshi B, Ray SK, Gyawali G, Pandey RP (2018) Understanding mechanism of photocatalytic microbial decontamination of environmental wastewater. Front Chem 6:1–6. <https://doi.org/10.3389/fchem.2018.00033>
291. Osman MEH, Abo-Shady AM, Gheda SF, Desoki SM, Elshobary ME (2023) Unlocking the potential of microalgae cultivated on wastewater combined with salinity stress to improve biodiesel production. Environ Sci Pollut Res 30:114610–114624. <https://doi.org/10.1007/s11356-023-30370-6>
292. Cao Q, Sun W, Yang T, Zhu Z, Jiang Y, Hu W et al (2022) The toxic effects of polystyrene microplastics on freshwater algae *Chlorella pyrenoidosa* depends on the different size of polystyrene microplastics. Chemosphere 308:136135. <https://doi.org/10.1016/j.chemosphere.2022.136135>
293. Castro-Castellon AT, Horton AA, Hughes JMR, Rampley C, Jeffers ES, Bussi G et al (2022) Ecotoxicity of microplastics to freshwater biota: considering exposure and hazard across trophic levels. Sci Total Environ 816:151638. <https://doi.org/10.1016/j.scitotenv.2021.151638>
294. Li S, Wang P, Zhang C, Zhou X, Yin Z, Hu T et al (2020) Influence of polystyrene microplastics on the growth, photosynthetic efficiency and aggregation of freshwater microalgae *Chlamydomonas reinhardtii*. Sci Total Environ 714:136767. <https://doi.org/10.1016/j.scitotenv.2020.136767>
295. Natarajan L, Soupam D, Dey S, Chandrasekaran N, Kundu R, Paul S et al (2022) Toxicity of polystyrene microplastics in freshwater algae *Scenedesmus obliquus*: effects of particle size and surface charge. Toxicol Rep 9:1953–1961. <https://doi.org/10.1016/j.toxrep.2022.10.013>
296. Nava V, Leoni B (2021) A critical review of interactions between microplastics, microalgae and aquatic ecosystem function. Water Res 188:116476. <https://doi.org/10.1016/j.watres.2020.116476>
297. Parsai T, Figueiredo N, Dalvi V, Martins M, Malik A, Kumar A (2022) Implication of microplastic toxicity on functioning of microalgae in aquatic system. Environ Pollut 308:119626. <https://doi.org/10.1016/j.envpol.2022.119626>
298. Prata JC, da Costa JP, Lopes I, Duarte AC, Rocha-Santos T (2019) Effects of microplastics on microalgae populations: a critical review. Sci Total Environ 665:400–405. <https://doi.org/10.1016/j.scitotenv.2019.02.132>
299. Rani-Borges B, Moschini-Carlos V, Pompêo M (2021) Microplastics and freshwater microalgae: what do we know so far? Aquat Ecol 55:363–377. <https://doi.org/10.1007/s10452-021-09834-9>
300. Sjollem SB, Redondo-Hasselerharm P, Leslie HA, Kraak MHS, Vethaak AD (2016) Do plastic particles affect microalgal photosynthesis and growth? Aquat Toxicol 170:259–261. <https://doi.org/10.1016/j.aquatox.2015.12.002>
301. Wu X, Wu H, Zhang A, Sekou K, Li Z, Ye J (2022) Influence of polystyrene microplastics on levofloxacin removal by microalgae from freshwater aquaculture wastewater. J Environ Manage 301:113865. <https://doi.org/10.1016/j.jenvman.2021.113865>
302. Wang W, Liu H, Liu H, Chen J, Xu X, Xia J et al (2023) Effects of polystyrene microparticles on growth and physiological metabolism of microalgae *Scenedesmus obliquus*. Sustain 15:11223. <https://doi.org/10.3390/su151411223>
303. Lagarde F, Olivier O, Zanella M, Daniel P, Hiard S, Caruso A (2016) Microplastic interactions with freshwater microalgae: hetero-aggregation and changes in plastic density appear strongly dependent on polymer type. Environ Pollut 215:331–339. <https://doi.org/10.1016/j.envpol.2016.05.006>
304. Priya AK, Jalil AA, Dutta K, Rajendran S, Vasseghian Y, Karimi-Maleh H et al (2022) Algal degradation of microplastic from the environment: mechanism, challenges, and future prospects. Algal Res 67:102848. <https://doi.org/10.1016/j.algal.2022.102848>
305. Ugya AY, Meguellati K, Aliyu AD, Abba A, Musa MA (2022) Microplastic stress induce bioresource production and response in microalgae: a concise review. Environ Pollut Bioavailab 34:51–60. <https://doi.org/10.1080/26395940.2022.2033138>
306. Demir-Yilmaz I, Yakovenko N, Roux C, Guiraud P, Collin F, Coudret C et al (2022) The role of microplastics in microalgae cells aggregation: a study at the molecular scale using atomic force microscopy. Sci Total Environ 832:155036. <https://doi.org/10.1016/j.scitotenv.2022.155036>
307. Cunha C, Faria M, Nogueira N, Ferreira A, Cordeiro N (2019) Marine vs freshwater microalgae exopolymers as biosolutions to microplastics pollution. Environ Pollut 249:372–380. <https://doi.org/10.1016/j.envpol.2019.03.046>
308. Govindan N, Bhuyar P, Feng HX, Rahim MHA, Maniam GP, Sundararaju S et al (2021) Evaluation of microalgae's plastic biodeterioration property by a consortium of *chlorella* sp. and cyanobacteria sp. Environ Res Eng Manag 77:86–98. <https://doi.org/10.5755/j01.erem.77.3.25317>
309. Cheng YR, Wang HY (2022) Highly effective removal of microplastics by microalgae *Scenedesmus abundans*. Chem Eng J 435:135079. <https://doi.org/10.1016/j.cej.2022.135079>
310. Kumar RV, Kanna GR, Elumalai S (2017) Bioremediation and biodegradation of polyethylene by green photosynthetic microalgae. J Bioremediat Biodegrad 8:1–8. <https://doi.org/10.4172/2155-6199.1000381>
311. Sarmah P, Rout J (2018) Efficient biodegradation of low-density polyethylene by cyanobacteria isolated from submerged polyethylene surface in domestic sewage water. Environ Sci Pollut Res 25:33508–33520. <https://doi.org/10.1007/s11356-018-3079-7>
312. Rempel A, Gutkoski JP, Nazari MT, Biolchi GN, Cavanhi VAF, Treichel H et al (2021) Current advances in microalgae-based bioremediation and other technologies for emerging contaminants treatment. Sci Total Environ 772:144918. <https://doi.org/10.1016/j.scitotenv.2020.144918>
313. Sanniyasi E, Gopal RK, Gunasekar DK, Raj PP (2021) Biodegradation of low-density polyethylene (LDPE) sheet by microalga *Uronema africanum*. Borge Sci Rep. <https://doi.org/10.1038/s41598-021-96315-6>
314. Gopalakrishnan K, Kashian DR (2022) Extracellular polymeric substances in green alga facilitate microplastic deposition. Chemosphere 286:131814. <https://doi.org/10.1016/j.chemosphere.2021.131814>
315. Zhou JL, Yang L, Huang KX, Chen DZ, Gao F (2022) Mechanisms and application of microalgae on removing emerging contaminants from wastewater: a review. Bioresour Technol 364:128049. <https://doi.org/10.1016/j.biortech.2022.128049>
316. Nolte TM, Hartmann NB, Kleijn JM, Garnæs J, van de Meent D, Jan Hendriks A et al (2017) The toxicity of plastic nanoparticles to green algae as influenced by surface modification, medium hardness and cellular adsorption. Aquat Toxicol 183:11–20. <https://doi.org/10.1016/j.aquatox.2016.12.005>
317. Sundbæk KB, Due I, Koch W, Villaro CG, Rasmussen NS, Holdt SL et al (2018) Sorption of fluorescent polystyrene microplastic particles to edible seaweed *Fucus vesiculosus*. J Appl Phycol 30:2923–2927. <https://doi.org/10.1007/s10811-018-1472-8>
318. Peller J, Nevers MB, Byappanahalli M, Nelson C, Ganesh Babu B, Evans MA et al (2021) Sequestration of microfibers and other microplastics by green algae, *Cladophora*, in the US Great Lakes. Environ Pollut 276:116695. <https://doi.org/10.1016/j.envpol.2021.116695>
319. Kiki C, Qiu Y, Wang Q, Ifon BE, Qin D, Chabi K et al (2022) Induced aging, structural change, and adsorption behavior modifications of microplastics by microalgae. Environ Int 66:107382. <https://doi.org/10.1016/j.envint.2022.107382>
320. Anand U, Dey S, Bontempi E, Ducoli S, Vethaak AD, Dey A et al (2023) Biotechnological methods to remove microplastics:

- a review. Environ Chem Lett 21:1787–1810. <https://doi.org/10.1007/s10311-022-01552-4>
321. Hadiyanto H, Khoironi A, Dianratri I, Suherman S, Muhammad F, Vaidyanathan S (2021) Interactions between polyethylene and polypropylene microplastics and *Spirulina* sp. microalgae in aquatic systems. Heliyon 7:e07676. <https://doi.org/10.1016/j.heliyon.2021.e07676>
322. Mitchell SM, Ullman JL, Teel AL, Watts RJ (2014) pH and temperature effects on the hydrolysis of three β -lactam antibiotics: ampicillin, cefalotin and cefoxitin. Sci Total Environ 466–467:547–555. <https://doi.org/10.1016/j.scitotenv.2013.06.027>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.