

# MA 765: Topics in Non-linear Algebra

Taught by Dr. Uwe Nagel

Notes by Narendran

University of Kentucky

Fall 2023

|          |                          |          |
|----------|--------------------------|----------|
| <b>1</b> | <b>Gröbner Basis</b>     | <b>1</b> |
| <b>2</b> | <b>Hilbert Functions</b> | <b>4</b> |

# 1 Gröbner Basis

Denote by  $\leq$  the coordinate wise partial order on  $\mathbb{N}_0^n$ .  $(a_1, \dots, a_n) \leq (b_1, \dots, b_n)$  if  $a_i \leq b_i$  for  $i \in [n]$ .

Divisibility is a partial order on monomials.

**Theorem 1.1** (Dickson's lemma). Every infinite subset of  $\mathbb{N}_0^n$  contains elements  $a, b$  with  $a < b$ .

*Proof.* The proof follows by induction. Let  $\mathcal{M} \subset \mathbb{N}_0^n$  be an infinite subset. For any  $i \in \mathbb{N}_0$  define  $\mathcal{M}_i$ ,

$$\mathcal{M}_i = \{a = (a_1, \dots, a_n) \in \mathbb{N}_0^{n-1} : (a, i) \in \mathcal{M}\}$$

If  $\mathcal{M}_i$  is finite, then look at  $\bigcup_i \in \mathbb{N}_0$ , which has finitely many and atleast one minimal element. Thus there is some  $j \in \mathbb{N}$  such that  $\bigcup_{i=0}^j \mathcal{M}_i$  with  $a \in \mathcal{M}_i$  for some  $i \leq j$ . Hence  $(a, i) < (b, k)$ . □

**Corollary 1.2.** For any  $\phi \neq \mathcal{M} \subset \mathbb{N}_0^n$  the set of minimal elements wrt  $<$  is nonempty and finite.

**1.3.** Monomial order on  $\mathbb{N}_0^n$  is a relation  $<$  such that

1. If  $a \neq b$  then  $a < b$  or  $b < a$ .
2. If  $A < b$  and  $b < c$  then  $a < c$ .
3.  $(0, \dots, 0) < a$  for any  $a \in \mathbb{N}_0^n$ .
4.  $a < b$  then  $a + c < b + c$  for any  $c \in \mathbb{N}_0^n$ .

First two conditions together imply that  $<$  is a total order.

**Remark 1.4.** 1. If  $a < b$ , then  $a < b$ , i.e., any monomial order refines coordinate wise partial order.

2. Any monomial order on  $\mathbb{N}_0^n$  gives total order on monomials in  $k[X_n]$

**Example 1.5.** Lexicographic order: Left most non zero entry of  $a - b$  is positive the  $a >_{lex} b$ .

degree lex order if either  $|a| > |b|$  or  $|a| = |b|$  and right most entry of  $a - b$  is negative.

degree reverse lex order, same as above but right most entry of  $a - b$  is negative.

**Proposition 1.6.** For any monomial order  $<$  on  $\mathbb{N}_0^n$  with  $\phi \neq \mathcal{M} \subset \mathbb{N}_0^n$  has a unique minimal element.

*Proof.* Dickson's lemma gives that  $\mathcal{M}$  has a finite non empty set of minimal elements wrt coordinate wise order. The minimal element of these wrt  $<$  is the desired element. □

**1.7.** Fix a monomial order  $<$  on  $k[X_n]$

1. The initial monomial  $\text{im}_{<}(f)$  of  $f = \sum c_a x^a$  is the largest monomial wrt  $<$  appearing in  $f$  with non zero coefficients.
2. The leading term  $\text{lt}_{<}(f) := c_a x^a$  with  $x^a = \text{im}(f)$

3. The initial ideal of an ideal  $I$  is  $\text{im}(I) = \langle \text{im}(f) : f \in I \rangle$ .

If  $G$  is a generating set for an ideal  $I$  then  $\langle \text{im}(g) : g \in G \rangle \subset \text{im}(I)$  and this inclusion can be strict.

**Proposition 1.8.** Let  $<$  be a monomial order on  $k[X_n]$ . Every ideal  $I$  has a finite subest  $\mathcal{G}$  such that  $\text{im}(I) = \langle \text{im}(g) : g \in \mathcal{G} \rangle$ .

Any such  $\mathcal{G}$  is called a Gröbner Basis of  $I$  wrt  $<$ .

*Proof.* The set of monomials in  $\text{im}(I)$  has a finite and nonempty subset of minimal elements wrt divisibility, say  $m_1, \dots, m_s$ . Thus  $\text{im}(I) = \langle m_1, \dots, m_s \rangle$ . Every monomial in  $\text{im}(I)$  is the initial monomial of some  $f \in I$ . hence there exists  $f_1, \dots, f_s \in I$  with  $\text{im}(f_i) = m_i$ . Thus  $\{f_i\}$  is a Gröbner basis.  $\square$

**Theorem 1.9.** If  $\mathcal{G}$  is a Gröbner basis of  $I$ , then  $I = \langle \mathcal{G} \rangle$

Note that Hilbert's basis theorem is a simple corollary of this.

*Proof.* We argue by contradiction. By 1.6, choose  $f \in I - \langle \mathcal{G} \rangle$  such that  $\text{im}(f)$  is minimal. Call  $\text{im}(f) = x^b$ .  $x^b \in \text{im}(I) = \langle \text{im}(g) : g \in \mathcal{G} \rangle$ . There exists  $g \in \mathcal{G}$  such that  $\text{im}(g) | x^b$ , say  $x^b = x^c \cdot \text{im}(g)$ .

$\text{im}(f - x^c \lambda g) < x^b = \text{im}(f)$  where  $\lambda = \text{lt}(f)/x^c \text{lt}(g) \in k$ . But  $f - x^c \lambda g \in I - \langle \mathcal{G} \rangle$ , which is a contradiction to the minimality.  $\square$

**Lemma 1.10.** Consider  $f, g_1, \dots, g_s \in k[X_n]$ , with  $g_i \neq 0$ . Then for any minimal order  $<$ , there exists  $q_1, \dots, q_s, r \in k[X_n]$  such that

1.  $f + \sum_{i=1}^s q_i g_i + r$
2.  $\text{im}(f) \geq \text{im}(q_i g_i) \forall i$  (note that  $\text{im}(f) = \text{im}(q_i g_i)$  for some  $i$ )
3.  $\text{im}(r)$  is not divisible by  $\text{im}(g_i)$  for any  $i$ .

We say  $f$  reduces to  $r$  by  $\{g_1, \dots, g_s\}$ .

**1.11 Division Algorithm.** Input:  $f, g_1, \dots, g_s$

Output:  $q_1, \dots, q_s, r$  satisfying the properties 1-3.

1. Set  $r = 0, p = gf, q_1 = \dots = q_s = 0$

2. While  $p \neq 0$  do :

If  $\text{im}(g_i)$  divides  $\text{im}(p)$  for some  $i \in [s]$ , then set  $q_i = q_i + \frac{\text{lt}(p)}{\text{lt}(g_i)}$  and  $p_i = p - \frac{\text{lt}(p)}{\text{lt}(g_i)} g_i$   
else set  $r = r + \text{lt}(p), p = p - \text{lt}(p)$

3. Return  $q_1, \dots, q_s, r$

**Example 1.12.** Consider Lexicographic ordering with  $x > y$  and  $f = x^2y + xy^2 + y^2$  and

$$g_1 = xy - 1, g_2 = y^2 - 1.$$

| $p$                            | $xy - 1$ | $y^2 - 1$ | $r$ |
|--------------------------------|----------|-----------|-----|
| $x^2y + xy^2 + y^2 - x^2y + x$ | $x$      |           |     |
| $xy^2 + y^2 + x - xy^2 + y$    | $y$      |           |     |
| $y^2 + x + y - x$              |          |           | $x$ |
| $y^2 + y - y^2 + 1$            | $1$      |           |     |
| $y + 1 - y$                    |          |           | $y$ |
| $1$                            |          |           | $1$ |

**Corollary 1.13.** Buchberger's criteria Let  $\mathcal{G}$  be a finite subset of  $I$ . Then  $\mathcal{G}$  is a Gröbner basis of  $I$  iff each  $f \in I$  can be reduced to 0 by  $\mathcal{G}$ .

*Proof.* If  $f$  reduces to  $r$  by  $\mathcal{G}$ , then  $\text{im}(g_i)$  does not divide  $\text{im}(r)$ , for all  $i$ . However  $f - r \in \langle \mathcal{G} \rangle$  and  $r \in I$  and  $\mathcal{G}$  is a Gröbner basis of  $I$ . So there exists some  $g \in \mathcal{G}$  such that  $\text{im}(r)$  is divisible by  $\text{im}(g)$ , which forces  $r = 0$ .

Conversely, we have  $f = \sum q_i g_i$  with  $g_i \in \mathcal{G}$  and  $\text{im}(q_i g_i) \leq \text{im}(f)$ . Hence equality for some  $i$  and so  $\text{im}(f) \in \langle \text{im}(g) : g \in \mathcal{G} \rangle$ .  $\square$

**Proposition 1.14.** Let  $<$  be a monomial order. Then

1. Let  $B$  be the set of monomials in  $k[X_n] - \text{im}(I)$ . Then  $\bar{B} \subset k[X_n]/I$  is a  $k$  vsp basis.
2. If  $\mathcal{G}$  is a Gröbner basis of  $I$ , then the remainder of  $f$  by  $\mathcal{G}$  is unique and does not depend on the choice of  $\mathcal{G}$ .

*Proof.* 1. If  $p = \sum \lambda_i m_i \in I$  with  $m_i \in B$ , then  $\text{im}(p) \in \text{im}(I)$ , but  $\text{im}(p) = \text{im}(m_i) \notin I$ . To show  $\bar{B}$  spans  $m \in k[X_n]$  such that  $\bar{m} \notin \text{span}(\bar{B})$ .

Take  $\min\{m\} = m$  where  $m \notin B$ . So we have  $m \in \text{im}(I)$ . There exists  $f \in I$  such that  $\text{im}(f) = m$ . So any monomial in  $f - \text{lt}(f) + I = f - \lambda m + I$  is in  $\text{span}(\bar{B})$ . So  $\lambda m + I = p - f + I \in \text{span}(\bar{B})$ . This leads to a contradiction  $\square$

**Definition 1.15.** 1. For terms  $\lambda x^a, \mu x^b$  ( $\lambda, \mu \in k$ ) denote by

$$\begin{aligned} \gcd(\lambda x^a, \mu x^b) &= \gcd(x^a, x^b) \\ \text{lcm}(\lambda x^a, \mu x^b) &= \text{lcm}(x^a, x^b) \end{aligned}$$

2. For  $0 \neq g, h \in k[x_n]$ , their  $s$ -polynomial (wrt monomial order  $<$ )

$$S(g, h) := \frac{\text{lt}(h)}{\gcd(\text{lt}(h), \text{lt}(g))} g = \frac{\text{lt}(g)}{\gcd(\text{lt}(h), \text{lt}(g))} h.$$

**1.16.** Buchberger's Algorithm for computing Gröbner basis Input:  $f_1, \dots, f_s \in k[X_n]$  in monomial order.

Output: Gröbner basis  $\mathcal{G}$  of  $I = \langle f_1, \dots, f_s \rangle$  wrt  $<$ .

1. Set  $\eta = \langle f_1 \dots, f_s \rangle$
2. Order the elements of  $\mathcal{G}$  as  $f_1, \dots, f_t$
3. For  $1 \leq i < j \leq t$  do: Reduce  $S(g_i, g_j)$  to  $r$  by  $g$ . If  $r \neq 0$ , then set  $\mathcal{G} := \mathcal{G} \cup \{r\}$  and go to step 2.
4. Return  $\mathcal{G}$

**Remark 1.17.** The algorithm computes a Gröbner basis. It terminate because in case  $r \neq 0$ ,  $\text{im}(r) \notin \langle \text{im}(g_1), \dots, \text{im}(g_t) \rangle$ .

**1.18.** Extension to submodules of finitely generated  $k[X_n]$  module So  $F = k[X_n]^r = \bigoplus_{i=1}^r k[X_n]e_i$ .

Monomials in  $G$  are of the form  $x^a e_i$  and terms are  $\lambda x^a e_i$  with  $\lambda \in k$ .

A monomial order on  $FF$  is a total order on the monomials satisfying:

If  $x^a \neq 1$ , then  $m_1 < m_2 \implies m_1 < x^a \cdot 1 < x^a m_2$ , for monomials  $m_i$  in  $F$ .

Given a monomial order on the polynomial ring  $k[X_n]$  and an order on  $\{e_i\}$ . We obtain a monomial ordering on  $F$  by ordering  $\mathbb{N}_0^n \times [r]$  or  $[r] \times \mathbb{N}_0^n$  lexicographically.

Dicksen's lemma can be extended to monomials in  $F$ . We also define  $\text{im}$ , It with respect to  $<$  analogously. There is also a division algorithm.

**Theorem 1.19.** 1. Every submodule  $M$  of  $F$  has finite Gröbner basis and the basis generates  $M$ .

2. If  $B$  is the set of monomials in  $F - \text{im}(M)$ , then  $\bar{B} \subset F/M$  form a  $k$  basis of  $F/M$ .

## 2 Hilbert Functions

**Definition 2.1.** Let  $I \subset k[X_n]$  be a monomial ideal. The Hilbert function of  $A = k[X_n]/I$  (or of  $I$ ) is

$$h_A : \mathbb{N}_0 \rightarrow \mathbb{Z}$$

$$a \mapsto h_A(j) = \dim_k[A]_j$$

where  $[A]_j$  is  $k$  vector space of images of polynomials of degree  $j$ , from  $k[X_n] \rightarrow A$ . (So  $h_A(j) = \text{number of monomials in } [k[X_n]]_j - I$ ).

It's generating function is the Hilbert series

$$H_A(z) = \sum_{j \geq 0} h_A(j) z^j$$

**Example 2.2.** 1. For  $I = 0$ , we get  $h_{k[X_n]}(j) = \binom{n+j-1}{j}$  and so

$$H_{k[X_n]}(z) = \sum_{j \geq 0} \binom{n+j-1}{j} z^j = \frac{1}{(1-z)^n}$$

2. If  $I = \langle x^a \rangle$ , then let  $e = \deg(x^a)$ .

3.

$$h_{k[X_n]/I}(j) = \begin{cases} h_{k[X_n]}(j) & j < e \\ h_{k[X_n]}(j) - h_{k[X_n]}(j - e) & j \geq e \end{cases}$$

Hence

$$H_{k[X_n]/I} = \frac{1 - z^e}{(1 - z)^n}$$

**Theorem 2.3.** For any proper monomial ideal  $I$  of  $k[X_n]$ , the Hilbert series of  $A = k[X_n]/I$  is a rational function of the form

$$H_A(z) = \frac{\kappa_A(z)}{(1 - z)^d}$$

with  $\kappa_A(z) \in \mathbb{Z}[z]$ ,  $\kappa_A(0) = 1$ ,  $\kappa_A(1) \neq 0$ ,  $d \in \mathbb{N}$ . (this expression is unique.)

The dimension of  $A$  is

$$\dim A := d$$

and the multiplicity of  $A$  (or degree of  $I$ ) is

$$\deg(I) = \mathcal{H}_A(1) > 0$$

There is a polynomial called Hilbert polynomial  $p_A$  of  $A$  such that  $h_A(j) = p_A(j)$  if  $j \gg 0$ . If  $d = \dim A > 0$ , then

$$\begin{aligned} p_A(z) &= \frac{\deg(I)}{(d-1)!} z^{d-1} + \text{lower order terms} \\ &= h_0(A) \binom{z+d-1}{d-1} + h_1(A) \binom{z+d-2}{d-2} + \dots + h_{d-1}(A) \binom{z}{0} \end{aligned}$$

where  $h_0(A) = \deg(I)$  and  $h_i(A)$  are integers. Note if  $p_A(z) \neq 0$  polynomial, then  $\deg p_A = d - 1 = \dim A - 1$ .

**Example 2.4.**

$I = 0$ , we get  $\dim k[X_n] = n$  and the multiplicity of  $k[X_n]$  is  $1 = \deg I$ .

$I = \langle x^a \rangle$  with  $\deg x = e$ , then  $\dim k[X_n]/I = n - 1$  and  $\deg(I) = \deg(x^a)$ .

*Proof of 2.3.* Inclusion exclusion principle:  $|\bigcup_{i=1}^s X_i| = \sum_{\phi \neq T \subset [s]} (-1)^{|T|+1} |X_T|$  where  $X_T = \bigcap_{i \in T} X_i$

Let  $I = \langle m_1, \dots, m_s \rangle = I$ , where  $m_i$  are monomials. Denote by  $X_i(j)$  set of degree  $j$  monomials in  $\langle m_i \rangle \subset k[X_n]$ . hence for  $T \subset [s]$ ,  $X_T(j) = \bigcap_{i \in T} X_i(j)$  is the set of deg  $j$  monomials that are divisible by  $m_T = \text{lcm}(m_i)_{i \in T}$ . Define  $e_t := \deg m_T$ . So

$$|X_T(j)| = \begin{cases} 0 & j < e_t \\ \binom{n-1+j-e_t}{n-1} & j \geq e_t \end{cases}$$

Thus

$$\sum_{j \geq 0} |X_T(j)| z^j = \sum_{j \geq e_t} \binom{n-1+j-e_t}{n-1} z^j = \sum_{k \geq 0} \binom{n-1+k}{n-1} z^{k+e_t} = z^{e_t} \frac{1}{(1-z)^n}$$

Since

$$\begin{aligned} h_A(j) &= \binom{n-1+j}{n-1} - \text{number of deg } j \text{ monomials in } I = \langle m_1 \dots, m_s \rangle \\ &= \binom{n-1+j}{n-1} - \left| \bigcup_{i \in [j]} X_i \right| \end{aligned}$$

We get

$$\begin{aligned} H_A(z) &= \sum_{j \geq 0} h_A(j) z^j = \sum_{j \geq 0} \binom{n-1+j}{n-1} z^j + \sum_{T \in [s]} (-1)^{|T|} \sum_{j \geq 0} |X_T(j)| z^j \\ &= \frac{1}{(1-z)^n} + \sum_{T \in [s]} (-1)^T \frac{z^{e_T}}{(1-z)^n} =: \frac{g(z)}{(1-z)^n} \end{aligned}$$

When  $g(z) \in \mathbb{Z}[z]$ , with  $g(0) = 1$  writing  $g(z) = (1-z)^v \kappa_A(z)$  with  $\kappa_A(z) \in \mathbb{Z}[z]$ , suitable  $v \in \mathbb{N}$  and  $\kappa_A(1) \neq 0, \kappa_A(0) = 1$ . Hence  $H_A(z) = \frac{\kappa_A(z)}{(1-z)^d}$  where  $d = n - v$ . Write

$$\kappa_A(z) = \sum_{k=0}^w c_k z^k \text{ with } c_k \in \mathbb{Z}$$

$$\sum_{j \geq 0} h_A(j) z^j = \left( \sum_{k=0}^w c_k z^k \right) \left( \sum_{l \geq 0} \binom{d-1+l}{d-1} z^l \right)$$

comparing coefficients in  $\deg j \gg 0$ , we get

$$\begin{aligned} h_A(j) &= \sum_{k+l=j} c_k \binom{d-1+l}{d-1} = \sum_{k=0}^w c_k \underbrace{\binom{d-1+j-k}{d-1}}_{\text{polynomial in } j-k \text{ variables of deg } d-1} \\ &= \underbrace{\left( \sum_{k=0}^w c_k \right)}_{\kappa_A(1)} \binom{j}{d-1} + \text{lower order terms} \\ &=: p_A(j) \text{ (Hilbert polynomial)} \end{aligned}$$

If  $h_A(j) = 0$  whenever  $j \gg 0$ , then by definition  $0 = d = \dim A$  and in this case  $p_A$  is the zero polynomial. Hence if  $d > 0$ , then  $h_A(j) > 0$  if  $j \gg 0$  and so the leading coefficient of  $p_A(z)$  must be positive, i.e.,  $\kappa_A(1) > 0$   $\square$

**Definition 2.5.** A monomial order is called degree compatible if

$$\deg(x^a) > \deg(x^b) \implies x^a > x^b$$

for any two monomials.

**2.6.** For any ideal  $I \subset k[X_n]$  and any  $t \in \mathbb{Z}$ , set

$$I_{\leq t} = \{f \in I : \deg f \leq t\}$$

It is a  $k$ -subspace of  $k[X_n]$ . Write  $\text{Mon}(k[X_n])$  for the set of monomials in  $k[X_n]$ .

**Lemma 2.7.** Let  $<$  be a degree compatible monomial order. For any ideal  $I \subset k[X_n] = S$ , one has

$$\dim_k \frac{k[X_n]_{\leq t}}{I_{\leq t}} = \text{number of monomials in } \frac{k[X_n]_{\leq t}}{\text{im}_{<}(I)} \\ = | \text{Mon}(k[X_n])_{\leq t} - \text{im}_{<}(t) |$$

*Proof.*  $B := \text{Mon}(k[X_n])_{\leq t} - \text{im}_{<}(I)$ . We claim

$$\bar{B} \subset \frac{k[X_n]_{\leq t}}{I_{\leq t}} \text{ is a } k\text{-basis}$$

$\overline{\text{Mon}(k[X_n]) - \text{im}(I)}$  is a  $k$ -basis of  $k[X_n] - \text{im}(I)$ , by 1.14.  $\bar{B}$  spans remainder of any  $F \in k[X_n]$  upon dividing by Gröbner basis of  $I$  wrt  $<$  satisfies  $\deg r \leq \deg f$ .  $\square$

For  $I \subset k[X_n]$ , the affine Hilbert function of  $A = k[X_n]/I$  is

$$h_A^a : \mathbb{N}_0 \longrightarrow \mathbb{Z} \\ j \mapsto h_A^a(j) = \dim_k \frac{k[X_n]_{\leq j}}{I_{\leq j}}$$

**Lemma 2.8.** For any degree compatible monomial order  $<$  on  $I$ , one has

$$h_{\frac{k[X_n]}{\text{im}_{<}(I)}}(j) = h_A^a(j) - h_A^1(j-1)$$

where  $A = k[X_n]/I$ .

*Proof.* By definition, we have

$$h_{\frac{k[X_n]}{\text{im}(I)}}(j) = | \text{Mon}(k[X_n])_j - \text{im}(I) |$$

$$2.7 \implies h_A^a(j) = | \text{Mon}(k[X_n])_{\leq j} - \text{im}(I) |$$

$\square$

**Remark 2.9.** If  $A \neq 0$ ,  $h_A^a(0) = 1$ . Then by 2.8 we have  $h_A^a(j) = \sum_{k=0}^j h_{k[X_n]/\text{im}(I)}(k)$ . It follows that generating function of  $h_A^a$  and  $h_{\frac{k[X_n]}{\text{im}_{<}(I)}}$  have analogous properties

**2.10.** We define dimension of  $A$ ,

$$\dim \frac{k[X_n]}{I} := \dim_k \frac{k[X_n]}{\text{im}(I)}$$

and the degree

$$\deg(I) = \deg(\text{im}(I))$$

where  $<$  is a degree compatible monomial order.

**2.11.** Let  $G = (G, +)$  be an abelian group. A  $G$ -graded ring  $R$  is a family of subgroups  $([R]_{a \in G}) \leq (R, +)$  such that

1.  $R = \oplus_{a \in G} [R]_a$  ( as  $\mathbb{Z}$ -modules )



$$2. [R]_a \cdot [R]_b \subset [R]_{a+b}$$

The elements of  $[R]_a$  are called homogeneous of degree  $a$ .

**Example 2.12.** Fine or  $\mathbb{Z}$ -grading of  $k[X_n]$ , where  $G = \mathbb{Z}^n$

$$[S]_a = \begin{cases} 0 & \text{if some } a_i < 0 \\ \{\lambda x^a : \lambda \in k\} & \end{cases}$$

**Definition 2.13.**  $R = G$ -graded ring

1.  $G$ -graded  $R$ -module is an  $R$ -module  $M$  with a decomposition  $([M]_a)_{a \in G}$  such that  $M = \bigoplus_{a \in G} [M]_a$  and  $[R]_a \cdot [M]_b \subset [M]_{a+b}$ .
2. A  $G$ -graded or simply graded submodule of such a graded  $M$  is a graded submodule  $N \subset M$  such that

$$[N]_a \subset [M]_a$$

**Lemma 2.14.** For an arbitrary submodule  $N$  of a  $G$  graded  $R$ -module  $M$  the following are equivalent

1.  $N$  is a graded submodule
2.  $N$  has a generating set consisting of homogeneous elements
3. If  $m = \sum_{a \in G} m_a$  with  $m_a \in [M]_a$ , then  $m \in N$  iff each  $m_a \in N$ .
4.  $M/N$  is a  $G$ -graded  $R$ -module with grading  $[M/N]_a := \frac{[M]_a + N}{N}$ .

*Proof.*

□

**Example 2.15.** 1.  $M$  and  $N$  are  $G$ -graded, then so is  $M \oplus N$  with grading  $[M \oplus N]_a := [M]_a \oplus [N]_a$  (as  $R$  modules). So if  $R$  is  $G$  graded, then so is  $R^n$ .

2.  $I \subset k[X_n]$  is a  $\mathbb{Z}^n$  graded submodule if  $I$  is a monomial ideal.

3. A  $\mathbb{Z}$ -graded or homogeneous ideal of  $k[X_n]$ , is an ideal that has generating set consisting of homogeneous polynomials. In that case  $k[X_n]/I$  is a graded module.

**2.16.** A homomorphism of  $G$ -graded modules or a  $G$ -graded homomorphism is a  $R$ -module homomorphism  $\phi : M \rightarrow N$  that is degree preserving,  $\phi([M]_a) \subset [N]_a$ .

For any  $a \in G$  and a  $G$ -graded module  $M$ , the module  $M(a)$  has the same module structure as  $M$ , but grading given by

$$[M(a)]_b := [M]_{a+b}$$

$M(a)$  is a degree  $a$  shift of  $M$ . (Note here that the convention is opposite of that in algebraic topology.)

**Example 2.17.** 1. Consider  $k[X_n]$  with standard grading.

$$\begin{aligned} \phi : k[X_n] &\rightarrow k[X_n] \\ f &\mapsto x_1^2 f \end{aligned}$$

is not a graded homomorphism. However define

$$\begin{aligned}\psi &: k[X_n](-2) \rightarrow k[X_n] \\ f &\mapsto x_1^2 f\end{aligned}$$

then  $f \in k[X_n](-2)$  has degree  $\deg f + 2$ . So  $f \in [k[X_n](-2)]_{\deg f + 2}$ .

2. For any  $a \neq \text{in } G$ ,  $R(a)$  is not a graded ring, (because identity is not in 0 dimension), but it is a graded  $R$ -module.

**Lemma 2.18.** If  $\phi : M \rightarrow N$  is a homomorphism of graded modules, then  $\ker \phi$ ,  $\text{im} \phi$ ,  $\text{coker} \phi$  are graded modules.

*Proof.*  $[\ker \phi]_a = \ker \phi \cap [M]_a$ , and  $[\text{im} \phi]_a = \text{im} \phi \cap [N]_a$ .

□

**Example 2.19.** 1. If  $M$  is a  $\mathbb{Z}$  graded module with generators  $m_1, \dots, m_t$  where  $d_i = \deg m_i$ , then

$$\begin{aligned}\phi &: \bigoplus_{i=1}^t R(-d_i) \rightarrow M \\ \begin{bmatrix} r_1 \\ \vdots \\ r_t \end{bmatrix} &\mapsto \sum_{i=1}^t r_i m_i\end{aligned}$$

is a homomorphism of graded  $R$ -modules and is surjective.

2. Consider  $I = \langle x^3, xy, y^4 \rangle \subset k[x, y] + S$  with standard grading.

$$\begin{aligned}\phi &: S(-3) \oplus S(-2) \oplus S(-4) \rightarrow I \\ \begin{bmatrix} f_1 \\ f_2 \\ f_3 \end{bmatrix} &\mapsto f_1 x^3 + f_2 xy + f_3 y^4\end{aligned}$$

$$\ker \phi : \left\langle \begin{bmatrix} y \\ -x^2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ y^3 \\ -x \end{bmatrix} \right\rangle \xleftarrow[\text{graded hom}]{\simeq} S(-4) \oplus S(-5)$$

There exists an exact sequence,

$$\begin{array}{ccccccc} & & & S(-3) & & & \\ & & & \oplus & & & \\ 0 \rightarrow & \begin{matrix} S(-4) \\ S(-5) \end{matrix} \oplus & \xrightarrow{\begin{bmatrix} y & 0 \\ -x^2 & y^3 \\ 0 & -x \end{bmatrix}} & \begin{matrix} S(-2) \\ \oplus \\ S(-4) \end{matrix} & \xrightarrow{\begin{bmatrix} x^3 & xy & y^4 \end{bmatrix}} & S & \rightarrow S/I \rightarrow 0 \end{array}$$

**Definition 2.20.** For any  $\mathbb{Z}$  graded module  $M$  over  $k[X_n]$  its Hilbert function is

$$\begin{aligned}h_m &: \mathbb{Z} \rightarrow \mathbb{Z} \\ j &\mapsto h_m(j) := \dim_k [M]_j\end{aligned}$$

assuming  $[M]_j$  is finitely generated for all  $j$ .

**Remark 2.21.** For any monomial ideal  $I \subset k[X_n]$ , 2.1 and 2.20 agree.

$$\dim_k[k[X_n]/I]_j = |\text{Mon}(k[X_n])_j - I|$$

**Proposition 2.22.** For every graded submodule  $M$  of a finitely generated free  $k[X_n]$  module  $F$  and any monomial order  $<$  of  $F$ ,

$$h_{F/M}(j) = \frac{h_F(j)}{\text{im}(M)} \quad \forall j \in \mathbb{Z}$$

**Corollary 2.23.** For any finitely generated  $k[X_n]$  submodule  $m \neq 0$ , Hilbert series is of the form

$$\begin{aligned} H_M(z) &:= \sum_{j \in \mathbb{Z}} h_M(j) z^j \\ &= \frac{\kappa_M(z) z^t}{(1-z)^d} \end{aligned}$$

where  $\kappa_M(z) \in \mathbb{Z}[z]$ ,  $\kappa_M(0) \neq 0$ ,  $\kappa_M(1) > 0$ ,  $t \in \mathbb{Z}$ .

There is a Hilbert polynomial  $p_M(z) \in \mathbb{Q}(z)$  such that

$$h_M(j) = p_M(j) \quad j \gg 0$$

(krull) dimension of  $M$  is defined as

$$\dim M = d$$

and the degree is

$$\deg(M) = \kappa_M(1)$$

*Proof.* Let  $M$  be generated by  $m_1, \dots, m_t$  of degree  $d_1, \dots, d_t$  respectively. Define

$$\begin{aligned} \phi : \bigoplus_{i=1}^t S(-d_i) &\twoheadrightarrow M \\ f &\mapsto x_1^2 f \end{aligned}$$

is not a graded homomorphism. However define

$$\begin{aligned} \psi : k[X_n](-2) &\rightarrow k[X_n] \\ \begin{bmatrix} f_1 \\ \vdots \\ f_t \end{bmatrix} &\mapsto \sum f_i m_i \end{aligned}$$

Set  $N = \ker \phi$  and we have a short exact sequence  $0 \rightarrow N \rightarrow F \rightarrow M \rightarrow 0$  and by rank nullity theorem we have  $h_M(j) = h_F(j) - h_N(j)$ .  $S$  and  $F$  have desired Hilbert series which are rational functions. So it is enough to show the same for  $N$ , which is same as  $h_{F/\text{im}(N)}$ .

$\text{im}(N)$  is generated by monomials. So

$$\frac{F}{\text{im}(N)} \cong \bigoplus_{i=1}^t \underbrace{\left( \frac{S}{J_i} \right)}_{\text{has the desired properties}} (-d_i)$$

for monomial ideals  $J_i$ . □

**Example 2.24.** 1.  $S = k[X_n], F = \bigoplus_{i=1}^t S(-d_i)$ . Then

$$H_{S(-d_i)}(z) = \frac{z_i^d}{(1-z)^n}$$

2. Consider  $I = \langle x^3, xy, y^4 \rangle \subset k[x, y] + S$  with standard grading.  $A = S/I$ . We have the exact sequence,

$$\begin{array}{ccccccc} & & S(-3) & & & & \\ & & \oplus & & & & \\ 0 \rightarrow & \oplus & \rightarrow S(-2) & \xrightarrow{\begin{smallmatrix} \phi \\ [x^3 \quad xy \quad y^4] \end{smallmatrix}} & S & \rightarrow A = S/I \rightarrow 0 \\ & S(-5) & \oplus & & & & \\ & & S(-4) & & & & \end{array}$$

$$\begin{aligned} H_A(z) &= H_S(z) - H_{S(-3) \oplus S(-2) \oplus S(-4)}(z) + H_{S(-4) \oplus S(-5)}(z) \\ &= H_S(z) - H_{S(-3)}(z) - H_{S(-2)}(z) + H_{S(-4)}(z) + H_{S(-5)}(z) - H_{S(-4)}(z) \\ &= \frac{1 - z^3 - z^2 + z^5}{(1-z)^2} \\ &= 1 + 3z + 2z^2 + z^3 \end{aligned}$$

So  $\dim A = 0$  and  $\deg M = 6$  (polynomial evaluated at 1).