

# PSTAT 5A Practice Worksheet 3 - SOLUTIONS

## Comprehensive Review: Probability, Counting, and Conditional Probability

Solution Key

2025-07-29

### Table of contents

<b>1</b>	<b>Section A: Probability - SOLUTIONS</b>	<b>1</b>
<b>2</b>	<b>Section B: Permutations and Combinations - SOLUTIONS</b>	<b>2</b>
<b>3</b>	<b>Section C: Conditional Probability - SOLUTIONS</b>	<b>2</b>
<b>4</b>	<b>Section D: Review - SOLUTIONS</b>	<b>4</b>

## 1 Section A: Probability - SOLUTIONS

*Estimated time: 8 minutes*

### Problem A1: Probability Distributions - SOLUTION

For a valid probability distribution, two conditions must be met:

1. All probabilities must be non-negative (  $\geq 0$  )
2. The sum of all probabilities must equal 1

**Analysis:**

**(a) Invalid**

- $\text{Sum} = 0.3 + 0.3 + 0.3 + 0.2 + 0.1 = 1.2 > 1$  The probabilities sum to more than 1, violating the second condition.

**(b) Valid**

- $\text{Sum} = 0 + 0 + 1 + 0 + 0 = 1$  All probabilities are non-negative and sum to 1. This represents a class where everyone receives a C.

**(c) Invalid**

- $\text{Sum} = 0.3 + 0.3 + 0.3 + 0 + 0 = 0.9 < 1$  The probabilities sum to less than 1, violating the second condition.

**(d) Invalid**

- Contains  $F = -0.1 < 0$  Although the sum would equal 1.0, the probability for grade F is negative, violating the first condition.

(e) **Valid**

- $\text{Sum} = 0.2 + 0.4 + 0.2 + 0.1 + 0.1 = 1.0$  All probabilities are non-negative and sum to 1.

(f) **Invalid**

- Contains  $B = -0.1 < 0$  Although the sum equals 1.0, the probability for grade B is negative, violating the first condition.

## 2 Section B: Permutations and Combinations - SOLUTIONS

*Estimated time: 15 minutes*

### Problem B1: Permutations and Combinations - SOLUTION

**Part (a):** How many 6-character passwords can be formed using 3 specific letters and 3 specific digits if repetitions are not allowed and letters must come before digits?

**Solution:** Since letters must come before digits, we have a fixed structure: LLL DDD

- Step 1: Arrange 3 letters in the first 3 positions
  - This is a permutation:  $P(3,3) = 3! = 6$  ways
- Step 2: Arrange 3 digits in the last 3 positions
  - This is a permutation:  $P(3,3) = 3! = 6$  ways
- Step 3: Apply multiplication principle
  - Total passwords  $= 6 \times 6 = \mathbf{36}$  passwords

**Part (b):** If the team wants to select 4 people from 12 employees to form a security committee where order doesn't matter, how many ways can this be done?

**Solution:** Since order doesn't matter, this is a combination problem.

$$C(12, 4) = \binom{12}{4} = \frac{12!}{4!(12-4)!} = \frac{12!}{4! \cdot 8!}$$

$$= \frac{12 \times 11 \times 10 \times 9}{4 \times 3 \times 2 \times 1} = \frac{11880}{24} = \mathbf{495 \text{ ways}}$$

## 3 Section C: Conditional Probability - SOLUTIONS

*Estimated time: 15 minutes*

### Problem B1: Conditional Probability and Medical Testing - SOLUTION

**Given Information:**

- $P(\text{has variant}) = 0.03$
- $P(\text{test positive} \mid \text{has variant}) = 0.95$  (sensitivity)
- $P(\text{test negative} \mid \text{no variant}) = 0.92$  (specificity)

- Therefore:  $P(\text{test positive} \mid \text{no variant}) = 1 - 0.92 = 0.08$

**Part (a):** What is the probability that a randomly selected person tests positive?

**Solution:**

Using the Law of Total Probability:

$$P(\text{test positive}) = P(\text{test positive} \mid \text{has variant}) \times P(\text{has variant}) + P(\text{test positive} \mid \text{no variant}) \times P(\text{no variant})$$

$$\begin{aligned} P(\text{test positive}) &= 0.95 \times 0.03 + 0.08 \times 0.97 \\ &= 0.0285 + 0.0776 = \mathbf{0.1061} \end{aligned}$$

**Part (b):** If someone tests positive, what is the probability they actually have the variant?

**Solution:** Using Bayes' Theorem:

$$\begin{aligned} P(\text{has variant} \mid \text{test positive}) &= \frac{P(\text{test positive} \mid \text{has variant}) \times P(\text{has variant})}{P(\text{test positive})} \\ &= \frac{0.95 \times 0.03}{0.1061} = \frac{0.0285}{0.1061} = \mathbf{0.2686} \end{aligned}$$

**Part (c):** If someone tests negative, what is the probability they actually don't have the variant?

**Solution:** First, find  $P(\text{test negative})$ :

$$P(\text{test negative}) = 1 - P(\text{test positive}) = 1 - 0.1061 = 0.8939$$

Using Bayes' Theorem:

$$\begin{aligned} P(\text{no variant} \mid \text{test negative}) &= \frac{P(\text{test negative} \mid \text{no variant}) \times P(\text{no variant})}{P(\text{test negative})} \\ &= \frac{0.92 \times 0.97}{0.8939} = \frac{0.8924}{0.8939} = \mathbf{0.9983} \end{aligned}$$

**Part (d) [Challenge]:** Two consecutive positive tests - what is the probability they actually have the variant?

**Solution:** Assuming independence between tests:

$$\begin{aligned} P(\text{two positive} \mid \text{has variant}) &= 0.95^2 = 0.9025 \\ P(\text{two positive} \mid \text{no variant}) &= 0.08^2 = 0.0064 \end{aligned}$$

$$P(\text{two positive}) = 0.9025 \times 0.03 + 0.0064 \times 0.97 = 0.027075 + 0.006208 = 0.033283$$

$$P(\text{has variant} \mid \text{two positive}) = \frac{0.027075}{0.033283} = \mathbf{0.8134}$$

### Problem C1: Advanced Counting with Restrictions - SOLUTION

**Part (a):** How many valid meal combinations are possible?

**Solution:** We need to consider cases based on the restrictions.

#### Case 1: Seafood appetizer is chosen

- 1 appetizer option (seafood)
- 7 main course options (cannot choose vegetarian)
- 5 dessert options
- Combinations:  $1 \times 7 \times 5 = 35$

#### Case 2: Non-seafood appetizer + chocolate dessert

- 5 appetizer options (non-seafood)
- 3 main course options (only beef or chicken allowed with chocolate)
- 1 dessert option (chocolate)
- Combinations:  $5 \times 3 \times 1 = 15$

#### Case 3: Non-seafood appetizer + non-chocolate dessert - 5 appetizer options (non-seafood)

- 8 main course options (no restrictions)
- 4 dessert options (non-chocolate)
- Combinations:  $5 \times 8 \times 4 = 160$

**Total valid combinations:**  $35 + 15 + 160 = \mathbf{210}$  combinations

**Part (b):** If customers choose randomly among valid combinations, what is the probability someone chooses the chocolate dessert?

**Solution:** Combinations with chocolate dessert: 15 (from Case 2 above) Total valid combinations: 210

$$P(\text{chocolate dessert}) = \frac{15}{210} = \frac{1}{14} = \mathbf{0.0714}$$

## 4 Section D: Review - SOLUTIONS

*Estimated time: 12 minutes*

### Problem B3: Daily Expenses - SOLUTION

**Given:**

- Coffee: Mean = \$1.40, SD = \$0.30
- Muffin: Mean = \$2.50, SD = \$0.15

- Prices are independent

**Part (a):** What is the mean and standard deviation of the amount she spends on breakfast daily?

**Solution:** For the sum of independent random variables:

**Mean of daily expenses:**

$$E[\text{Daily}] = E[\text{Coffee}] + E[\text{Muffin}] = \$1.40 + \$2.50 = \mathbf{\$3.90}$$

**Variance of daily expenses:**

$$\text{Var}[\text{Daily}] = \text{Var}[\text{Coffee}] + \text{Var}[\text{Muffin}] = (0.30)^2 + (0.15)^2 = 0.09 + 0.0225 = 0.1125$$

**Standard deviation of daily expenses:**

$$SD[\text{Daily}] = \sqrt{0.1125} = \mathbf{\$0.3354}$$

**Part (b):** What is the mean and standard deviation of the amount she spends on breakfast weekly (7 days)?

**Solution:** For the sum of 7 independent daily expenses:

**Mean of weekly expenses:**

$$E[\text{Weekly}] = 7 \times E[\text{Daily}] = 7 \times \$3.90 = \mathbf{\$27.30}$$

**Variance of weekly expenses:**

$$\text{Var}[\text{Weekly}] = 7 \times \text{Var}[\text{Daily}] = 7 \times 0.1125 = 0.7875$$

**Standard deviation of weekly expenses:**

$$SD[\text{Weekly}] = \sqrt{0.7875} = \mathbf{\$0.8874}$$