

CEE 260/MIE 273: Probability and Statistics in Civil Engineering

Lecture 3a: Introduction: Random variables

Prof. Oke

September 19, 2025

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

- Bayes' Theorem for two events:

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

- Bayes' Theorem for two events:

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

- Bayes' Theorem for two events:

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)} \quad (2)$$

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

- Bayes' Theorem for two events:

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)} \quad (2)$$

For multiple events:

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

- Bayes' Theorem for two events:

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)} \quad (2)$$

For multiple events:

$$P(E_1|A) = \frac{P(A|E_1)P(E_1)}{P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n)} \quad (3)$$

M2c Recap: Conditional Probability and Bayes' Theorem

- Total probability:

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n) \quad (1)$$

- Bayes' Theorem for two events:

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)} \quad (2)$$

For multiple events:

$$P(E_1|A) = \frac{P(A|E_1)P(E_1)}{P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \cdots + P(A|E_n)P(E_n)} \quad (3)$$

$$= \frac{P(A|E_1)P(E_1)}{P(A)} \quad (4)$$

Module 3: Probability Distributions

Module 3: Probability Distributions

- M3a: Introduction: Random Variables

Module 3: Probability Distributions

- M3a: Introduction: Random Variables
- M3b: Normal Distribution

Module 3: Probability Distributions

- M3a: Introduction: Random Variables
- M3b: Normal Distribution
- M3c: Lognormal and Exponential Distributions

Module 3: Probability Distributions

- M3a: Introduction: Random Variables
- M3b: Normal Distribution
- M3c: Lognormal and Exponential Distributions
- M3d: Binomial Distribution

Module 3: Probability Distributions

- M3a: Introduction: Random Variables
- M3b: Normal Distribution
- M3c: Lognormal and Exponential Distributions
- M3d: Binomial Distribution
- M3e: Poisson Distribution

Module 3: Probability Distributions

- M3a: Introduction: Random Variables
- M3b: Normal Distribution
- M3c: Lognormal and Exponential Distributions
- M3d: Binomial Distribution
- M3e: Poisson Distribution
- M3f: Joint Distributions and further topics

Objectives and outline

Objectives and outline

- Understand random variables

Objectives and outline

- Understand random variables
- Distinguish between discrete and continuous random variables

Objectives and outline

- Understand random variables
- Distinguish between discrete and continuous random variables
- Compute measures of centrality and dispersion, as well as sketch PMFs, PDFs and CDFs

Objectives and outline

- Understand random variables
- Distinguish between discrete and continuous random variables
- Compute measures of centrality and dispersion, as well as sketch PMFs, PDFs and CDFs

Objectives and outline

- Understand random variables
- Distinguish between discrete and continuous random variables
- Compute measures of centrality and dispersion, as well as sketch PMFs, PDFs and CDFs

Random variables

Random variables

Definitions

- A random variable (r.v.) represents the values of the outcomes in a sample space

Random variables

Definitions

- A random variable (r.v.) represents the values of the outcomes in a sample space (e.g. the outcome of die roll: $X = \{1, 2, 3, 4, 5, 6\}$)

Random variables

Definitions

- A random variable (r.v.) represents the values of the outcomes in a sample space (e.g. the outcome of die roll: $X = \{1, 2, 3, 4, 5, 6\}$)
- A random variable is a function that uniquely maps events in a sample space to the set of real numbers.

A random variable X may be:

Random variables

Definitions

- A random variable (r.v.) represents the values of the outcomes in a sample space (e.g. the outcome of die roll: $X = \{1, 2, 3, 4, 5, 6\}$)
- A random variable is a function that uniquely maps events in a sample space to the set of real numbers.

A random variable X may be:

- *Discrete*
- *Continuous*

Describing random variables

Describing random variables

Central values

- Mean

Describing random variables

Central values

- Mean
- Median

Describing random variables

Central values

- Mean
- Median
- Mode

Describing random variables

Central values

- Mean
- Median
- Mode

Describing random variables

Central values

- Mean
- Median
- Mode

Measures of dispersion

- Variance

Describing random variables

Central values

- Mean
- Median
- Mode

Measures of dispersion

- Variance
- Standard deviation

Describing random variables

Central values

- Mean
- Median
- Mode

Measures of dispersion

- Variance
- Standard deviation
- Coefficient of variation (COV)

Probability distribution

Probability distribution

A probability distribution governs the values of a random variable.

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF **discrete random variable**

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF **discrete random variable**
- probability density function, PDF

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF **discrete random variable**
- probability density function, PDF

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF **discrete random variable**
- probability density function, PDF **continuous random variable**

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF **discrete random variable**
- probability density function, PDF **continuous random variable**
- cumulative distribution function, CDF

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF **discrete random variable**
- probability density function, PDF **continuous random variable**
- cumulative distribution function, CDF

Probability distribution

A probability distribution governs the values of a random variable.
It can be described by the following functions:

- probability mass function, PMF discrete random variable
- probability density function, PDF continuous random variable
- cumulative distribution function, CDF discrete/continuous random variable

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X$$

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv$$

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x)$$

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

The CDF satisfies the basic axioms of probability:

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

The CDF satisfies the basic axioms of probability:

① $F_X(-\infty) = 0$ and $F_X(\infty) = 1$

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

The CDF satisfies the basic axioms of probability:

- ① $F_X(-\infty) = 0$ and $F_X(\infty) = 1$
- ② $F_X(x) \geq 0 \quad \forall x$ and is nondecreasing with x .

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

The CDF satisfies the basic axioms of probability:

- ① $F_X(-\infty) = 0$ and $F_X(\infty) = 1$
- ② $F_X(x) \geq 0 \quad \forall x$ and is nondecreasing with x .
- ③ $F_X(x)$ is continuous to the right with x .

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

The CDF satisfies the basic axioms of probability:

- ① $F_X(-\infty) = 0$ and $F_X(\infty) = 1$
- ② $F_X(x) \geq 0 \quad \forall x$ and is nondecreasing with x .
- ③ $F_X(x)$ is continuous to the right with x .

Cumulative distribution function (CDF)

The CDF of a random variable X is given by

$$F_X \equiv P(X \leq x) \quad \text{for all } x \quad (5)$$

The CDF satisfies the basic axioms of probability:

- ① $F_X(-\infty) = 0$ and $F_X(\infty) = 1$
- ② $F_X(x) \geq 0 \quad \forall x$ and is nondecreasing with x .
- ③ $F_X(x)$ is continuous to the right with x .

Note that the symbol \forall means “for all”

Probability mass function (PMF)

Probability mass function (PMF)

The PMF is given by

Probability mass function (PMF)

The PMF is given by

$$p_X(x_i) \equiv P(X = x_i) \quad \forall x \quad (6)$$

Probability mass function (PMF)

The PMF is given by

$$p_X(x_i) \equiv P(X = x_i) \quad \forall x \quad (6)$$

CDF of discrete random
variable



Probability mass function (PMF)

The PMF is given by

$$p_X(x_i) \equiv P(X = x_i) \quad \forall x \quad (6)$$

CDF of discrete random variable

$$\begin{aligned} F_X(x) &= \sum_{x_i \leq x} P(X = x_i) \\ &= \sum_{x_i \leq x} p_X(x_i) \end{aligned}$$



Probability mass function (PMF)



The PMF is given by

$$p_X(x_i) \equiv P(X = x_i) \quad \forall x \quad (6)$$

CDF of discrete random variable

$$\begin{aligned} F_X(x) &= \sum_{x_i \leq x} P(X = x_i) \\ &= \sum_{x_i \leq x} p_X(x_i) \end{aligned}$$

The probability masses in a PMF sum up to 1.



Probability density function (PDF)

Probability density function (PDF)

The PDF is denoted $f_X(x)$ such that the probability of X in the interval $(a, b]$ is:

Probability density function (PDF)

The PDF is denoted $f_X(x)$ such that the probability of X in the interval $(a, b]$ is:

$$P(a < X \leq b) = \int_a^b f_X(x) dx \quad (7)$$

Probability density function (PDF)

The PDF is denoted $f_X(x)$ such that the probability of X in the interval $(a, b]$ is:

$$P(a < X \leq b) = \int_a^b f_X(x) dx \quad (7)$$

CDF of continuous random variable

Probability density function (PDF)

The PDF is denoted $f_X(x)$ such that the probability of X in the interval $(a, b]$ is:

$$P(a < X \leq b) = \int_a^b f_X(x) dx \quad (7)$$

CDF of continuous random variable

$$\begin{aligned} F_X(x) &= P(X \leq x) \\ &= \int_{-\infty}^x f_X(\tau) d\tau \end{aligned}$$

Probability density function (PDF)

The PDF is denoted $f_X(x)$ such that the probability of X in the interval $(a, b]$ is:

$$P(a < X \leq b) = \int_a^b f_X(x) dx \quad (7)$$



CDF of continuous random variable

$$\begin{aligned} F_X(x) &= P(X \leq x) \\ &= \int_{-\infty}^x f_X(\tau) d\tau \end{aligned}$$



It follows that the PDF is the derivative of the CDF:

$$f_X(x) = \frac{dF_X(x)}{dx} \quad (8)$$

PDF (cont.)



PDF (cont.)



The total area under a PDF is 1.

Further derivations

① Continuous case:

$$P(a < X \leq b) = \int_{-\infty}^b f_X(x) dx - \int_{-\infty}^a f_X(x) dx \quad (9)$$

Further derivations

① Continuous case:

$$P(a < X \leq b) = \int_{-\infty}^b f_X(x) dx - \int_{-\infty}^a f_X(x) dx \quad (9)$$

② Discrete case:

$$P(a < X \leq b) = \sum_{x_i \leq b} p_X(x_i) - \sum_{x_i \leq a} p_X(x_i) \quad (10)$$

Further derivations

① Continuous case:

$$P(a < X \leq b) = \int_{-\infty}^b f_X(x) dx - \int_{-\infty}^a f_X(x) dx \quad (9)$$

② Discrete case:

$$P(a < X \leq b) = \sum_{x_i \leq b} p_X(x_i) - \sum_{x_i \leq a} p_X(x_i) \quad (10)$$

③ For all random variables:

$$P(a < X \leq b) = F_X(b) - F_X(a) \quad (11)$$

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months.

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN
 - NOO

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN
 - NOO
 - NON

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN
 - NOO
 - NON
 - NNO

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN
 - NOO
 - NON
 - NNO
 - NNN

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN
 - NOO
 - NON
 - NNO
 - NNN

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.




Figure: PMF

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OON
 - OOO
 - ONO
 - ONN
 - NOO
 - NON
 - NNO
 - NNN

Example 1: Operating condition of bulldozers

Each of 3 bulldozers equally likely to operational or nonoperational after 6 months. Plot the PMF and CDF of the random variable X which represents the operating condition of the bulldozers after 6 months.

- Let the outcomes be O (operational) and N (nonoperational)
- There are $2 \times 2 \times 2 = 8$ possibilities:
 - OOO
 - ONN
 - NOO
 - NON
 - NNO
 - NNN




Figure: PMF




Figure: CDF

Mean and variance

Mean and variance

Mean

Weighted average or expected value

Mean and variance

Mean

Weighted average or expected value

$$E(X) = \sum_i x_i p_X(x_i) \quad \text{discrete case}$$

Mean and variance

Mean

Weighted average or expected value

$$E(X) = \sum_i x_i p_X(x_i) \quad \text{discrete case} \quad (12)$$

Variance

Mean and variance

Mean

Weighted average or expected value

$$E(X) = \sum_i x_i p_X(x_i) \quad \text{discrete case} \quad (12)$$

Variance

In the discrete case:

Mean and variance

Mean

Weighted average or expected value

$$E(X) = \sum_i x_i p_X(x_i) \quad \text{discrete case} \quad (12)$$

Variance

In the discrete case:

$$\text{Var}(X) = \sum_i (x_i - \mu_X)^2 p_X(x_i) \quad (13)$$

Mean and variance

Mean

Weighted average or expected value

$$E(X) = \sum_i x_i p_X(x_i) \quad \text{discrete case} \quad (12)$$

Variance

In the discrete case:

$$\text{Var}(X) = \sum_i (x_i - \mu_X)^2 p_X(x_i) \quad (13)$$

Expanding results in:

$$\text{Var}(X) = E(X^2) - \mu_X^2 \quad (14)$$

Measures of dispersion (cont.)

Measures of dispersion (cont.)

Standard deviation

Measures of dispersion (cont.)

Standard deviation

The standard deviation is convenient as it has the same unit as the random variable:

Measures of dispersion (cont.)

Standard deviation

The standard deviation is convenient as it has the same unit as the random variable:

$$\sigma_X =$$

Measures of dispersion (cont.)

Standard deviation

The standard deviation is convenient as it has the same unit as the random variable:

$$\sigma_X = \sqrt{\text{Var}(X)} \quad (15)$$

Measures of dispersion (cont.)

Standard deviation

The standard deviation is convenient as it has the same unit as the random variable:

$$\sigma_X = \sqrt{\text{Var}(X)} \quad (15)$$

Coefficient of variation

The COV gives the deviation relative to the mean. It is unitless.

Measures of dispersion (cont.)

Standard deviation

The standard deviation is convenient as it has the same unit as the random variable:

$$\sigma_X = \sqrt{\text{Var}(X)} \quad (15)$$

Coefficient of variation

The COV gives the deviation relative to the mean. It is unitless.

$$\delta_X = \frac{\sigma_X}{\mu_X} \quad (16)$$


Example 2: Bulldozers revisited

Example 2: Bulldozers revisited

You are given the PMF (probability mass function) of the operating condition X of bulldozers after 6 months.


Example 2: Bulldozers revisited

You are given the PMF (probability mass function) of the operating condition X of bulldozers after 6 months.



Example 2: Bulldozers revisited


You are given the PMF (probability mass function) of the operating condition X of bulldozers after 6 months.




Find the mean, variance, standard deviation and coefficient of variation of X .

Example 2: Bulldozers revisited (cont.)

Example 2: Bulldozers revisited (cont.)




Example 2: Bulldozers revisited (cont.)



(a) $\mu_X = E(X) = 0 \left(\frac{1}{8}\right) + 1 \left(\frac{3}{8}\right) + 2 \left(\frac{3}{8}\right) + 3 \left(\frac{1}{8}\right) = 1.5.$


Example 2: Bulldozers revisited (cont.)



(a) $\mu_X = E(X) = 0 \left(\frac{1}{8}\right) + 1 \left(\frac{3}{8}\right) + 2 \left(\frac{3}{8}\right) + 3 \left(\frac{1}{8}\right) = 1.5.$

(b) $Var(X) = [0^2 \left(\frac{1}{8}\right) + 1^2 \left(\frac{3}{8}\right) + 2^2 \left(\frac{3}{8}\right) + 3^2 \left(\frac{1}{8}\right)] - (1.5)^2 = 0.75$

Example 2: Bulldozers revisited (cont.)




(a) $\mu_X = E(X) = 0 \left(\frac{1}{8}\right) + 1 \left(\frac{3}{8}\right) + 2 \left(\frac{3}{8}\right) + 3 \left(\frac{1}{8}\right) = 1.5.$

(b) $Var(X) = [0^2 \left(\frac{1}{8}\right) + 1^2 \left(\frac{3}{8}\right) + 2^2 \left(\frac{3}{8}\right) + 3^2 \left(\frac{1}{8}\right)] - (1.5)^2 = 0.75$

(c) $\sigma_X = \sqrt{0.75} = 0.866$

Example 2: Bulldozers revisited (cont.)



(a) $\mu_X = E(X) = 0 \left(\frac{1}{8}\right) + 1 \left(\frac{3}{8}\right) + 2 \left(\frac{3}{8}\right) + 3 \left(\frac{1}{8}\right) = 1.5.$

(b) $Var(X) = [0^2 \left(\frac{1}{8}\right) + 1^2 \left(\frac{3}{8}\right) + 2^2 \left(\frac{3}{8}\right) + 3^2 \left(\frac{1}{8}\right)] - (1.5)^2 = 0.75$

(c) $\sigma_X = \sqrt{0.75} = 0.866$

(d) $\delta_X = \frac{0.866}{1.50} = 0.577$

Mean and variance

Mean and variance

These include the mean, median and mode.

Mean and variance

These include the mean, median and mode.

- Mean: weighted average or expected value

Mean and variance

These include the mean, median and mode.

- Mean: weighted average or expected value

Mean and variance

These include the mean, median and mode.

- Mean: weighted average or expected value

$$E(X) = \int_{-\infty}^{\infty} xf_X(x)dx \quad \text{continuous case} \quad (17)$$

Variance

Mean and variance

These include the mean, median and mode.

- Mean: weighted average or expected value

$$E(X) = \int_{-\infty}^{\infty} xf_X(x)dx \quad \text{continuous case} \quad (17)$$

Variance

In the continuous case:

Mean and variance

These include the mean, median and mode.

- Mean: weighted average or expected value

$$E(X) = \int_{-\infty}^{\infty} xf_X(x)dx \quad \text{continuous case} \quad (17)$$

Variance

In the continuous case:

$$\text{Var}(X) = \int_{-\infty}^{\infty} (x - \mu_X)^2 f_X(x) dx \quad (18)$$

Mean and variance

These include the mean, median and mode.

- Mean: weighted average or expected value

$$E(X) = \int_{-\infty}^{\infty} xf_X(x)dx \quad \text{continuous case} \quad (17)$$

Variance

In the continuous case:


$$\text{Var}(X) = \int_{-\infty}^{\infty} (x - \mu_X)^2 f_X(x) dx \quad (18)$$

Expanding both equations results in:

$$\text{Var}(X) = E(X^2) - \mu_X^2 \quad (19)$$


Example 3: Loaded beam

Example 3: Loaded beam




Consider the beam under a 100-kg load. If the load is equally likely to be placed anywhere along the 10m span of the beam,

Example 3: Loaded beam



Consider the beam under a 100-kg load. If the load is equally likely to be placed anywhere along the 10m span of the beam, then the PDF of the load position X is **uniformly distributed** in $0 < x \leq 10$, i.e.:


Example 3: Loaded beam



Consider the beam under a 100-kg load. If the load is equally likely to be placed anywhere along the 10m span of the beam, then the PDF of the load position X is **uniformly distributed** in $0 < x \leq 10$, i.e.:

$$f_X(c) = \begin{cases} c & 0 < x \leq 10 \end{cases}$$


Example 3: Loaded beam



Consider the beam under a 100-kg load. If the load is equally likely to be placed anywhere along the 10m span of the beam, then the PDF of the load position X is **uniformly distributed** in $0 < x \leq 10$, i.e.:

$$f_X(c) = \begin{cases} c & 0 < x \leq 10 \\ 0 & \text{otherwise} \end{cases} \quad (20)$$

Example 3: Loaded beam



Consider the beam under a 100-kg load. If the load is equally likely to be placed anywhere along the 10m span of the beam, then the PDF of the load position X is **uniformly distributed** in $0 < x \leq 10$, i.e.:

$$f_X(c) = \begin{cases} c & 0 < x \leq 10 \\ 0 & \text{otherwise} \end{cases} \quad (20)$$

- (a) Plot the PDF of X .
- (b) Solve the integral for the CDF and plot.
- (c) Find $P(2 < X \leq 5)$.


Example 3: Loaded beam (cont.)

Example 3: Loaded beam (cont.)

(a) The area under the PDF must be 1. Thus, c must be $\frac{1}{10}$.

Example 3: Loaded beam (cont.)

- (a) The area under the PDF must be 1. Thus, c must be $\frac{1}{10}$.



Example 3: Loaded beam (cont.)

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

$$F_X =$$

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

$$F_X = \int_0^x c dx$$

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

$$F_X = \int_0^x c dx = cx$$

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

$$F_X = \int_0^x c dx = cx = \frac{x}{10}$$

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

$$F_X = \int_0^x c dx = cx = \frac{x}{10} \quad 0 < x \leq 10$$

Example 3: Loaded beam (cont.)

(b) The CDF is given by:

$$F_X = \int_0^x c dx = cx = \frac{x}{10} \quad 0 < x \leq 10$$




Figure E3.2b CDF of X.

Example 3: Loaded beam (cont.)

Example 3: Loaded beam (cont.)

(c) To compute $P(2 < X \leq 5)$, we use the CDF:

Example 3: Loaded beam (cont.)

(c) To compute $P(2 < X \leq 5)$, we use the CDF:

$$P(2 < X \leq 5) = F_X(5) - F_X(2)$$

Example 3: Loaded beam (cont.)

(c) To compute $P(2 < X \leq 5)$, we use the CDF:

$$\begin{aligned} P(2 < X \leq 5) &= F_X(5) - F_X(2) \\ &= \frac{5 - 2}{10} = \end{aligned}$$

Example 3: Loaded beam (cont.)

(c) To compute $P(2 < X \leq 5)$, we use the CDF:

$$\begin{aligned} P(2 < X \leq 5) &= F_X(5) - F_X(2) \\ &= \frac{5 - 2}{10} = 0.3 \end{aligned}$$

Example 4: Useful life of machines

The useful life T of welding machines is a random variable with an **exponential distribution**. The PDF and CDF are:

$$f_T(t) = \lambda e^{-\lambda t} \quad t \geq 0$$

Example 4: Useful life of machines

The useful life T of welding machines is a random variable with an **exponential distribution**. The PDF and CDF are:

$$\begin{aligned}f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

Example 4: Useful life of machines

The useful life T of welding machines is a random variable with an **exponential distribution**. The PDF and CDF are:

$$\begin{aligned}f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$




Figure E3.3a Exponential PDF of useful life T .

Example 4: Useful life of machines

The useful life T of welding machines is a random variable with an **exponential distribution**. The PDF and CDF are:

$$\begin{aligned}f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$




Figure E3.3a Exponential PDF of useful life T .




Figure E3.3b CDF of useful life T .

Example 4: Useful life of machines

The useful life T of welding machines is a random variable with an **exponential distribution**. The PDF and CDF are:

$$\begin{aligned}f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$




Figure E3.3a Exponential PDF of useful life T .




Figure E3.3b CDF of useful life T .

- (a) Find the mean of this distribution
- (b) Find the median
- (c) Show that the variance is $\frac{1}{\lambda^2}$

Example 4: Useful life of machines

PDF :

Example 4: Useful life of machines

$$\text{PDF : } f_T(t) = \lambda e^{-\lambda t} \quad t \geq 0$$

Example 4: Useful life of machines

$$\text{PDF : } f_T(t) = \lambda e^{-\lambda t} \quad t \geq 0$$

CDF :

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

(a) The mean is given by $\mu_T = E(T) = \int_0^{\infty} t \lambda e^{-\lambda t} dt$.

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

- (a) The mean is given by $\mu_T = E(T) = \int_0^\infty t \lambda e^{-\lambda t} dt$. We use integration by parts: $\int u dv = uv - \int v du$.

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

- (a) The mean is given by $\mu_T = E(T) = \int_0^\infty t \lambda e^{-\lambda t} dt$. We use integration by parts: $\int u dv = uv - \int v du$.

$$\mu_T = \int_0^\infty t \lambda e^{-\lambda t} dt$$

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

- (a) The mean is given by $\mu_T = E(T) = \int_0^\infty t \lambda e^{-\lambda t} dt$. We use integration by parts: $\int u dv = uv - \int v du$.

$$\begin{aligned}\mu_T &= \int_0^\infty t \lambda e^{-\lambda t} dt \\ &= \lambda \int_0^\infty \lambda e^{-\lambda t} dt\end{aligned}$$

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

- (a) The mean is given by $\mu_T = E(T) = \int_0^\infty t \lambda e^{-\lambda t} dt$. We use integration by parts: $\int u dv = uv - \int v du$.

$$\begin{aligned}\mu_T &= \int_0^\infty t \lambda e^{-\lambda t} dt \\ &= \lambda \int_0^\infty t e^{-\lambda t} dt \\ &= \lambda \left[t \left(-\frac{1}{\lambda} e^{-\lambda t} \right) \right]_0^\infty - \left[-\frac{1}{\lambda} e^{-\lambda t} \right]_0^\infty\end{aligned}$$

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

- (a) The mean is given by $\mu_T = E(T) = \int_0^\infty t \lambda e^{-\lambda t} dt$. We use integration by parts: $\int u dv = uv - \int v du$.

$$\begin{aligned}\mu_T &= \int_0^\infty t \lambda e^{-\lambda t} dt \\ &= \lambda \int_0^\infty t e^{-\lambda t} dt \\ &= \lambda \left[t \left(-\frac{1}{\lambda} e^{-\lambda t} \right) \right]_0^\infty - \left[-\frac{1}{\lambda} e^{-\lambda t} \right]_0^\infty \\ &= \lambda \left(0 + \frac{1}{\lambda} \frac{e^{-\lambda t}}{\lambda} \Big|_0^\infty \right)\end{aligned}$$

Example 4: Useful life of machines

$$\begin{aligned}\text{PDF : } f_T(t) &= \lambda e^{-\lambda t} & t \geq 0 \\ \text{CDF : } F_T(t) &= 1 - e^{-\lambda t} & t \geq 0\end{aligned}$$

- (a) The mean is given by $\mu_T = E(T) = \int_0^\infty t \lambda e^{-\lambda t} dt$. We use integration by parts: $\int u dv = uv - \int v du$.

$$\begin{aligned}\mu_T &= \int_0^\infty t \lambda e^{-\lambda t} dt \\ &= \lambda \int_0^\infty t e^{-\lambda t} dt \\ &= \lambda \left[t \left(-\frac{1}{\lambda} e^{-\lambda t} \right) \right]_0^\infty - \left[-\frac{1}{\lambda} e^{-\lambda t} \right]_0^\infty \\ &= \lambda \left(0 + \frac{1}{\lambda} \frac{e^{-\lambda t}}{\lambda} \Big|_0^\infty \right) = \frac{\lambda}{\lambda^2} = \frac{1}{\lambda}\end{aligned}$$

Recap

- Random variables

Reading

- Open Intro Statistics Section 3.4 (Random variables)
- Open Intro Statistics Section 3.5 (Continuous distributions)

Recap

- Random variables
- Probability mass/density function (PMF/PDF) and cumulative distribution function (CDF)

Reading

- Open Intro Statistics Section 3.4 (Random variables)
- Open Intro Statistics Section 3.5 (Continuous distributions)

Recap

- Random variables
- Probability mass/density function (PMF/PDF) and cumulative distribution function (CDF)
- Measures of centrality

Reading

- Open Intro Statistics Section 3.4 (Random variables)
- Open Intro Statistics Section 3.5 (Continuous distributions)

Recap

- Random variables
- Probability mass/density function (PMF/PDF) and cumulative distribution function (CDF)
- Measures of centrality
- Measures of dispersion

Reading

- Open Intro Statistics Section 3.4 (Random variables)
- Open Intro Statistics Section 3.5 (Continuous distributions)

Skewness

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

In the discrete case:

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

In the discrete case:

$$E(X - \mu_X)^3 = \sum_i (x_i - \mu_X)^3 p_X(x_i) \quad (21)$$

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

In the discrete case:

$$E(X - \mu_X)^3 = \sum_i (x_i - \mu_X)^3 p_X(x_i) \quad (21)$$

In the continuous case:

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

In the discrete case:

$$E(X - \mu_X)^3 = \sum_i (x_i - \mu_X)^3 p_X(x_i) \quad (21)$$

In the continuous case:

$$E(X - \mu_X)^3 = \int_{-\infty}^{\infty} (x - \mu_X)^3 f_X(x) dx \quad (22)$$

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

In the discrete case:

$$E(X - \mu_X)^3 = \sum_i (x_i - \mu_X)^3 p_X(x_i) \quad (21)$$

In the continuous case:

$$E(X - \mu_X)^3 = \int_{-\infty}^{\infty} (x - \mu_X)^3 f_X(x) dx \quad (22)$$

For convenience, the skewness coefficient is also used (unitless):

Skewness

The skewness or symmetry of a distribution is measured by the third central moment:

In the discrete case:

$$E(X - \mu_X)^3 = \sum_i (x_i - \mu_X)^3 p_X(x_i) \quad (21)$$

In the continuous case:

$$E(X - \mu_X)^3 = \int_{-\infty}^{\infty} (x - \mu_X)^3 f_X(x) dx \quad (22)$$

For convenience, the skewness coefficient is also used (unitless):

$$\theta = \frac{E(X - \mu_X)^3}{\sigma^3} \quad (23)$$

Skewness (cont.)

Skewness (cont.)

- Positive skewness is characterized by a long right tail (right-skewed)

Skewness (cont.)


- Positive skewness is characterized by a long right tail (right-skewed)
- Negative skewness is characterized by a long left tail (left-skewed)

Skewness (cont.)

- Positive skewness is characterized by a long right tail (right-skewed)
- Negative skewness is characterized by a long left tail (left-skewed)

Skewness (cont.)

- Positive skewness is characterized by a long right tail (right-skewed)
- Negative skewness is characterized by a long left tail (left-skewed)



Kurtosis

Kurtosis

This is the measure of peakedness in a distribution.

Kurtosis

This is the measure of peakedness in a distribution. It is the fourth central moment:

Kurtosis

This is the measure of peakedness in a distribution. It is the fourth central moment:

In the discrete case:

Kurtosis

This is the measure of peakedness in a distribution. It is the fourth central moment:

In the discrete case:

$$E(X - \mu_X)^4 = \sum_i (x_i - \mu_X)^4 p_X(x_i) \quad (24)$$

Kurtosis

This is the measure of peakedness in a distribution. It is the fourth central moment:

In the discrete case:

$$E(X - \mu_X)^4 = \sum_i (x_i - \mu_X)^4 p_X(x_i) \quad (24)$$

In the continuous case:

Kurtosis

This is the measure of peakedness in a distribution. It is the fourth central moment:

In the discrete case:

$$E(X - \mu_X)^4 = \sum_i (x_i - \mu_X)^4 p_X(x_i) \quad (24)$$



In the continuous case:

$$E(X - \mu_X)^4 = \int_{-\infty}^{\infty} (x - \mu_X)^4 f_X(x) dx \quad (25)$$


Skewness vs. kurtosis

Skewness vs. kurtosis

Skewness (S_{sk})



Kurtosis (S_{ku})



Source: Bonyar, A (2015) "Application of localization factor for the detection of tin oxidation with AFM" DOI: 10.1109/SIITME.2015.7342289

Generalized expectation

The mathematical expectation can be defined for a function g of random variable X :

$$E[g(X)] = \sum_i g(x_i) p_X(x_i) \quad \text{discrete case} \quad (26)$$

Generalized expectation

The mathematical expectation can be defined for a function g of random variable X :

$$E[g(X)] = \sum_i g(x_i) p_X(x_i) \quad \text{discrete case} \quad (26)$$

$$E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx \quad \text{continuous case} \quad (27)$$