

# CEE 260/MIE 273: Probability and Statistics in Civil Engineering

## Lecture M2c: Conditional Probability and Bayes' Theorem

**Prof. Jimi Oke**

UMassAmherst  

---

College of Engineering

September 18, 2025

# Outline

- ① Conditional probability
- ② Independent events
- ③ Multiplication rule
- ④ Theorem of total probability
- ⑤ Bayes' theorem
- ⑥ Outlook

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(E_1 \cup E_2 \cup \cdots \cup E_n) = P(E_1) + P(E_2) + \cdots + P(E_n) \quad (\text{Mutually exclusive})$$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \cdots \cup E_n) = P(E_1) + P(E_2) + \cdots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \cdots \cup E_n) = P(E_1) + P(E_2) + \cdots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)



# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$
  - Permutations of  $n$  objects:  $n! = n(n-1)(n-2) \dots (2)(1)$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$
  - Permutations of  $n$  objects:  $n! = n(n-1)(n-2) \dots (2)(1)$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$
  - Permutations of  $n$  objects:  $n! = n(n-1)(n-2) \dots (2)(1)$
  - Permutations (arrangements) of a subset of  $k$  items chosen from set of  $n$  items:  $n!/(n-k)!$



# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$
  - Permutations of  $n$  objects:  $n! = n(n-1)(n-2) \dots (2)(1)$
  - Permutations (arrangements) of a subset of  $k$  items chosen from set of  $n$  items:  $n!/(n-k)!$

# Recap from Lecture 2b: Theory of Probability

- Three axioms of probability:

$$P(E) \geq 0 \quad \text{and} \quad P(E) \leq 1$$

$$P(S) = 1$$

$$P(E_1 \cup E_2 \cup \dots \cup E_n) = P(E_1) + P(E_2) + \dots + P(E_n) \quad (\text{Mutually exclusive})$$

- Addition rule:  $P(A \cup B) = P(A) + P(B) - P(AB)$
- Addition rule for mutually exclusive events:  $P(A \cup B) = P(A) + P(B)$   
(Axiom 3)
- Counting
  - Fundamental principle of counting: number of outcomes for  $1, \dots, k$  events, each with  $n_1, \dots, n_k$  possibilities is  $n_1 \times \dots \times n_k$
  - Permutations of  $n$  objects:  $n! = n(n-1)(n-2) \dots (2)(1)$
  - Permutations (arrangements) of a subset of  $k$  items chosen from set of  $n$  items:  $n!/(n-k)!$
  - Combinations (distinct; order not important) of group of  $k$  items chosen from set of  $n$  items:  $n!/(k!(n-k)!)$

# Objectives of today's lecture

# Objectives of today's lecture

- Understand conditional probability

# Objectives of today's lecture

- Understand conditional probability
- Grasp the concept of statistical independence

# Objectives of today's lecture

- Understand conditional probability
- Grasp the concept of statistical independence
- Apply the multiplication rule and understand its relation to conditional probability

# Objectives of today's lecture

- Understand conditional probability
- Grasp the concept of statistical independence
- Apply the multiplication rule and understand its relation to conditional probability
- Understand total probability

# Objectives of today's lecture

- Understand conditional probability
- Grasp the concept of statistical independence
- Apply the multiplication rule and understand its relation to conditional probability
- Understand total probability
- Understand Bayes' Theorem and learn how to apply it to solving inverse probability problems



# Conditional probability

## Definition

The probability of an event  $E_1$  is conditional upon another event  $E_2$  if the occurrence of  $E_1$  is dependent on the occurrence of  $E_2$ :

# Conditional probability

## Definition

The probability of an event  $E_1$  is conditional upon another event  $E_2$  if the occurrence of  $E_1$  is dependent on the occurrence of  $E_2$ :

$$P(E_1|E_2) = \text{the conditional probability of } E_1 \text{ given } E_2$$

# Conditional probability

## Definition

The probability of an event  $E_1$  is conditional upon another event  $E_2$  if the occurrence of  $E_1$  is dependent on the occurrence of  $E_2$ :

$$P(E_1|E_2) = \text{the conditional probability of } E_1 \text{ given } E_2 \\ \text{(OR the probability of } E_1 \text{ conditioned on } E_2)$$

# Conditional probability

## Definition

The probability of an event  $E_1$  is conditional upon another event  $E_2$  if the occurrence of  $E_1$  is dependent on the occurrence of  $E_2$ :

$$\begin{aligned} P(E_1|E_2) &= \text{the conditional probability of } E_1 \text{ given } E_2 \\ &\quad \text{(OR the probability of } E_1 \text{ conditioned on } E_2) \\ P(E_2|E_1) &= \end{aligned}$$

# Conditional probability

## Definition

The probability of an event  $E_1$  is conditional upon another event  $E_2$  if the occurrence of  $E_1$  is dependent on the occurrence of  $E_2$ :


$$\begin{aligned} P(E_1|E_2) &= \text{the conditional probability of } E_1 \text{ given } E_2 \\ &\quad \text{(OR the probability of } E_1 \text{ conditioned on } E_2) \\ P(E_2|E_1) &= \text{the conditional probability of } E_2 \text{ given } E_1 \end{aligned}$$

# Conditional probability (cont.)

We can think of  $P(E_1|E_2)$  as the probability of realizing sample points of  $E_1$  within the subsample space of  $E_2$ .

# Conditional probability (cont.)


We can think of  $P(E_1|E_2)$  as the probability of realizing sample points of  $E_1$  within the subsample space of  $E_2$ .



$$P(S) = 1$$

# Conditional probability (cont.)


We can think of  $P(E_1|E_2)$  as the probability of realizing sample points of  $E_1$  within the subsample space of  $E_2$ .






# Conditional probability (cont.)

We can think of  $P(E_1|E_2)$  as the probability of realizing sample points of  $E_1$  within the subsample space of  $E_2$ .



# Conditional probability (cont.)

We can think of  $P(E_1|E_2)$  as the probability of realizing sample points of  $E_1$  within the subsample space of  $E_2$ .



# Conditional probability (cont.)

Thus,

# Conditional probability (cont.)


Thus,

$$P(E_1|E_2) = \frac{P(E_1 E_2)}{P(E_2)} \quad (1)$$

# Conditional probability (cont.)

Thus,

$$P(E_1|E_2) = \frac{P(E_1 E_2)}{P(E_2)} \quad (1)$$




$$P(S) = 1$$

# Conditional probability (cont.)

Thus,


$$P(E_1|E_2) = \frac{P(E_1 E_2)}{P(E_2)} \quad (1)$$



# Conditional probability (cont.)

Thus,


$$P(E_1|E_2) = \frac{P(E_1 E_2)}{P(E_2)} \quad (1)$$



# Conditional probability (cont.)

Thus,

$$P(E_1|E_2) = \frac{P(E_1 E_2)}{P(E_2)} \quad (1)$$





# Useful conditional probability relations

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

$$P(B|A) = \frac{P(BA)}{P(A)} =$$

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

$$P(B|A) = \frac{P(BA)}{P(A)} = \frac{P(AB)}{P(A)} \quad (4)$$

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

$$P(B|A) = \frac{P(BA)}{P(A)} = \frac{P(AB)}{P(A)} \quad (4)$$

$$P(\bar{A}|B) = 1 - P(A|B) \quad (5)$$

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

$$P(B|A) = \frac{P(BA)}{P(A)} = \frac{P(AB)}{P(A)} \quad (4)$$

$$P(\bar{A}|B) = 1 - P(A|B) \quad (5)$$

- A useful mnemonic: think of “|” as the division sign.

# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

$$P(B|A) = \frac{P(BA)}{P(A)} = \frac{P(AB)}{P(A)} \quad (4)$$

$$P(\bar{A}|B) = 1 - P(A|B) \quad (5)$$

- A useful mnemonic: think of “|” as the division sign.
  - This means that the event to the right is what goes in the denominator.



# Useful conditional probability relations

$$P(A|S) = P(A) \quad (2)$$

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (3)$$

$$P(B|A) = \frac{P(BA)}{P(A)} = \frac{P(AB)}{P(A)} \quad (4)$$

$$P(\bar{A}|B) = 1 - P(A|B) \quad (5)$$

- A useful mnemonic: think of “|” as the division sign.
  - This means that the event to the right is what goes in the denominator.
  - The numerator is simply the intersection of both events

# Example 1: Highway routes

## Example 1: Highway routes

There are two routes from City A to City B. One or both routes may be closed due to severe weather.

## Example 1: Highway routes


There are two routes from City A to City B. One or both routes may be closed due to severe weather.

We denote  $E_1$  as the event Route 1 is open, and  $E_2$  as the event Route 2 is open.

# Example 1: Highway routes

There are two routes from City A to City B. One or both routes may be closed due to severe weather.


We denote  $E_1$  as the event Route 1 is open, and  $E_2$  as the event Route 2 is open.



# Example 1: Highway routes

There are two routes from City A to City B. One or both routes may be closed due to severe weather.

We denote  $E_1$  as the event Route 1 is open, and  $E_2$  as the event Route 2 is open.




During severe weather, the probabilities the routes will be open are:

## Example 1: Highway routes

There are two routes from City A to City B. One or both routes may be closed due to severe weather.

We denote  $E_1$  as the event Route 1 is open, and  $E_2$  as the event Route 2 is open.




During severe weather, the probabilities the routes will be open are:

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

# Example 1: Highway routes

There are two routes from City A to City B. One or both routes may be closed due to severe weather.

We denote  $E_1$  as the event Route 1 is open, and  $E_2$  as the event Route 2 is open.



During severe weather, the probabilities the routes will be open are:

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (a) What is probability that Route 1 is open during a storm given that Route 2 is also open?
- (b) What is probability that Route 2 is open during a storm conditioned on the event that Route 1 is also open?
- (c) What is probability that Route 1 is closed during a storm given that Route 2 is also closed?



# Example 1: Highway routes (cont.)

## Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (a) The probability that Route 1 is open ( $E_1$ ) during a storm given that Route 2 is also open ( $E_2$ ) is denoted as:

## Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (a) The probability that Route 1 is open ( $E_1$ ) during a storm given that Route 2 is also open ( $E_2$ ) is denoted as:

$$P(E_1|E_2) = \frac{P(E_1 E_2)}{P(E_2)}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (a) The probability that Route 1 is open ( $E_1$ ) during a storm given that Route 2 is also open ( $E_2$ ) is denoted as:

$$\begin{aligned} P(E_1|E_2) &= \frac{P(E_1 E_2)}{P(E_2)} \\ &= \frac{0.40}{0.50} = \end{aligned}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (a) The probability that Route 1 is open ( $E_1$ ) during a storm given that Route 2 is also open ( $E_2$ ) is denoted as:

$$\begin{aligned} P(E_1|E_2) &= \frac{P(E_1 E_2)}{P(E_2)} \\ &= \frac{0.40}{0.50} = \boxed{0.80} \end{aligned}$$

# Example 1: Highway routes (cont.)

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$



# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (b) The probability that Route 2 is open ( $E_2$ ) during a storm conditioned on the event that Route 1 is also open ( $E_1$ ) is denoted as:

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (b) The probability that Route 2 is open ( $E_2$ ) during a storm conditioned on the event that Route 1 is also open ( $E_1$ ) is denoted as:

$$P(E_2|E_1) = \frac{P(E_2 E_1)}{P(E_1)}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (b) The probability that Route 2 is open ( $E_2$ ) during a storm conditioned on the event that Route 1 is also open ( $E_1$ ) is denoted as:

$$\begin{aligned} P(E_2|E_1) &= \frac{P(E_2 E_1)}{P(E_1)} \\ &= \frac{0.40}{0.75} = \end{aligned}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (b) The probability that Route 2 is open ( $E_2$ ) during a storm conditioned on the event that Route 1 is also open ( $E_1$ ) is denoted as:

$$\begin{aligned} P(E_2|E_1) &= \frac{P(E_2 E_1)}{P(E_1)} \\ &= \frac{0.40}{0.75} = \boxed{0.53} \end{aligned}$$

## Example 1: Highway routes (cont.)

## Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

## Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\bar{E}_1$ ) during a storm given that Route 2 is also closed ( $\bar{E}_2$ ) is denoted as:

## Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$



# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$P(\overline{E}_1 \overline{E}_2) = 1 - P(\overline{\overline{E}_1 \overline{E}_2})$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \end{aligned}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \end{aligned}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \end{aligned}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \quad \text{addition rule} \\ &= 1 - (0.75 + 0.50 - 0.40) \\ &= 1 - 0.85 = \end{aligned}$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \quad \text{addition rule} \\ &= 1 - (0.75 + 0.50 - 0.40) \\ &= 1 - 0.85 = 0.15 \end{aligned}$$

Thus, we obtain

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \quad \text{addition rule} \\ &= 1 - (0.75 + 0.50 - 0.40) \\ &= 1 - 0.85 = 0.15 \end{aligned}$$

Thus, we obtain

$$P(\overline{E}_1 | \overline{E}_2) =$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\overline{E}_1$ ) during a storm given that Route 2 is also closed ( $\overline{E}_2$ ) is denoted as:

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)}$$

$$\begin{aligned} P(\overline{E}_1 \overline{E}_2) &= 1 - P(\overline{\overline{E}_1 \overline{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \quad \text{addition rule} \\ &= 1 - (0.75 + 0.50 - 0.40) \\ &= 1 - 0.85 = 0.15 \end{aligned}$$

Thus, we obtain

$$P(\overline{E}_1 | \overline{E}_2) = \frac{P(\overline{E}_1 \overline{E}_2)}{P(\overline{E}_2)} =$$



# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\bar{E}_1$ ) during a storm given that Route 2 is also closed ( $\bar{E}_2$ ) is denoted as:

$$P(\bar{E}_1 | \bar{E}_2) = \frac{P(\bar{E}_1 \bar{E}_2)}{P(\bar{E}_2)}$$

$$\begin{aligned} P(\bar{E}_1 \bar{E}_2) &= 1 - P(\overline{\bar{E}_1 \bar{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \quad \text{addition rule} \\ &= 1 - (0.75 + 0.50 - 0.40) \\ &= 1 - 0.85 = 0.15 \end{aligned}$$

Thus, we obtain

$$P(\bar{E}_1 | \bar{E}_2) = \frac{P(\bar{E}_1 \bar{E}_2)}{P(\bar{E}_2)} = \frac{0.15}{0.50} =$$

# Example 1: Highway routes (cont.)

$$P(E_1) = 0.75 \quad P(E_2) = 0.50 \quad P(E_1 E_2) = 0.40$$

- (c) The probability that Route 1 is closed ( $\bar{E}_1$ ) during a storm given that Route 2 is also closed ( $\bar{E}_2$ ) is denoted as:

$$P(\bar{E}_1 | \bar{E}_2) = \frac{P(\bar{E}_1 \bar{E}_2)}{P(\bar{E}_2)}$$

$$\begin{aligned} P(\bar{E}_1 \bar{E}_2) &= 1 - P(\overline{\bar{E}_1 \bar{E}_2}) \quad \text{complementary events} \\ &= 1 - P(E_1 \cup E_2) \quad \text{De Morgan's Rule} \\ &= 1 - [P(E_1) + P(E_2) - P(E_1 E_2)] \quad \text{addition rule} \\ &= 1 - (0.75 + 0.50 - 0.40) \\ &= 1 - 0.85 = 0.15 \end{aligned}$$

Thus, we obtain

$$P(\bar{E}_1 | \bar{E}_2) = \frac{P(\bar{E}_1 \bar{E}_2)}{P(\bar{E}_2)} = \frac{0.15}{0.50} = \boxed{0.30}$$

# The addition rule

# The addition rule

Conditioning equally applies to probabilities of events on the same subsample space.

# The addition rule

Conditioning equally applies to probabilities of events on the same subsample space.

$$P(A \cup B) = P(A) + P(B) - P(AB) \quad (6)$$

# The addition rule

Conditioning equally applies to probabilities of events on the same subsample space.

$$P(A \cup B) = P(A) + P(B) - P(AB) \quad (6)$$

$$P(A \cup B|E) = P(A|E) + P(B|E) - P(AB|E) \quad (7)$$

# Statistically independent events

# Statistically independent events

## Statistical independence



# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession.

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession. The result of the first roll does not affect the result of the other.

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession. The result of the first roll does not affect the result of the other.
- The event that it will be cloudy in Amherst tomorrow and the event that the number of births in California tomorrow will increase compared to that of today.

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession. The result of the first roll does not affect the result of the other.
- The event that it will be cloudy in Amherst tomorrow and the event that the number of births in California tomorrow will increase compared to that of today. (Knowledge of one event cannot improve prediction of the other.)



# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession. The result of the first roll does not affect the result of the other.
- The event that it will be cloudy in Amherst tomorrow and the event that the number of births in California tomorrow will increase compared to that of today. (Knowledge of one event cannot improve prediction of the other.)
- The ages of a random sample of Pioneer Valley residents

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession. The result of the first roll does not affect the result of the other.
- The event that it will be cloudy in Amherst tomorrow and the event that the number of births in California tomorrow will increase compared to that of today. (Knowledge of one event cannot improve prediction of the other.)
- The ages of a random sample of Pioneer Valley residents
- Generally, all elements of a random sample are assumed independent

# Statistically independent events

## Statistical independence

Two events are statistically independent if the occurrence of one does not depend on the occurrence or non-occurrence of the other:

$$P(E_1|E_2) = P(E_1) \quad (8)$$

$$P(E_2|E_1) = P(E_2) \quad (9)$$

## Examples of independent events

- The outcomes of a die rolled twice in succession. The result of the first roll does not affect the result of the other.
- The event that it will be cloudy in Amherst tomorrow and the event that the number of births in California tomorrow will increase compared to that of today. (Knowledge of one event cannot improve prediction of the other.)
- The ages of a random sample of Pioneer Valley residents
- Generally, all elements of a random sample are assumed independent

## Example 2: Rolling two dice

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

$$P(X = 1) = \frac{1}{6}$$

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

$$P(X = 1) = \frac{1}{6}$$

(b) What is the probability that both  $X$  and  $Y$  are 1?



## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

$$P(X = 1) = \frac{1}{6}$$

(b) What is the probability that both  $X$  and  $Y$  are 1?

The outcome of  $X$  has no bearing on the outcome of  $Y$ . Thus:

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

$$P(X = 1) = \frac{1}{6}$$

(b) What is the probability that both  $X$  and  $Y$  are 1?

The outcome of  $X$  has no bearing on the outcome of  $Y$ . Thus:

$$P(X = 1 \cap Y = 1) = P(X = 1) \times P(Y = 1)$$

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

$$P(X = 1) = \frac{1}{6}$$

(b) What is the probability that both  $X$  and  $Y$  are 1?

The outcome of  $X$  has no bearing on the outcome of  $Y$ . Thus:

$$\begin{aligned} P(X = 1 \cap Y = 1) &= P(X = 1) \times P(Y = 1) \\ &= \frac{1}{6} \times \frac{1}{6} = \end{aligned}$$

## Example 2: Rolling two dice

Let  $X$  and  $Y$  represent the outcomes of rolling two dice.

(a) What is the probability that the first die  $X$  is 1?

$$P(X = 1) = \frac{1}{6}$$

(b) What is the probability that both  $X$  and  $Y$  are 1?

The outcome of  $X$  has no bearing on the outcome of  $Y$ . Thus:

$$\begin{aligned} P(X = 1 \cap Y = 1) &= P(X = 1) \times P(Y = 1) \\ &= \frac{1}{6} \times \frac{1}{6} = \frac{1}{36} \end{aligned}$$

## Example 2: Rolling two dice (cont.)

## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$


## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$P(Y = 1|X = 1) = \frac{P(Y = 1 \cap X = 1)}{P(X = 1)}$$

## Example 2: Rolling two dice (cont.)


(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$\begin{aligned} P(Y = 1|X = 1) &= \frac{P(Y = 1 \cap X = 1)}{P(X = 1)} \\ &= \frac{P(Y = 1) \times P(X = 1)}{P(X = 1)} \end{aligned}$$





## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$\begin{aligned} P(Y = 1|X = 1) &= \frac{P(Y = 1 \cap X = 1)}{P(X = 1)} \\ &= \frac{P(Y = 1) \times P(X = 1)}{P(X = 1)} \end{aligned}$$


## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$\begin{aligned} P(Y = 1|X = 1) &= \frac{P(Y = 1 \cap X = 1)}{P(X = 1)} \\ &= \frac{P(Y = 1) \times P(X = 1)}{P(X = 1)} \end{aligned}$$


## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$\begin{aligned}P(Y = 1|X = 1) &= \frac{P(Y = 1 \cap X = 1)}{P(X = 1)} \\&= \frac{P(Y = 1) \times P(X = 1)}{P(X = 1)} \\&= P(Y = 1)\end{aligned}$$

## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$\begin{aligned}P(Y = 1|X = 1) &= \frac{P(Y = 1 \cap X = 1)}{P(X = 1)} \\&= \frac{P(Y = 1) \times P(X = 1)}{P(X = 1)} \\&= P(Y = 1)\end{aligned}$$

(d) Why is  $P(Y = 1|X = 1) = P(Y = 1)$ ?

## Example 2: Rolling two dice (cont.)

(c) Use the conditional probability formula to find  $P(Y = 1|X = 1)$

$$\begin{aligned}P(Y = 1|X = 1) &= \frac{P(Y = 1 \cap X = 1)}{P(X = 1)} \\&= \frac{P(Y = 1) \times P(X = 1)}{P(X = 1)} \\&= P(Y = 1)\end{aligned}$$

(d) Why is  $P(Y = 1|X = 1) = P(Y = 1)$ ?

$X$  and  $Y$  are independent events.

# The multiplication rule

# The multiplication rule

The probability of the intersection of two events is

# The multiplication rule

The probability of the intersection of two events is

$$P(E_1 E_2) = P(E_1 | E_2) P(E_2) \quad (10)$$



# The multiplication rule

The probability of the intersection of two events is

$$P(E_1 E_2) = P(E_1 | E_2) P(E_2) \quad (10)$$

Equivalently:

# The multiplication rule

The probability of the intersection of two events is

$$P(E_1 E_2) = P(E_1 | E_2) P(E_2) \quad (10)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (11)$$

# The multiplication rule

The probability of the intersection of two events is

$$P(E_1 E_2) = P(E_1 | E_2) P(E_2) \quad (10)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (11)$$

## Statistically independent events

IF  $A$  and  $B$  are statistically independent, the multiplication rule becomes:

# The multiplication rule

The probability of the intersection of two events is

$$P(E_1 E_2) = P(E_1 | E_2) P(E_2) \quad (10)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (11)$$

## Statistically independent events

IF  $A$  and  $B$  are statistically independent, the multiplication rule becomes:

$$P(AB) = P(A)P(B) \quad (12)$$

# The multiplication rule

The probability of the intersection of two events is

$$P(E_1 E_2) = P(E_1 | E_2) P(E_2) \quad (10)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (11)$$

## Statistically independent events

IF  $A$  and  $B$  are statistically independent, the multiplication rule becomes:

$$P(AB) = P(A)P(B) \quad (12)$$

That is, the **joint probability** of two statistically independent events is the **product** of their individual probabilities.

# The multiplication rule: three events

The probability of the joint occurrence of three events is:

# The multiplication rule: three events

The probability of the joint occurrence of three events is:

$$\begin{aligned}P(ABC) &= P(A|BC)P(BC) \\ &= P(A|BC)P(B|C)P(C)\end{aligned}\tag{13}$$

# The multiplication rule: three events

The probability of the joint occurrence of three events is:

$$\begin{aligned}P(ABC) &= P(A|BC)P(BC) \\ &= P(A|BC)P(B|C)P(C)\end{aligned}\tag{13}$$

## Statistically independent events

If the three events are statistically independent, then:



# The multiplication rule: three events

The probability of the joint occurrence of three events is:

$$\begin{aligned}P(ABC) &= P(A|BC)P(BC) \\ &= P(A|BC)P(B|C)P(C)\end{aligned}\tag{13}$$

## Statistically independent events

If the three events are statistically independent, then:

$$P(ABC) =$$

# The multiplication rule: three events

The probability of the joint occurrence of three events is:

$$\begin{aligned}P(ABC) &= P(A|BC)P(BC) \\ &= P(A|BC)P(B|C)P(C)\end{aligned}\tag{13}$$

## Statistically independent events

If the three events are statistically independent, then:

$$P(ABC) = P(A)P(B)P(C)\tag{14}$$

# The multiplication rule: conditional probabilities

Equally applies to probabilities of events conditioned on the same subsample space.

# The multiplication rule: conditional probabilities

Equally applies to probabilities of events conditioned on the same subsample space.

$$P(AB|E) = P(A|B|E)P(B|E) \quad (15)$$

# The multiplication rule: conditional probabilities

Equally applies to probabilities of events conditioned on the same subsample space.

$$P(AB|E) = P(A|B|E)P(B|E) \quad (15)$$

## Statistically independent events

If  $A$  and  $B$  are statistically independent, the multiplication rule for two events conditioned on the same space becomes:

# The multiplication rule: conditional probabilities

Equally applies to probabilities of events conditioned on the same subsample space.

$$P(AB|E) = P(A|B|E)P(B|E) \quad (15)$$

## Statistically independent events

If  $A$  and  $B$  are statistically independent, the multiplication rule for two events conditioned on the same space becomes:

$$P(AB|E) = P(A|E)P(B|E) \quad (16)$$

## Example 3: Airline industry strikes

## Example 3: Airline industry strikes

The airline industry in a certain country is subject to labor strikes by the pilots (event  $A$ ), mechanics (event  $B$ ) or flight attendants (event  $C$ ).



## Example 3: Airline industry strikes

The airline industry in a certain country is subject to labor strikes by the pilots (event  $A$ ), mechanics (event  $B$ ) or flight attendants (event  $C$ ).

The probability of strikes by each of the individual groups in the next 3 years is given by:

## Example 3: Airline industry strikes

The airline industry in a certain country is subject to labor strikes by the pilots (event  $A$ ), mechanics (event  $B$ ) or flight attendants (event  $C$ ).

The probability of strikes by each of the individual groups in the next 3 years is given by:

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

## Example 3: Airline industry strikes

The airline industry in a certain country is subject to labor strikes by the pilots (event  $A$ ), mechanics (event  $B$ ) or flight attendants (event  $C$ ).

The probability of strikes by each of the individual groups in the next 3 years is given by:

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

Assuming that events  $A$ ,  $B$  and  $C$  are statistically independent, find

- (a) the probability that all groups will strike in the next 3 years
- (b) the probability of a labor strike in the airline industry in the next 3 years.

## Example 3: Airline industry strikes (cont.)

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

(a) The probability that all 3 groups will strike in the next 3 years is given by:

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

(a) The probability that all 3 groups will strike in the next 3 years is given by:

$$P(ABC) = P(A)P(B)P(C)$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

(a) The probability that all 3 groups will strike in the next 3 years is given by:

$$P(ABC) = P(A)P(B)P(C) \quad \text{statistical independence}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

(a) The probability that all 3 groups will strike in the next 3 years is given by:

$$\begin{aligned} P(ABC) &= P(A)P(B)P(C) \quad \text{statistical independence} \\ &= 0.03(0.05)(0.06) \end{aligned}$$



## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

(a) The probability that all 3 groups will strike in the next 3 years is given by:

$$\begin{aligned} P(ABC) &= P(A)P(B)P(C) \quad \text{statistical independence} \\ &= 0.03(0.05)(0.06) \\ &= 0.00009 = \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

(a) The probability that all 3 groups will strike in the next 3 years is given by:

$$\begin{aligned} P(ABC) &= P(A)P(B)P(C) \quad \text{statistical independence} \\ &= 0.03(0.05)(0.06) \\ &= 0.00009 = \boxed{9.0 \times 10^{-5}} \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:



## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$P(A \cup B \cup C) = 1 - P(\overline{A \cup B \cup C})$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$P(A \cup B \cup C) = 1 - P(\overline{A \cup B \cup C}) \quad \text{complement rule}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) \quad \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) \quad \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) \quad \text{de Morgan's rule} \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) \quad \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) \quad \text{de Morgan's rule} \\ &= 1 - P(\overline{A})P(\overline{B})P(\overline{C}) \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) && \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) && \text{de Morgan's rule} \\ &= 1 - P(\overline{A})P(\overline{B})P(\overline{C}) && \text{statistical independence} \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

**(Step 1)** Formulate the desired probability:

$$P(A \cup B \cup C)$$

**(Step 2)** Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) && \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) && \text{de Morgan's rule} \\ &= 1 - P(\overline{A})P(\overline{B})P(\overline{C}) && \text{statistical independence} \\ &= 1 - (1 - 0.03)(1 - 0.05)(1 - 0.06) \end{aligned}$$

## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

**(Step 1)** Formulate the desired probability:

$$P(A \cup B \cup C)$$

**(Step 2)** Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) && \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) && \text{de Morgan's rule} \\ &= 1 - P(\overline{A})P(\overline{B})P(\overline{C}) && \text{statistical independence} \\ &= 1 - (1 - 0.03)(1 - 0.05)(1 - 0.06) \\ &= 1 - (0.97)(0.95)(0.94) \end{aligned}$$



## Example 3: Airline industry strikes (cont.)

$$P(A) = 0.03 \quad P(B) = 0.05 \quad P(C) = 0.06$$

**(b)** The probability of a labor strike is the probability that any combination of the groups will strike.

(Step 1) Formulate the desired probability:

$$P(A \cup B \cup C)$$

(Step 2) Use the complement rule, de Morgan's rule and multiplication rule for statistically independent events:

$$\begin{aligned} P(A \cup B \cup C) &= 1 - P(\overline{A \cup B \cup C}) && \text{complement rule} \\ &= 1 - P(\overline{A} \overline{B} \overline{C}) && \text{de Morgan's rule} \\ &= 1 - P(\overline{A})P(\overline{B})P(\overline{C}) && \text{statistical independence} \\ &= 1 - (1 - 0.03)(1 - 0.05)(1 - 0.06) \\ &= 1 - (0.97)(0.95)(0.94) \\ &= \boxed{0.134} \end{aligned}$$

# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$


# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$




# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) =$$


# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) = P(AE_1) +$$


# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) = P(AE_1) + P(AE_2)$$




# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) = P(AE_1) + P(AE_2) + \dots$$


# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) = P(AE_1) + P(AE_2) + P(AE_3) + \dots + P(AE_n)$$


# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) = P(AE_1) + P(AE_2) + P(AE_3) + \dots + P(AE_n)$$

Note that:


# Total probability

Useful in situations where the probability of an event cannot be directly determined but its conditional probabilities are known.

## Theorem of total probability

The probability of an event  $A$  conditioned on the mutually exclusive and collectively exhaustive events  $E_1, E_2, \dots, E_n$  is given by

$$P(A) = P(A|E_1)P(E_1) + P(A|E_2)P(E_2) + \dots + P(A|E_n)P(E_n) \quad (17)$$



$$P(A) = P(AE_1) + P(AE_2) + P(AE_3) + \dots + P(AE_n)$$

Note that:

$$P(AE_1) = P(A|E_1)P(E_1), \text{ etc.}$$

## Example 4: Flooding and snow accumulation

## Example 4: Flooding and snow accumulation

The flooding of a river in the spring season (event  $F$ ) will depend on the accumulation of snow in the mountain during the past winter.

## Example 4: Flooding and snow accumulation

The flooding of a river in the spring season (event  $F$ ) will depend on the accumulation of snow in the mountain during the past winter. The accumulation of snow may be heavy ( $H$ ), normal ( $N$ ) or light ( $L$ ).

## Example 4: Flooding and snow accumulation

The flooding of a river in the spring season (event  $F$ ) will depend on the accumulation of snow in the mountain during the past winter. The accumulation of snow may be heavy ( $H$ ), normal ( $N$ ) or light ( $L$ ).

Given:



## Example 4: Flooding and snow accumulation

The flooding of a river in the spring season (event  $F$ ) will depend on the accumulation of snow in the mountain during the past winter. The accumulation of snow may be heavy ( $H$ ), normal ( $N$ ) or light ( $L$ ).

Given:

$$P(F|H) = 0.90 \quad P(F|N) = 0.40 \quad P(F|L) = 0.10$$

and in the winter

## Example 4: Flooding and snow accumulation

The flooding of a river in the spring season (event  $F$ ) will depend on the accumulation of snow in the mountain during the past winter. The accumulation of snow may be heavy ( $H$ ), normal ( $N$ ) or light ( $L$ ).

Given:

$$P(F|H) = 0.90 \quad P(F|N) = 0.40 \quad P(F|L) = 0.10$$

and in the winter

$$P(H) = 0.20 \quad P(N) = 0.50 \quad P(L) = 0.30$$

## Example 4: Flooding and snow accumulation

The flooding of a river in the spring season (event  $F$ ) will depend on the accumulation of snow in the mountain during the past winter. The accumulation of snow may be heavy ( $H$ ), normal ( $N$ ) or light ( $L$ ).

Given:

$$P(F|H) = 0.90 \quad P(F|N) = 0.40 \quad P(F|L) = 0.10$$

and in the winter

$$P(H) = 0.20 \quad P(N) = 0.50 \quad P(L) = 0.30$$

Find the probability of flooding in the river during the spring season  $P(F)$ .

## Example 4: Flooding and snow accumulation (cont.)

## Example 4: Flooding and snow accumulation (cont.)

We use the theorem of total probability:

## Example 4: Flooding and snow accumulation (cont.)

We use the theorem of total probability:

$$P(F) = P(F|H)P(H) + P(F|N)P(N) + P(F|L)P(L)$$

## Example 4: Flooding and snow accumulation (cont.)

We use the theorem of total probability:

$$\begin{aligned} P(F) &= P(F|H)P(H) + P(F|N)P(N) + P(F|L)P(L) \\ &= 0.90(0.20) + 0.40(0.50) + 0.10(0.30) \end{aligned}$$

## Example 4: Flooding and snow accumulation (cont.)

We use the theorem of total probability:

$$\begin{aligned} P(F) &= P(F|H)P(H) + P(F|N)P(N) + P(F|L)P(L) \\ &= 0.90(0.20) + 0.40(0.50) + 0.10(0.30) \\ &= 0.41 \end{aligned}$$



# Derivation of Bayes' theorem

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (19)$$

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (19)$$

We combine both equations to obtain:

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (19)$$

We combine both equations to obtain:

$$P(A|B)P(B) = P(B|A)P(A) \quad (20)$$

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (19)$$

We combine both equations to obtain:

$$P(A|B)P(B) = P(B|A)P(A) \quad (20)$$

Then, we obtain the **inverse probability** of the conditioning event:



# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (19)$$

We combine both equations to obtain:

$$P(A|B)P(B) = P(B|A)P(A) \quad (20)$$

Then, we obtain the **inverse probability** of the conditioning event:

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)} \quad (21)$$

# Derivation of Bayes' theorem

Recall from the multiplication rule that:

$$P(AB) = P(A|B)P(B) \quad (18)$$

Equivalently:

$$P(AB) = P(B|A)P(A) \quad (19)$$

We combine both equations to obtain:

$$P(A|B)P(B) = P(B|A)P(A) \quad (20)$$

Then, we obtain the **inverse probability** of the conditioning event:

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)} \quad (21)$$

This is in essence Bayes' Theorem

# Bayes' theorem

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} =$$

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$



Rev. Thomas Bayes  
(1701-61)

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$



Rev. Thomas Bayes  
(1701-61)



# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$



Rev. Thomas Bayes  
(1701-61)

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

If the event  $E$  can be conditioned on only two events  $A$  and  $B$ , then:

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

If the event  $E$  can be conditioned on only two events  $A$  and  $B$ , then:

$$P(A|E) =$$

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

If the event  $E$  can be conditioned on only two events  $A$  and  $B$ , then:

$$P(A|E) = \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \quad (23)$$

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

If the event  $E$  can be conditioned on only two events  $A$  and  $B$ , then:

$$P(A|E) = \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \quad (23)$$

$$P(B|E) =$$

# Bayes' theorem

Bayes' Theorem allows for the computation of an inverse probability, e.g. given  $P(E|A)$ , can we find  $P(A|E)$ ?

$$P(A_i|E) = \frac{P(E|A_i)P(A_i)}{\sum_{j=1}^n P(E|A_j)P(A_j)} = \frac{P(E|A_i)P(A_i)}{P(E)} \quad (22)$$

- **posterior probability:**  $P(A_i|E)$
- **likelihood:**  $P(E|A_i)$
- **prior:**  $P(A_i)$
- **evidence (total probability):**  $P(E)$



Rev. Thomas Bayes  
(1701-61)

If the event  $E$  can be conditioned on only two events  $A$  and  $B$ , then:

$$P(A|E) = \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \quad (23)$$

$$P(B|E) = \frac{P(E|B)P(B)}{P(E|A)P(A) + P(E|B)P(B)} \quad (24)$$



## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies.

## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies. Company *a* delivers 600 truckloads a day while Company *b* delivers 400 truckloads a day.

## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies. Company *a* delivers 600 truckloads a day while Company *b* delivers 400 truckloads a day. From prior experience, 3% of Company *a*'s material is expected to be substandard while 1% of Company *b*'s material is expected to be substandard.

## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies. Company *a* delivers 600 truckloads a day while Company *b* delivers 400 truckloads a day. From prior experience, 3% of Company *a*'s material is expected to be substandard while 1% of Company *b*'s material is expected to be substandard.

We define:

## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies. Company *a* delivers 600 truckloads a day while Company *b* delivers 400 truckloads a day. From prior experience, 3% of Company *a*'s material is expected to be substandard while 1% of Company *b*'s material is expected to be substandard.

We define:

$A$  = aggregates supplied by Company *a*

## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies. Company *a* delivers 600 truckloads a day while Company *b* delivers 400 truckloads a day. From prior experience, 3% of Company *a*'s material is expected to be substandard while 1% of Company *b*'s material is expected to be substandard.

We define:

$A$  = aggregates supplied by Company *a*

$B$  = aggregates supplied by Company *b*

## Example 5: Construction supplies

Aggregates for the construction of a reinforced concrete building are supplied by two companies. Company *a* delivers 600 truckloads a day while Company *b* delivers 400 truckloads a day. From prior experience, 3% of Company *a*'s material is expected to be substandard while 1% of Company *b*'s material is expected to be substandard.

We define:

$A$  = aggregates supplied by Company *a*

$B$  = aggregates supplied by Company *b*

$E$  = aggregates are substandard

## Example 5: Construction supplies (cont.)




## Example 5: Construction supplies (cont.)

- (a) Draw a Venn diagram and convince yourself that  
 $P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$


## Example 5: Construction supplies (cont.)

- (a) Draw a Venn diagram and convince yourself that  
 $P(A) = 0.60$ ,  $P(B) = 0.40$ ,  $P(E|A) = 0.03$ ,  $P(E|B) = 0.01$




## Example 5: Construction supplies (cont.)

- (a) Draw a Venn diagram and convince yourself that  
 $P(A) = 0.60$ ,  $P(B) = 0.40$ ,  $P(E|A) = 0.03$ ,  $P(E|B) = 0.01$



## Example 5: Construction supplies (cont.)

- (a) Draw a Venn diagram and convince yourself that  
 $P(A) = 0.60$ ,  $P(B) = 0.40$ ,  $P(E|A) = 0.03$ ,  $P(E|B) = 0.01$



## Example 5: Construction supplies (cont.)

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

## Example 5: Construction supplies (cont.)


$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(a)  $P(E|A) = \frac{P(EA)}{P(A)}$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(a)  $P(E|A) = \frac{P(EA)}{P(A)}$






## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$


(a)  $P(E|A) = \frac{P(EA)}{P(A)}$



## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$


(a)  $P(E|A) = \frac{P(EA)}{P(A)}$



## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(a)  $P(E|A) = \frac{P(EA)}{P(A)}$



## Example 5: Construction supplies (cont.)

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

## Example 5: Construction supplies (cont.)


$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(a)  $P(E|B) = \frac{P(EB)}{P(B)}$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$


(a)  $P(E|B) = \frac{P(EB)}{P(B)}$



## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(a)  $P(E|B) = \frac{P(EB)}{P(B)}$






## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$


(a)  $P(E|B) = \frac{P(EB)}{P(B)}$



## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(a)  $P(E|B) = \frac{P(EB)}{P(B)}$



## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$P(E) = P(E|A)P(A) +$$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$P(E) = P(E|A)P(A) + P(E|B)P(B)$$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$\begin{aligned} P(E) &= P(E|A)P(A) + P(E|B)P(B) \\ &= (0.03)(0.6) + \end{aligned}$$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$\begin{aligned} P(E) &= P(E|A)P(A) + P(E|B)P(B) \\ &= (0.03)(0.6) + (0.01)(0.4) \end{aligned}$$



## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$\begin{aligned} P(E) &= P(E|A)P(A) + P(E|B)P(B) \\ &= (0.03)(0.6) + (0.01)(0.4) \\ &= 0.018 + \end{aligned}$$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$\begin{aligned} P(E) &= P(E|A)P(A) + P(E|B)P(B) \\ &= (0.03)(0.6) + (0.01)(0.4) \\ &= 0.018 + 0.004 \end{aligned}$$

## Example 5: Construction supplies (cont.)

$$P(A) = 0.60, P(B) = 0.40, P(E|A) = 0.03, P(E|B) = 0.01$$

(b) Show that  $P(E) = 0.022$

We use the Theorem of Total Probability:

$$\begin{aligned} P(E) &= P(E|A)P(A) + P(E|B)P(B) \\ &= (0.03)(0.6) + (0.01)(0.4) \\ &= 0.018 + 0.004 \\ &= \boxed{0.022} \end{aligned}$$

## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

$$P(A|E) = \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)}$$

## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

$$\begin{aligned}
 P(A|E) &= \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \\
 &= \frac{P(E|A)P(A)}{P(E)} \quad \text{Denominator: total probability}
 \end{aligned}$$

## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

$$\begin{aligned} P(A|E) &= \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \\ &= \frac{P(E|A)P(A)}{P(E)} \quad \text{Denominator: total probability} \\ &= \frac{0.03 \times 0.60}{0.022} \end{aligned}$$

## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

$$\begin{aligned} P(A|E) &= \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \\ &= \frac{P(E|A)P(A)}{P(E)} \quad \text{Denominator: total probability} \\ &= \frac{0.03 \times 0.60}{0.022} \equiv \frac{\text{likelihood} \times \text{prior}}{\text{evidence}} \end{aligned}$$



## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

$$\begin{aligned} P(A|E) &= \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \\ &= \frac{P(E|A)P(A)}{P(E)} \quad \text{Denominator: total probability} \\ &= \frac{0.03 \times 0.60}{0.022} \equiv \frac{\text{likelihood} \times \text{prior}}{\text{evidence}} \\ &= 0.818 \approx \boxed{0.82} \end{aligned}$$

## Example 5: Construction supplies (cont.)

(c) Show that the probability  $P(A|E) = 0.82$  and discuss its significance

Here, we use Bayes' Theorem:

$$\begin{aligned}
 P(A|E) &= \frac{P(E|A)P(A)}{P(E|A)P(A) + P(E|B)P(B)} \\
 &= \frac{P(E|A)P(A)}{P(E)} \quad \text{Denominator: total probability} \\
 &= \frac{0.03 \times 0.60}{0.022} \equiv \frac{\text{likelihood} \times \text{prior}}{\text{evidence}} \\
 &= 0.818 \approx \boxed{0.82}
 \end{aligned}$$

$P(A|E)$  is the posterior probability of  $A$  having observed  $E$ . In other words, having prior knowledge of  $A$  (i.e.  $P(A)$  and the likelihood of  $E$  given  $A$ , we can update our knowledge of  $A$  (posterior probability) based on these observations.

# Recap

- **Conditional probability:**

# Recap

- **Conditional probability:**

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

$$P(AB) = P(A)P(B) \quad (26)$$



# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

$$P(AB) = P(A)P(B) \quad (26)$$

Generally, the joint probability (intersection) of any number of independent events is the product of their individual probabilities:

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

$$P(AB) = P(A)P(B) \quad (26)$$

Generally, the joint probability (intersection) of any number of independent events is the product of their individual probabilities:

$$P(E_1 \cap E_2 \cap \cdots \cap E_n) = P(E_1)P(E_2) \cdots P(E_n) \quad (27)$$

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

$$P(AB) = P(A)P(B) \quad (26)$$

Generally, the joint probability (intersection) of any number of independent events is the product of their individual probabilities:

$$P(E_1 \cap E_2 \cap \cdots \cap E_n) = P(E_1)P(E_2) \cdots P(E_n) \quad (27)$$

- **Multiplication rule:**

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

$$P(AB) = P(A)P(B) \quad (26)$$

Generally, the joint probability (intersection) of any number of independent events is the product of their individual probabilities:

$$P(E_1 \cap E_2 \cap \cdots \cap E_n) = P(E_1)P(E_2) \cdots P(E_n) \quad (27)$$

- **Multiplication rule:**

# Recap

- **Conditional probability:**

$$P(A|B) = \frac{P(AB)}{P(B)} \quad (25)$$

- **Independent events:**

$$P(AB) = P(A)P(B) \quad (26)$$

Generally, the joint probability (intersection) of any number of independent events is the product of their individual probabilities:

$$P(E_1 \cap E_2 \cap \cdots \cap E_n) = P(E_1)P(E_2) \cdots P(E_n) \quad (27)$$

- **Multiplication rule:**

$$P(AB) = P(A|B)P(B) = P(B|A)P(A) \quad (28)$$

# Recap (cont.)

- Total probability:

$$P(E) = P(E|A_1)P(A_1) + P(E|A_2)P(A_2) + \cdots + P(E|A_n)P(A_n) \quad (29)$$

# Recap (cont.)

- Total probability:

$$P(E) = P(E|A_1)P(A_1) + P(E|A_2)P(A_2) + \cdots + P(E|A_n)P(A_n) \quad (29)$$

- Bayes' Theorem:

$$P(A_1|E) = \frac{P(E|A_1)P(A_1)}{P(E|A_1)P(A_1) + P(E|A_2)P(A_2) + \cdots + P(E|A_n)P(A_n)} \quad (30)$$