

CEE 260/MIE 273: Probability and Statistics in Civil Engineering

Lecture 3B: The Normal Distribution

Prof. Oke

UMassAmherst

College of Engineering

September 25, 2025

Outline

- ① The normal distribution
- ② Standard normal distribution
- ③ Computing normal probabilities
- ④ More Examples
- ⑤ Outlook

Reading: OpenIntro Statistics 4.1

Normal (Gaussian) distribution

Definition

Normal (Gaussian) distribution

Definition

Denoted as $\mathcal{N}(\mu, \sigma^2)$ or $\mathcal{N}(\mu, \sigma)$, the normal distribution is continuous with PDF:

Normal (Gaussian) distribution

Definition

Denoted as $\mathcal{N}(\mu, \sigma^2)$ or $\mathcal{N}(\mu, \sigma)$, the normal distribution is continuous with PDF:

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

Normal (Gaussian) distribution

Definition

Denoted as $\mathcal{N}(\mu, \sigma^2)$ or $\mathcal{N}(\mu, \sigma)$, the normal distribution is continuous with PDF:

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} \equiv \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right] \quad -\infty < x < \infty \quad (1)$$

Normal (Gaussian) distribution

Definition

Denoted as $\mathcal{N}(\mu, \sigma^2)$ or $\mathcal{N}(\mu, \sigma)$, the normal distribution is continuous with PDF:

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} \equiv \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right] \quad -\infty < x < \infty \quad (1)$$

where μ and σ^2 (or σ) are its parameters (mean and variance (or standard deviation)).



Normal (Gaussian) distribution

Definition

Denoted as $\mathcal{N}(\mu, \sigma^2)$ or $\mathcal{N}(\mu, \sigma)$, the normal distribution is continuous with PDF:

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} \equiv \frac{1}{\sigma\sqrt{2\pi}} \exp\left[-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right] \quad -\infty < x < \infty \quad (1)$$

where μ and σ^2 (or σ) are its parameters (mean and variance (or standard deviation)).



Example 1: Normal distribution parameters

Example 1: Normal distribution parameters

Example 1: Normal distribution parameters

- (a) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$.
What are the mean and standard deviation of this distribution?

Example 1: Normal distribution parameters

- (a) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$.
What are the mean and standard deviation of this distribution? **Mean:** $\mu = 0$,
standard deviation: $\sigma = 1$

Example 1: Normal distribution parameters

- (a) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$.
What are the mean and standard deviation of this distribution? **Mean: $\mu = 0$, standard deviation: $\sigma = 1$**
- (b) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 2, \sigma^2 = 4)$.
What are the mean and variance of this distribution?

Example 1: Normal distribution parameters

- (a) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$.
What are the mean and standard deviation of this distribution? **Mean:** $\mu = 0$,
standard deviation: $\sigma = 1$
- (b) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 2, \sigma^2 = 4)$.
What are the mean and variance of this distribution? **Mean:** $\mu = 2$, **variance:**
 $\sigma^2 = 4$

Example 1: Normal distribution parameters

- (a) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$.
What are the mean and standard deviation of this distribution? **Mean: $\mu = 0$, standard deviation: $\sigma = 1$**
- (b) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 2, \sigma^2 = 4)$.
What are the mean and variance of this distribution? **Mean: $\mu = 2$, variance: $\sigma^2 = 4$**




Figure: Standard normal distribution $\mathcal{N}(0, 1)$

Example 1: Normal distribution parameters

- (a) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 0, \sigma^2 = 1)$.
What are the mean and standard deviation of this distribution? **Mean: $\mu = 0$, standard deviation: $\sigma = 1$**
- (b) A random variable X is normally distributed as: $X \sim \mathcal{N}(\mu = 2, \sigma^2 = 4)$.
What are the mean and variance of this distribution? **Mean: $\mu = 2$, variance: $\sigma^2 = 4$**





Figure: Standard normal distribution $\mathcal{N}(0, 1)$ **Figure:** Normal distribution with $\mu = 2$, $\sigma = 2$

CDF of a normal distribution

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx =$$

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx =$$

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx = \int_{-\infty}^x \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] dx \quad (2)$$

- There is no closed-form solution to this integral

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx = \int_{-\infty}^x \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] dx \quad (2)$$

- There is no closed-form solution to this integral
- So, in texts/tables, we denote the **standard normal** CDF as $\Phi(z)$, where:

$$\Phi(z) = \int_{-\infty}^z f_Z(z) \frac{1}{1\sqrt{2\pi}} \exp \left[-\frac{1}{2} z^2 \right] dz$$

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx = \int_{-\infty}^x \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] dx \quad (2)$$

- There is no closed-form solution to this integral
- So, in texts/tables, we denote the **standard normal** CDF as $\Phi(z)$, where:

$$\Phi(z) = \int_{-\infty}^z f_Z(z) \frac{1}{1\sqrt{2\pi}} \exp \left[-\frac{1}{2} z^2 \right] dz$$

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx = \int_{-\infty}^x \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] dx \quad (2)$$

- There is no closed-form solution to this integral
- So, in texts/tables, we denote the **standard normal** CDF as $\Phi(z)$, where:

$$\Phi(z) = \int_{-\infty}^z f_Z(z) \frac{1}{1\sqrt{2\pi}} \exp \left[-\frac{1}{2} z^2 \right] dz = P(Z \leq z) \quad (3)$$

where

CDF of a normal distribution

- The CDF of a normal distribution is the integral of the PDF:

$$F_X(x) = P(X \leq x) = \int_{-\infty}^x f_X(x) dx = \int_{-\infty}^x \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] dx \quad (2)$$

- There is no closed-form solution to this integral
- So, in texts/tables, we denote the **standard normal** CDF as $\Phi(z)$, where:

$$\Phi(z) = \int_{-\infty}^z f_Z(z) \frac{1}{1\sqrt{2\pi}} \exp \left[-\frac{1}{2} z^2 \right] dz = P(Z \leq z) \quad (3)$$

where

$$z = \frac{x - \mu}{\sigma} \quad (4)$$

- The standardized normal variable z is often referred to as the Z -score

Standard normal distribution

If a random variable X has a normal distribution $\mathcal{N}(\mu, \sigma^2)$, then the r.v. Z has a **standard normal distribution** if

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1) \quad (5)$$

Standard normal distribution

If a random variable X has a normal distribution $\mathcal{N}(\mu, \sigma^2)$, then the r.v. Z has a **standard normal distribution** if

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1) \quad (5)$$

The **standardized normal** therefore has a mean of 0 and variance of 1.

Standard normal distribution

If a random variable X has a normal distribution $\mathcal{N}(\mu, \sigma^2)$, then the r.v. Z has a **standard normal distribution** if

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1) \quad (5)$$

The **standardized normal** therefore has a mean of 0 and variance of 1. Its PDF is thus:

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2} \quad -\infty < z < \infty \quad (6)$$

Standard normal distribution

If a random variable X has a normal distribution $\mathcal{N}(\mu, \sigma^2)$, then the r.v. Z has a **standard normal distribution** if

$$Z = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1) \quad (5)$$

The **standardized normal** therefore has a mean of 0 and variance of 1. Its PDF is thus:

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2} \quad -\infty < z < \infty \quad (6)$$

The CDF Φ of the standard normal variate Z is given by:

$$\Phi(z) = F_Z(z) \equiv P(Z \leq z) \quad (7)$$

Example 2: Computing the Z -score

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

$$z = \frac{x - \mu}{\sigma} =$$

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

$$z = \frac{x - \mu}{\sigma} = \frac{5 - 10}{\sqrt{4}}$$

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{5 - 10}{\sqrt{4}} \\ &= -\frac{5}{2} = \boxed{-2.5} \end{aligned}$$

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{5 - 10}{\sqrt{4}} \\ &= -\frac{5}{2} = \boxed{-2.5} \end{aligned}$$

This means that the sample $x = 5$ is 2.5 standard deviations below the mean

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{5 - 10}{\sqrt{4}} \\ &= -\frac{5}{2} = \boxed{-2.5} \end{aligned}$$

This means that the sample $x = 5$ is 2.5 standard deviations below the mean




Figure: Normal distribution with $\mu = 10$, $\sigma^2 = 4$, and $x = 5$ indicated

Example 2: Computing the Z-score

If $X \sim \mathcal{N}(10, \sigma^2 = 4)$, what is the Z-score of a sample $x = 5$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{5 - 10}{\sqrt{4}} \\ &= -\frac{5}{2} = \boxed{-2.5} \end{aligned}$$

This means that the sample $x = 5$ is 2.5 standard deviations below the mean




Figure: Normal distribution with $\mu = 10$, $\sigma^2 = 4$, and $x = 5$ indicated




Figure: Standard normal distribution with $z = -2.5$ indicated

Example 3: Computing the Z -score

Example 3: Computing the Z -score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z -score of a sample $x = 8$?

Example 3: Computing the Z-score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z-score of a sample $x = 8$?

$$z = \frac{x - \mu}{\sigma} =$$

Example 3: Computing the Z-score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z-score of a sample $x = 8$?

$$z = \frac{x - \mu}{\sigma} = \frac{8 - 6}{\sqrt{9}}$$

Example 3: Computing the Z-score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z-score of a sample $x = 8$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{8 - 6}{\sqrt{9}} \\ &= \frac{2}{3} = \boxed{0.67} \end{aligned}$$

Example 3: Computing the Z-score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z-score of a sample $x = 8$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{8 - 6}{\sqrt{9}} \\ &= \frac{2}{3} = \boxed{0.67} \end{aligned}$$

This means that the sample $x = 8$ is 0.67 standard deviations above the mean

Example 3: Computing the Z-score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z-score of a sample $x = 8$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{8 - 6}{\sqrt{9}} \\ &= \frac{2}{3} = \boxed{0.67} \end{aligned}$$

This means that the sample $x = 8$ is 0.67 standard deviations above the mean




Figure: Normal distribution with $\mu = 6$, $\sigma^2 = 9$, and $x = 8$ indicated

Example 3: Computing the Z-score

If $X \sim \mathcal{N}(\mu = 6, \sigma^2 = 9)$, what is the Z-score of a sample $x = 8$?

$$\begin{aligned} z &= \frac{x - \mu}{\sigma} = \frac{8 - 6}{\sqrt{9}} \\ &= \frac{2}{3} = \boxed{0.67} \end{aligned}$$

This means that the sample $x = 8$ is 0.67 standard deviations above the mean




Figure: Normal distribution with $\mu = 6$, $\sigma^2 = 9$, and $x = 8$ indicated




Figure: Standard normal distribution with $z = 0.67$ indicated

Example 4: Finding the normal variate from a Z -score

Example 4: Finding the normal variate from a Z -score

A normal r.v. $X \sim \mathcal{N}(\mu = 2, \sigma = 3)$ has a Z -score of $z = 1.25$.

Example 4: Finding the normal variate from a Z -score

A normal r.v. $X \sim \mathcal{N}(\mu = 2, \sigma = 3)$ has a Z -score of $z = 1.25$. What is the corresponding x value?

Example 4: Finding the normal variate from a Z-score

A normal r.v. $X \sim \mathcal{N}(\mu = 2, \sigma = 3)$ has a Z-score of $z = 1.25$. What is the corresponding x value?




Figure: Standard normal distribution with $z = 1.25$ indicated

Example 4: Finding the normal variate from a Z-score

A normal r.v. $X \sim \mathcal{N}(\mu = 2, \sigma = 3)$ has a Z-score of $z = 1.25$. What is the corresponding x value?




Figure: Standard normal distribution with $z = 1.25$ indicated




Figure: Normal distribution with $\mu = 2$, $\sigma = 3$, and $x = 5.75$ indicated

$$z = \frac{x - \mu}{\sigma}$$

Example 4: Finding the normal variate from a Z-score

A normal r.v. $X \sim \mathcal{N}(\mu = 2, \sigma = 3)$ has a Z-score of $z = 1.25$. What is the corresponding x value?




Figure: Standard normal distribution with $z = 1.25$ indicated




Figure: Normal distribution with $\mu = 2$, $\sigma = 3$, and $x = 5.75$ indicated

$$z = \frac{x - \mu}{\sigma}$$

$$x = \mu + z\sigma$$

Example 4: Finding the normal variate from a Z-score

A normal r.v. $X \sim \mathcal{N}(\mu = 2, \sigma = 3)$ has a Z-score of $z = 1.25$. What is the corresponding x value?




Figure: Standard normal distribution with $z = 1.25$ indicated




Figure: Normal distribution with $\mu = 2$, $\sigma = 3$, and $x = 5.75$ indicated

$$z = \frac{x - \mu}{\sigma}$$

$$x = \mu + z\sigma = 2 + 1.25(3) = \boxed{5.75}$$

68-95-99.7 rule

68-95-99.7 rule

The probabilities of a normal r.v. within ± 1 , ± 2 and ± 3 standard deviations are 68.3%, 95.4% and 99.7%, respectively.

68-95-99.7 rule

The probabilities of a normal r.v. within ± 1 , ± 2 and ± 3 standard deviations are 68.3%, 95.4% and 99.7%, respectively. This is known as the **68-95-99.7 rule** or the **empirical rule**.




Figure: Standard normal distribution with color-coded regions showing probability percentages

Probability of a normal random variable

Probability of a normal random variable

The probability that a normal r.v. lies within a certain interval is given by the *area* under the PDF in that interval.

Probability of a normal random variable

The probability that a normal r.v. lies within a certain interval is given by the *area* under the PDF in that interval.




Figure: Standard normal distribution with $z = 0.5$ indicated and $P(Z \leq 0.5)$ shaded

Probability of a normal random variable

The probability that a normal r.v. lies within a certain interval is given by the *area* under the PDF in that interval.




Figure: Standard normal distribution with $z = 0.5$ indicated and $P(Z \leq 0.5)$ shaded

- Recall that the area under the PDF within a given interval is the CDF evaluated in that range

Probability of a normal random variable

The probability that a normal r.v. lies within a certain interval is given by the *area* under the PDF in that interval.




Figure: Standard normal distribution with $z = 0.5$ indicated and $P(Z \leq 0.5)$ shaded

- Recall that the area under the PDF within a given interval is the CDF evaluated in that range
- Thus, in the above figure: $p = P(Z \leq z_p) = \Phi(z_p)$

Probability of a normal random variable (cont.)

Probability of a normal random variable (cont.)

Given a normal r.v. $X \sim \mathcal{N}(\mu, \sigma^2)$:

Probability of a normal random variable (cont.)

Given a normal r.v. $X \sim \mathcal{N}(\mu, \sigma^2)$:

$$P(a < X \leq b) = \frac{1}{\sigma\sqrt{2\pi}} \int_a^b e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx \quad (8)$$

Probability of a normal random variable (cont.)

Given a normal r.v. $X \sim \mathcal{N}(\mu, \sigma^2)$:

$$P(a < X \leq b) = \frac{1}{\sigma\sqrt{2\pi}} \int_a^b e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx \quad (8)$$

Substituting $z = \frac{x-\mu}{\sigma}$ and $dx = \sigma dz$, we obtain:

Probability of a normal random variable (cont.)

Given a normal r.v. $X \sim \mathcal{N}(\mu, \sigma^2)$:

$$P(a < X \leq b) = \frac{1}{\sigma\sqrt{2\pi}} \int_a^b e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx \quad (8)$$

Substituting $z = \frac{x-\mu}{\sigma}$ and $dx = \sigma dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(a-\mu)/\sigma}^{(b-\mu)/\sigma} e^{-\frac{1}{2}z^2} dz \quad (9)$$

Probability of a normal random variable (cont.)

Given a normal r.v. $X \sim \mathcal{N}(\mu, \sigma^2)$:

$$P(a < X \leq b) = \frac{1}{\sigma\sqrt{2\pi}} \int_a^b e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx \quad (8)$$

Substituting $z = \frac{x-\mu}{\sigma}$ and $dx = \sigma dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(a-\mu)/\sigma}^{(b-\mu)/\sigma} e^{-\frac{1}{2}z^2} dz \quad (9)$$

Recognizing that the integrand is the PDF of a standard normal distribution, we have:

Probability of a normal random variable (cont.)

Given a normal r.v. $X \sim \mathcal{N}(\mu, \sigma^2)$:

$$P(a < X \leq b) = \frac{1}{\sigma\sqrt{2\pi}} \int_a^b e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} dx \quad (8)$$

Substituting $z = \frac{x-\mu}{\sigma}$ and $dx = \sigma dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(a-\mu)/\sigma}^{(b-\mu)/\sigma} e^{-\frac{1}{2}z^2} dz \quad (9)$$

Recognizing that the integrand is the PDF of a standard normal distribution, we have:

$$P(a < X \leq b) = \Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right) \quad (10)$$

Example 5a: Normal probabilities

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200?

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200?

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200? The Z-score is $z = \frac{1200-1100}{200} = 0.5$

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200? The Z-score is $z = \frac{1200-1100}{200} = 0.5$

Thus,

$$P(X \geq 1200) = 1 - \Phi(.5) =$$

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200? The Z-score is $z = \frac{1200-1100}{200} = 0.5$

Thus,

$$P(X \geq 1200) = 1 - \Phi(.5) = 1 - .695 = \boxed{.3085}$$

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200? The Z-score is $z = \frac{1200-1100}{200} = 0.5$

Thus,

$$P(X \geq 1200) = 1 - \Phi(.5) = 1 - .695 = \boxed{.3085}$$

In Python, you can compute this probability using the `scipy.stats` library:

```
import scipy.stats as stats  
p = 1 - stats.norm.cdf(1200, 1100, 200)
```

The first 3 arguments of `stats.norm.cdf` are the value, mean (`textttloc`), and standard deviation (`textttscale`), respectively.

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200? The Z-score is $z = \frac{1200-1100}{200} = 0.5$

Thus,

$$P(X \geq 1200) = 1 - \Phi(.5) = 1 - .695 = \boxed{.3085}$$

In Python, you can compute this probability using the `scipy.stats` library:

```
import scipy.stats as stats
p = 1 - stats.norm.cdf(1200, 1100, 200)
```

The first 3 arguments of `stats.norm.cdf` are the value, mean (`textttloc`), and standard deviation (`textttscale`), respectively. OR, you can use the Z-score (default `mean=0`, `std=1`):

```
import scipy.stats as stats
p = 1 - stats.norm.cdf(0.5)
```

Example 5a: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (a) What is the probability that a randomly selected student has a score that is at least 1200? The Z-score is $z = \frac{1200-1100}{200} = 0.5$

Thus,



$$P(X \geq 1200) = 1 - \Phi(.5) = 1 - .695 = \boxed{.3085}$$

In Python, you can compute this probability using the `scipy.stats` library:

```
import scipy.stats as stats  
p = 1 - stats.norm.cdf(1200, 1100, 200)
```

The first 3 arguments of `stats.norm.cdf` are the value, mean (`textttloc`), and standard deviation (`textttscale`), respectively. OR, you can use the Z-score (default `mean=0`, `std=1`):

```
import scipy.stats as stats  
p = 1 - stats.norm.cdf(0.5)
```



Example 5b: Normal probabilities

Example 5b: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

Example 5b: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (b) What is the probability that another randomly selected student's score is greater than 600 but less than 1200?

Example 5b: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (b) What is the probability that another randomly selected student's score is greater than 600 but less than 1200?

$$P(600 \leq X < 1200) = \Phi\left(\frac{1200 - 1100}{200}\right) - \Phi\left(\frac{600 - 1100}{200}\right)$$

Example 5b: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (b) What is the probability that another randomly selected student's score is greater than 600 but less than 1200?

$$\begin{aligned} P(600 \leq X < 1200) &= \Phi\left(\frac{1200 - 1100}{200}\right) - \Phi\left(\frac{600 - 1100}{200}\right) \\ &= \Phi(.5) - \Phi(-2.5) \end{aligned}$$

Example 5b: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (b) What is the probability that another randomly selected student's score is greater than 600 but less than 1200?

$$\begin{aligned} P(600 \leq X < 1200) &= \Phi\left(\frac{1200 - 1100}{200}\right) - \Phi\left(\frac{600 - 1100}{200}\right) \\ &= \Phi(.5) - \Phi(-2.5) = \boxed{.6853} \end{aligned}$$

In Python:

```
from scipy.stats import norm  
p = norm.cdf(1200, 1100, 200)  
    - norm.cdf(600, 1100, 200)
```

Example 5b: Normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (b) What is the probability that another randomly selected student's score is greater than 600 but less than 1200?

$$\begin{aligned} P(600 \leq X < 1200) &= \Phi\left(\frac{1200 - 1100}{200}\right) - \Phi\left(\frac{600 - 1100}{200}\right) \\ &= \Phi(.5) - \Phi(-2.5) = \boxed{.6853} \end{aligned}$$

In Python:

```
from scipy.stats import norm
p = norm.cdf(1200, 1100, 200)
    - norm.cdf(600, 1100, 200)
```

Example 5b: Normal probabilities


SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

- (b) What is the probability that another randomly selected student's score is greater than 600 but less than 1200?

$$\begin{aligned} P(600 \leq X < 1200) &= \Phi\left(\frac{1200 - 1100}{200}\right) - \Phi\left(\frac{600 - 1100}{200}\right) \\ &= \Phi(.5) - \Phi(-2.5) = \boxed{.6853} \end{aligned}$$

In Python:

```
from scipy.stats import norm  
p = norm.cdf(1200, 1100, 200)  
    - norm.cdf(600, 1100, 200)
```



Example 5b: Normal probabilities (cont)

Example 5b: Normal probabilities (cont)

(b) OR, you can use the Z-scores (default mean=0, std=1):

```
import scipy.stats as stats  
p = norm.cdf(0.5) - norm.cdf(-2.5)
```




Figure: Standard normal PDF with probability area shaded

Example 5b: Normal probabilities (cont)

(b) OR, you can use the Z-scores (default mean=0, std=1):

```
import scipy.stats as stats  
p = norm.cdf(0.5) - norm.cdf(-2.5)
```




Figure: Standard normal PDF with probability area shaded




Figure: Standard normal CDF with corresponding probability values marked

Example 6: Inverse normal probabilities

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

(c) If the probability of an SAT score lower than x is 0.4, find x .

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

(c) If the probability of an SAT score lower than x is 0.4, find x .

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

(c) If the probability of an SAT score lower than x is 0.4, find x .

$$z = \Phi^{-1}(.4) = -.2533 = \frac{x - \mu}{\sigma}$$

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

(c) If the probability of an SAT score lower than x is 0.4, find x .

$$z = \Phi^{-1}(.4) = -.2533 = \frac{x - \mu}{\sigma}$$

$$\therefore x = z\sigma + \mu = -.2533(200) + 1100 = \boxed{1049}$$

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

(c) If the probability of an SAT score lower than x is 0.4, find x .

$$z = \Phi^{-1}(.4) = -.2533 = \frac{x - \mu}{\sigma}$$

$$\therefore x = z\sigma + \mu = -.2533(200) + 1100 = \boxed{1049}$$

```
from scipy.stats import norm  
p = norm.ppf(0.4, 1100, 200)
```

Example 6: Inverse normal probabilities

SAT scores are normally distributed as $X \sim \mathcal{N}(1100, \sigma = 200)$.

(c) If the probability of an SAT score lower than x is 0.4, find x .

$$z = \Phi^{-1}(.4) = -.2533 = \frac{x - \mu}{\sigma}$$

$$\therefore x = z\sigma + \mu = -.2533(200) + 1100 = \boxed{1049}$$

```
from scipy.stats import norm  
p = norm.ppf(0.4, 1100, 200)
```




Figure: Normal distribution with $\mu = 1100$, $\sigma = 200$, and $P(X \leq 1049)$ shaded

Example 6: Inverse normal probabilities (cont.)

Example 6: Inverse normal probabilities (cont.)

- (c) You can think of the inverse CDF as finding the x value that corresponds to a given percentile.

Example 6: Inverse normal probabilities (cont.)

- (c) You can think of the inverse CDF as finding the x value that corresponds to a given percentile.




Figure: Normal distribution CDF showing the 40th percentile at $x = 1049$

More on the normal CDF

More on the normal CDF




Figure: Comparison of normal distribution CDFs with different parameters

More on the normal CDF




Figure: Comparison of normal distribution CDFs with different parameters

- The standard CDF is the blue curve in the above figure

More on the normal CDF




Figure: Comparison of normal distribution CDFs with different parameters

- The standard normal CDF is the blue curve in the above figure
- Quantiles can be read off the plot (e.g. the median is the value of X corresponding to the y value of 0.5)

More on the normal CDF




Figure: Comparison of normal distribution CDFs with different parameters

- The standard normal CDF is the blue curve in the above figure
- Quantiles can be read off the plot (e.g. the median is the value of X corresponding to the y value of 0.5)
- $\Phi(-z) = 1 - \Phi(z)$

More on the normal CDF




Figure: Comparison of normal distribution CDFs with different parameters

- The standard normal CDF is the blue curve in the above figure
- Quantiles can be read off the plot (e.g. the median is the value of X corresponding to the y value of 0.5)
- $\Phi(-z) = 1 - \Phi(z)$
- $z = \Phi^{-1}(p) = -\Phi^{-1}(1 - p)$

Example 7: Probability of flooding

Example 7: Probability of flooding

The drainage from a community during a storm is a normal random variable estimated to have a mean of 1.2 million gallons per day (mgd) and an SD of 0.4 mgd. If the storm drain system is designed with a maximum drainage capacity of 1.5 mgd:

Example 7: Probability of flooding

The drainage from a community during a storm is a normal random variable estimated to have a mean of 1.2 million gallons per day (mgd) and an SD of 0.4 mgd. If the storm drain system is designed with a maximum drainage capacity of 1.5 mgd:

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?
- (b) Find $P(1.0 < X \leq 1.6)$.
- (c) Find the 90th-percentile drainage load from the community during a storm.

Example 7: Probability of flooding (cont.)

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution





Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(X > 1.5)$ shaded

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution



$$P(X > 1.5) = 1 - P(X \leq 1.5)$$


Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(X > 1.5)$ shaded

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution



$$\begin{aligned} P(X > 1.5) &= 1 - P(X \leq 1.5) \\ &= 1 - \Phi\left(\frac{1.5 - 1.2}{0.4}\right) \end{aligned}$$


Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(X > 1.5)$ shaded

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution



$$\begin{aligned} P(X > 1.5) &= 1 - P(X \leq 1.5) \\ &= 1 - \Phi\left(\frac{1.5 - 1.2}{0.4}\right) \\ &= 1 - \Phi(0.75) \end{aligned}$$


Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(X > 1.5)$ shaded

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution



$$\begin{aligned} P(X > 1.5) &= 1 - P(X \leq 1.5) \\ &= 1 - \Phi\left(\frac{1.5 - 1.2}{0.4}\right) \\ &= 1 - \Phi(0.75) \\ &= 1 - 0.7734 = \boxed{0.227} \end{aligned}$$

Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(X > 1.5)$ shaded

Example 7: Probability of flooding (cont.)

Given $\mu = 1.2$ and $\sigma = 0.4$.

- (a) What is the underlying probability of flooding during a storm that is assumed in the design of the drainage system?

Solution




Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(X > 1.5)$ shaded

$$\begin{aligned} P(X > 1.5) &= 1 - P(X \leq 1.5) \\ &= 1 - \Phi\left(\frac{1.5 - 1.2}{0.4}\right) \\ &= 1 - \Phi(0.75) \\ &= 1 - 0.7734 = \boxed{0.227} \end{aligned}$$

In Python: `1 - norm.cdf(1.5, 1.2, 0.4)`

Example 7: Probability of flooding (cont.)

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution





Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X < 1.6)$ shaded

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution




$$p = \Phi\left(\frac{1.6 - 1.2}{0.4}\right) - \Phi\left(\frac{1.0 - 1.2}{0.4}\right)$$

Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X < 1.6)$ shaded

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution



$$\begin{aligned} p &= \Phi\left(\frac{1.6 - 1.2}{0.4}\right) \\ &\quad - \Phi\left(\frac{1.0 - 1.2}{0.4}\right) \\ &= \Phi(1.0) - \Phi(-0.5) \end{aligned}$$

Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X < 1.6)$ shaded

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution




Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X < 1.6)$ shaded

$$\begin{aligned} p &= \Phi\left(\frac{1.6 - 1.2}{0.4}\right) \\ &\quad - \Phi\left(\frac{1.0 - 1.2}{0.4}\right) \\ &= \Phi(1.0) - \Phi(-0.5) \\ &= 0.8413 - [1 - \Phi(0.5)] \end{aligned}$$

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution




Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X < 1.6)$ shaded

$$\begin{aligned} p &= \Phi\left(\frac{1.6 - 1.2}{0.4}\right) \\ &\quad - \Phi\left(\frac{1.0 - 1.2}{0.4}\right) \\ &= \Phi(1.0) - \Phi(-0.5) \\ &= 0.8413 - [1 - \Phi(0.5)] \\ &= 0.8413 - (1 - 0.6915) \end{aligned}$$

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution




Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X \leq 1.6)$ shaded

$$\begin{aligned} p &= \Phi\left(\frac{1.6 - 1.2}{0.4}\right) \\ &\quad - \Phi\left(\frac{1.0 - 1.2}{0.4}\right) \\ &= \Phi(1.0) - \Phi(-0.5) \\ &= 0.8413 - [1 - \Phi(0.5)] \\ &= 0.8413 - (1 - 0.6915) \\ &= 0.8413 - 0.3085 \end{aligned}$$

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution




Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X \leq 1.6)$ shaded

$$\begin{aligned} p &= \Phi\left(\frac{1.6 - 1.2}{0.4}\right) \\ &\quad - \Phi\left(\frac{1.0 - 1.2}{0.4}\right) \\ &= \Phi(1.0) - \Phi(-0.5) \\ &= 0.8413 - [1 - \Phi(0.5)] \\ &= 0.8413 - (1 - 0.6915) \\ &= 0.8413 - 0.3085 \\ &= 0.5328 \approx \boxed{0.533} \end{aligned}$$

Example 7: Probability of flooding (cont.)

(b) Find $p = P(1.0 < X \leq 1.6)$:

Solution




Figure: Normal distribution with $\mu = 1.2$, $\sigma = 0.4$, and $P(1.0 < X \leq 1.6)$ shaded

$$\begin{aligned} p &= \Phi\left(\frac{1.6 - 1.2}{0.4}\right) \\ &\quad - \Phi\left(\frac{1.0 - 1.2}{0.4}\right) \\ &= \Phi(1.0) - \Phi(-0.5) \\ &= 0.8413 - [1 - \Phi(0.5)] \\ &= 0.8413 - (1 - 0.6915) \\ &= 0.8413 - 0.3085 \\ &= 0.5328 \approx \boxed{0.533} \end{aligned}$$

In Python: `norm.cdf(1.6, 1.2, 0.4) - norm.cdf(1.0, 1.2, 0.4)`

Example 7: Probability of flooding (cont.)

Example 7: Probability of flooding (cont.)

- (c) Find the 90th-percentile drainage load from the community during a storm.

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.




Figure: Normal CDF showing the 90th percentile at $x = 1.71$

Example 7: Probability of flooding (cont.)

Example 7: Probability of flooding (cont.)

- (c) Find the 90th-percentile drainage load from the community during a storm.

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

Example 7: Probability of flooding (cont.)

- (c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$P(X \leq x_{0.90})$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$P(X \leq x_{0.90}) = \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$P(X \leq x_{0.90}) = \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90$$
$$\Rightarrow \frac{x_{0.90} - 1.2}{0.40}$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$\begin{aligned} P(X \leq x_{0.90}) &= \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90 \\ \Rightarrow \frac{x_{0.90} - 1.2}{0.40} &= \Phi^{-1}(0.90) \end{aligned}$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$\begin{aligned} P(X \leq x_{0.90}) &= \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90 \\ \Rightarrow \frac{x_{0.90} - 1.2}{0.40} &= \Phi^{-1}(0.90) = 1.28 \end{aligned}$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$\begin{aligned} P(X \leq x_{0.90}) &= \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90 \\ \Rightarrow \frac{x_{0.90} - 1.2}{0.40} &= \Phi^{-1}(0.90) = 1.28 \\ &\therefore x_{0.90} \end{aligned}$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$\begin{aligned}P(X \leq x_{0.90}) &= \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90 \\ \Rightarrow \frac{x_{0.90} - 1.2}{0.40} &= \Phi^{-1}(0.90) = 1.28 \\ \therefore x_{0.90} &= 1.28(0.40) + 1.2 =\end{aligned}$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$\begin{aligned}P(X \leq x_{0.90}) &= \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90 \\ \Rightarrow \frac{x_{0.90} - 1.2}{0.40} &= \Phi^{-1}(0.90) = 1.28 \\ \therefore x_{0.90} &= 1.28(0.40) + 1.2 = 1.71 \text{ mgd}\end{aligned}$$

Example 7: Probability of flooding (cont.)

(c) Find the 90th-percentile drainage load from the community during a storm.

Solution

$$\begin{aligned}P(X \leq x_{0.90}) &= \Phi\left(\frac{x_{0.90} - 1.2}{0.40}\right) = 0.90 \\ \Rightarrow \frac{x_{0.90} - 1.2}{0.40} &= \Phi^{-1}(0.90) = 1.28 \\ \therefore x_{0.90} &= 1.28(0.40) + 1.2 = 1.71 \text{ mgd}\end{aligned}$$

In Python:

```
from scipy.stats import norm  
p90 = norm.ppf(0.9, 1.2, 0.4)
```

gives 1.7095 mgd

Example 8: Steel beam reliability

Example 8: Steel beam reliability

Assume the variability E in the lengths of steel beams is normally distributed. What is the precision (in terms of σ) required for a reliability of 99.7%, given that the specified tolerance for a construction project is ± 5 mm?

Example 8: Steel beam reliability

Assume the variability E in the lengths of steel beams is normally distributed. What is the precision (in terms of σ) required for a reliability of 99.7%, given that the specified tolerance for a construction project is ± 5 mm?

Definitions:

- **Precision:** in physical terms is the inverse of the variance (i.e. higher precision means lower variance). In this context, all you need to do is find the standard deviation σ .

Example 8: Steel beam reliability

Assume the variability E in the lengths of steel beams is normally distributed. What is the precision (in terms of σ) required for a reliability of 99.7%, given that the specified tolerance for a construction project is ± 5 mm?

Definitions:

- **Precision:** in physical terms is the inverse of the variance (i.e. higher precision means lower variance). In this context, all you need to do is find the standard deviation σ .
- **Reliability:** probability that the deviation in the length of a beam meets (falls within) the specified tolerance

Recap of normal distribution

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (12)$$

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (12)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (12)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol Φ (“phi”) is used to represent the CDF of the **standard normal distribution**, whose values can be looked up in a table.

Recap of normal distribution

- The **PDF** of the normal distribution (parameters μ and σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (11)$$

- The parameters of a normal distribution $\mathcal{N}(\mu, \sigma^2)$ correspond to its mean and variance, respectively.
- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “Z-score”:

$$Z = \frac{X - \mu}{\sigma} \quad (12)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol Φ (“phi”) is used to represent the CDF of the *standard normal distribution*, whose values can be looked up in a table.
- In Python, the `scipy.stats.norm.cdf(x, mu, sigma)` and `scipy.stats.norm.ppf(p, mu, sigma)` can be used to compute probabilities and inverse CDFs of the normal distribution, respectively.