

CEE 260/MIE 273: Probability and Statistics in Civil Engineering

Lecture M3c: Lognormal and Exponential Distributions

Jimi Oke

UMassAmherst

College of Engineering

October 6, 2026

Outline

① Introduction

② Lognormal distribution

③ Exponential distribution

④ Outlook

Recap of normal distribution

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol Φ (“phi”) is used to represent the CDF of the **standard normal distribution**, whose values can be looked up in a table.

Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean μ and variance σ^2) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol Φ (“phi”) is used to represent the CDF of the *standard normal distribution*, whose values can be looked up in a table.
- In Python, the `norm.cdf(x, mu, sigma)` and `norm.ppf(p, mu, sigma)` can be used to compute probabilities and inverse CDFs of the normal distribution, respectively.

Using the standard normal CDF probability table

Using the standard normal CDF probability table


- First convert the random variable to its Z -score

Using the standard normal CDF probability table

- First convert the random variable to its Z -score
- Find the corresponding value in the table

Using the standard normal CDF probability table

- First convert the random variable to its Z -score
- Find the corresponding value in the table



Objectives of today's lecture

Objectives of today's lecture

- Lognormal distribution

Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions

Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions
- Exponential distribution

Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions
- Exponential distribution
- Memoryless property of exponential distribution

PDF of the lognormal distribution

Definition

PDF of the lognormal distribution

Definition

A random variable X that is lognormally distributed with the parameters μ and σ (denoted $X \sim \mathcal{LN}(\mu, \sigma)$) has the PDF:

PDF of the lognormal distribution

Definition

A random variable X that is lognormally distributed with the parameters μ and σ (denoted $X \sim \mathcal{LN}(\mu, \sigma)$) has the PDF:


$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$

PDF of the lognormal distribution

Definition

A random variable X that is lognormally distributed with the parameters μ and σ (denoted $X \sim \mathcal{LN}(\mu, \sigma)$) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$




PDF of the lognormal distribution

Definition

A random variable X that is lognormally distributed with the parameters μ and σ (denoted $X \sim \mathcal{LN}(\mu, \sigma)$) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$




PDF of the lognormal distribution

Definition

A random variable X that is lognormally distributed with the parameters μ and σ (denoted $X \sim \mathcal{LN}(\mu, \sigma)$) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[-\frac{1}{2} \left(\frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$



Mean, median and variance of a lognormal distribution

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of X is given by:

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of X is given by:

$$\mathbb{V}(X) =$$

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of X is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of X is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of X is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

Notes

- μ and σ are the mean and standard deviation of the associated normal distribution of $\ln(X)$.

Mean, median and variance of a lognormal distribution

Let X be a lognormally distributed random variable with parameters μ and σ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of X is given by

$$\mathbb{E}(X) = \exp \left[\mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of X is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of X is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

Notes

- μ and σ are the mean and standard deviation of the associated normal distribution of $\ln(X)$.
- Thus, if $X \sim \mathcal{LN}(\mu, \sigma)$, then $\ln(X) \sim \mathcal{N}(\mu, \sigma)$

Example 1: Mean and variance of lognormal distribution (1)

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\Rightarrow \ln(5) =$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\Rightarrow \ln(5) = \mu$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\Rightarrow \ln(5) = \mu$$

$\therefore$ The mean is given by

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\begin{aligned} \therefore \text{The mean is given by } \mathbb{E}(X) &= e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21} \\ &= 5(e^{0.21}) \end{aligned}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\begin{aligned} \therefore \text{The mean is given by } \mathbb{E}(X) &= e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21} \\ &= 5(e^{0.21}) = \boxed{6.17 \text{ days}} \end{aligned}$$

Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What are the mean and variance of its distribution?

Solution

First, we find the parameter μ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\begin{aligned} \therefore \text{The mean is given by } \mathbb{E}(X) &= e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21} \\ &= 5(e^{0.21}) = \boxed{6.17 \text{ days}} \end{aligned}$$

(Note: $\ln$ denotes the natural logarithm and is pronounced "lawn")

Example 1: Mean and variance of lognormal distribution (2)

Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

The variance is given by:

Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

The variance is given by:

$$\mathbb{V}(X) = (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2])$$

Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

The variance is given by:

$$\begin{aligned}\mathbb{V}(X) &= (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2]) \\ &= (\exp(0.42) - 1)(\exp(2 \ln(5) + 0.42))\end{aligned}$$

Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

The variance is given by:

$$\begin{aligned}
 \mathbb{V}(X) &= (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2]) \\
 &= (\exp(0.42) - 1)(\exp(2 \ln(5) + 0.42)) \\
 &= \boxed{19.86 \text{ days}^2}
 \end{aligned}$$

Relationship between normal and lognormal distributions

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma)$$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies$$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable X is **lognormally** distributed with the **parameters** μ and σ , then:
 - $\mu = \mathbb{E}(\ln(X))$ and $\sigma^2 = \mathbb{V}(\ln(X))$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable X is **lognormally** distributed with the **parameters** μ and σ , then:
 - $\mu = \mathbb{E}(\ln(X))$ and $\sigma^2 = \mathbb{V}(\ln(X))$
 - $\ln(X)$ is **normally** distributed with the same parameters.

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable X is **lognormally** distributed with the **parameters** μ and σ , then:
 - $\mu = \mathbb{E}(\ln(X))$ and $\sigma^2 = \mathbb{V}(\ln(X))$
 - $\ln(X)$ is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma)$$

Relationship between normal and lognormal distributions

- If random variable X is **normally** distributed with the parameters μ and σ^2 , then:
 - $\mu = \mathbb{E}(X)$ and $\sigma^2 = \mathbb{V}(X)$
 - e^X is **lognormally** distributed with the same parameters:


$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable X is **lognormally** distributed with the **parameters** μ and σ , then:
 - $\mu = \mathbb{E}(\ln(X))$ and $\sigma^2 = \mathbb{V}(\ln(X))$
 - $\ln(X)$ is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma) \quad (8)$$

Relationship between normal and lognormal (cont.)

Relationship between normal and lognormal (cont.)



Positive skewness of lognormal distribution

Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed

Positive skewness of lognormal distribution


- The lognormal distribution is positively skewed
- Its mean is always greater than its median

Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median

Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median



Probability of a lognormal random variate

Given a r.v. X that is lognormally distributed with parameters μ and σ^2 :

$$P(a < X \leq b) =$$

Probability of a lognormal random variate

Given a r.v. X that is lognormally distributed with parameters μ and σ^2 :

$$P(a < X \leq b) = \frac{1}{\sigma_X \sqrt{2\pi}}$$

Probability of a lognormal random variate

Given a r.v. X that is lognormally distributed with parameters μ and σ^2 :

$$P(a < X \leq b) = \frac{1}{\sigma_X \sqrt{2\pi}} \int_a^b \exp \left[-\frac{1}{2} \left(\frac{\ln(x) - \mu}{\sigma} \right)^2 \right] dx \quad (9)$$

Probability of a lognormal random variate

Given a r.v. X that is lognormally distributed with parameters μ and σ^2 :

$$P(a < X \leq b) = \frac{1}{\sigma x \sqrt{2\pi}} \int_a^b \exp \left[-\frac{1}{2} \left(\frac{\ln(x) - \mu}{\sigma} \right)^2 \right] dx \quad (9)$$




Figure: Lognormal distribution with $\mu = 0, \sigma = 1$, showing $P(a < X < b)$ where $a = 2$ and $b = 4$

Probability of a lognormal random variate (cont.)

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma}$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b)$$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma}$$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b)$$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right)$$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right) -$$

Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right) - \Phi\left(\frac{\ln a - \mu}{\sigma}\right) \quad (11)$$


Probability of a lognormal random variate (cont.)

Substituting $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$, we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right) - \Phi\left(\frac{\ln a - \mu}{\sigma}\right) \quad (11)$$



Example 2: Probability of incubation period

Example 2: Probability of incubation period

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What is the probability that a randomly selected person will show symptoms within 7 days of exposure (i.e., $P(X \leq 7)$)?

Example 2: Probability of incubation period

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and $\sigma^2 = 0.42$. What is the probability that a randomly selected person will show symptoms within 7 days of exposure (i.e., $P(X \leq 7)$)?




Figure: Lognormal distribution with $\mu = \ln(5)$, $\sigma^2 = 0.42$, showing $P(X \leq 7)$

Example 2: Probability of incubation period (cont.)

Example 2: Probability of incubation period (cont.)

Solution (using Python)

Example 2: Probability of incubation period (cont.)

Solution (using Python)

We want to find: $P(X \leq 7) = F(7)$; thus we use the `scipy.stats.lognorm.cdf` function:

Example 2: Probability of incubation period (cont.)

Solution (using Python)

We want to find: $P(X \leq 7) = F(7)$; thus we use the `scipy.stats.lognorm.cdf` function:

```
from scipy import stats
import numpy as np
p = stats.lognorm.cdf(7, s=0.648, scale=5)
```

Example 2: Probability of incubation period (cont.)

Solution (using Python)

We want to find: $P(X \leq 7) = F(7)$; thus we use the `scipy.stats.lognorm.cdf` function:

```
from scipy import stats
import numpy as np
p = stats.lognorm.cdf(7, s=0.648, scale=5)
```

where:

- $s = \sigma = 0.648 = \sqrt{0.42}$ (shape parameter/standard deviation of underlying normal)
- $\text{scale} = \exp(\mu) = \exp(\ln(5)) = 5$ (scale parameter/median)

The result is: $p = 0.691$, i.e., about 69.1% of the people will show symptoms within 7 days of exposure.

Example 2: Probability of incubation period (cont.)

Example 2: Probability of incubation period (cont.)

Solution (using tables)

Example 2: Probability of incubation period (cont.)

Solution (using tables)

We want to find: $P(X \leq 7) = F(7)$; thus we use the relationship:

Example 2: Probability of incubation period (cont.)

Solution (using tables)

We want to find: $P(X \leq 7) = F(7)$; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

Example 2: Probability of incubation period (cont.)

Solution (using tables)

We want to find: $P(X \leq 7) = F(7)$; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

where $\mu = \ln(5) = 1.609$ and $\sigma = \sqrt{0.42} = 0.648$. Thus:

Example 2: Probability of incubation period (cont.)

Solution (using tables)

We want to find: $P(X \leq 7) = F(7)$; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

where $\mu = \ln(5) = 1.609$ and $\sigma = \sqrt{0.42} = 0.648$. Thus:

$$\begin{aligned} P(X \leq 7) &= \Phi\left(\frac{\ln 7 - 1.609}{0.648}\right) \\ &= \Phi(0.539) \\ &\approx 0.7054 \text{ (from standard normal table)} \end{aligned}$$

Example 2: Probability of incubation period (cont.)

Solution (using tables)

We want to find: $P(X \leq 7) = F(7)$; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

where $\mu = \ln(5) = 1.609$ and $\sigma = \sqrt{0.42} = 0.648$. Thus:

$$\begin{aligned} P(X \leq 7) &= \Phi\left(\frac{\ln 7 - 1.609}{0.648}\right) \\ &= \Phi(0.539) \\ &\approx 0.7054 \text{ (from standard normal table)} \end{aligned}$$

The result is: $P(X \leq 7) \approx 0.7054$, i.e., about 70.54% of the people will show symptoms within 7 days of exposure.

Example 3: Equipment breakdown

Example 3: Equipment breakdown

The lifetime X of a major oil platform equipment is lognormally distributed with a Median(X) = 6 months and $\sigma = 0.30$. To ensure 95% reliability, determine the desired interval x_0 for maintenance.

Example 3: Equipment breakdown

The lifetime X of a major oil platform equipment is lognormally distributed with a Median(X) = 6 months and $\sigma = 0.30$. To ensure 95% reliability, determine the desired interval x_0 for maintenance.

Given: $\mu = \ln 6 = 1.792$ and $\sigma = 0.30$, we want to find x_0 such that:

Example 3: Equipment breakdown

The lifetime X of a major oil platform equipment is lognormally distributed with a Median(X) = 6 months and $\sigma = 0.30$. To ensure 95% reliability, determine the desired interval x_0 for maintenance.

Given: $\mu = \ln 6 = 1.792$ and $\sigma = 0.30$, we want to find x_0 such that:

$$P(X > x_0) = 1 - P(X \leq x_0) =$$

Example 3: Equipment breakdown

The lifetime X of a major oil platform equipment is lognormally distributed with a Median(X) = 6 months and $\sigma = 0.30$. To ensure 95% reliability, determine the desired interval x_0 for maintenance.

Given: $\mu = \ln 6 = 1.792$ and $\sigma = 0.30$, we want to find x_0 such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$

Example 3: Equipment breakdown

The lifetime X of a major oil platform equipment is lognormally distributed with a Median(X) = 6 months and $\sigma = 0.30$. To ensure 95% reliability, determine the desired interval x_0 for maintenance.

Given: $\mu = \ln 6 = 1.792$ and $\sigma = 0.30$, we want to find x_0 such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95 \implies P(X \leq x_0) = 0.05$$

Example 3: Equipment breakdown

The lifetime X of a major oil platform equipment is lognormally distributed with a $\text{Median}(X) = 6$ months and $\sigma = 0.30$. To ensure 95% reliability, determine the desired interval x_0 for maintenance.

Given: $\mu = \ln 6 = 1.792$ and $\sigma = 0.30$, we want to find x_0 such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95 \implies P(X \leq x_0) = 0.05$$




Figure: Lognormal CDF showing the 5th percentile at $x = x_0 = 3.66$

Example 3: Equipment breakdown (cont.)

Example 3: Equipment breakdown (cont.)

Solution (using Python)

Example 3: Equipment breakdown (cont.)

Solution (using Python)

$$P(X \leq x_0) = 0.05$$

Example 3: Equipment breakdown (cont.)

Solution (using Python)

$P(X \leq x_0) = 0.05$ implies:

$$x_0 = F^{-1}(0.05)$$

Example 3: Equipment breakdown (cont.)

Solution (using Python)

$P(X \leq x_0) = 0.05$ implies:

$$x_0 = F^{-1}(0.05)$$

Code:

```
from scipy import stats
import numpy as np
x0 = stats.lognorm.ppf(q=0.05, s=0.30, scale=6)
```

Example 3: Equipment breakdown (cont.)

Solution (using Python)

$P(X \leq x_0) = 0.05$ implies:

$$x_0 = F^{-1}(0.05)$$

Code:

```
from scipy import stats
import numpy as np
x0 = stats.lognorm.ppf(q=0.05, s=0.30, scale=6)
```

Note the following about the *arguments* `stats.lognorm.ppf` function:

- The first, q , is the cumulative probability or quantile (0.05)
- The second, s , is the shape parameter σ (0.30)
- The third, `scale`, is e^{μ} (median) (6)

This returns $x_0 = 3.66$ months.

Example 3: Equipment breakdown (cont.)

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) =$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 =$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 =$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Therefore, the required inspection interval is:

$$x_0 =$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Therefore, the required inspection interval is:

$$x_0 = e^{1.297} =$$

Example 3: Equipment breakdown (cont.)

Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Therefore, the required inspection interval is:

$$x_0 = e^{1.297} = 3.66 \text{ months}$$

Modeling probabilities of elapsed times

Modeling probabilities of elapsed times

Consider the random variable X which represents the *number of arrivals* at a restaurant within a given time interval.

Modeling probabilities of elapsed times

Consider the random variable X which represents the *number of arrivals* at a restaurant within a given time interval.



Modeling probabilities of elapsed times

Consider the random variable X which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of X in t time units can be modeled by the Poisson distribution with a rate parameter λt

Modeling probabilities of elapsed times

Consider the random variable X which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of X in t time units can be modeled by the Poisson distribution with a rate parameter λt

Now consider the variable Y representing the **elapsed time** between successive arrivals.

Modeling probabilities of elapsed times

Consider the random variable X which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of X in t time units can be modeled by the Poisson distribution with a rate parameter λt

Now consider the variable Y representing the **elapsed time** between successive arrivals.

- What is the probability the time between the third and fourth arrivals is less than y minutes, for instance?

Modeling probabilities of elapsed times

Consider the random variable X which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of X in t time units can be modeled by the Poisson distribution with a rate parameter λt

Now consider the variable Y representing the **elapsed time** between successive arrivals.

- What is the probability the time between the third and fourth arrivals is less than y minutes, for instance?
- This is modeled by the **exponential distribution** with parameter λ .

PDF of exponential distribution

PDF of exponential distribution

Definition

PDF of exponential distribution

Definition

A random variable X that is exponentially distributed with parameter λ has the PDF:

PDF of exponential distribution

Definition

A random variable X that is exponentially distributed with parameter λ has the PDF:

$$f_X(x) =$$

PDF of exponential distribution

Definition

A random variable X that is exponentially distributed with parameter λ has the PDF:

$$f_X(x) = \lambda e^{-\lambda x}$$

PDF of exponential distribution

Definition

A random variable X that is exponentially distributed with parameter λ has the PDF:


$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0 \quad (12)$$

PDF of exponential distribution

Definition

A random variable X that is exponentially distributed with parameter λ has the PDF:

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0 \quad (12)$$



CDF of the exponential distribution

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) =$$

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x)$$

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$


$$F_X(x) = 1 - e^{-\lambda x}$$

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$




CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$




Note that $P(X \leq x) = 1 - e^{-\lambda x}$,

CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$



Note that $P(X \leq x) = 1 - e^{-\lambda x}$, while $P(X > x) = 1 - (1 - e^{-\lambda x}) = e^{-\lambda x}$

Mean and variance of the exponential distribution

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

$$\mathbb{E}(X)$$

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

Variance

The variance of X is given by:

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

Variance

The variance of X is given by:

$$\mathbb{V}(X)$$

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

Variance

The variance of X is given by:

$$\mathbb{V}(X) =$$

Mean and variance of the exponential distribution

Let $X \sim \text{Exponential}(\lambda)$.

Mean

The mean of X is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

Variance

The variance of X is given by:

$$\mathbb{V}(X) = \frac{1}{\lambda^2} \quad (14)$$

Example 3: Waiting for a flight

Example 3: Waiting for a flight

The delay time T of a flight is exponentially distributed with $\lambda = 2$ (delays per hour).

Example 3: Waiting for a flight

The delay time T of a flight is exponentially distributed with $\lambda = 2$ (delays per hour). Answer the following questions:

- (a) What is the mean delay (waiting) time, $\mathbb{E}(T)$?
- (b) What is the variance of the delay time $\mathbb{V}(T)$?
- (c) Find the probability that a flight will be delayed by no more than 10 minutes.
- (d) Assuming you have been waiting for a flight for an hour, what is the probability that the flight will be delayed for an additional 30 minutes? (i.e. Find $P(T > 1.5 | T > 1)$).

Example 3: Waiting for a flight (cont.)

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) =$$

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda}$$

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2}$$

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2}$$

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2} = \frac{1}{2^2} =$$

Example 3: Waiting for a flight (cont.)

Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2} = \frac{1}{2^2} = \boxed{0.25\text{hr}^2}$$

Example 3: Waiting for a flight (cont.)

Example 3: Waiting for a flight (cont.)

Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ($\frac{1}{6}$ hr) is given by:

Example 3: Waiting for a flight (cont.)

Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ($\frac{1}{6}$ hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right]$$

Example 3: Waiting for a flight (cont.)

Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ($\frac{1}{6}$ hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] =$$

Example 3: Waiting for a flight (cont.)

Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ($\frac{1}{6}$ hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] = 1 - \exp\left[-\frac{1}{3}\right]$$

Example 3: Waiting for a flight (cont.)

Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ($\frac{1}{6}$ hr) is given by:


$$\begin{aligned} P\left(T \leq \frac{1}{6}\right) &= 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] = 1 - \exp\left[-\frac{1}{3}\right] \\ &= \boxed{0.283} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ($\frac{1}{6}$ hr) is given by:

$$\begin{aligned} P\left(T \leq \frac{1}{6}\right) &= 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] = 1 - \exp\left[-\frac{1}{3}\right] \\ &= \boxed{0.283} \end{aligned}$$



Example 3: Waiting for a flight (cont.)

Solution

- (c) In python, we can use the `scipy.stats.expon.cdf` function to find $P(T \leq 1/6)$:

Example 3: Waiting for a flight (cont.)

Solution

(c) In python, we can use the `scipy.stats.expon.cdf` function to find $P(T \leq 1/6)$:

```
from scipy import stats
import numpy as np
p = stats.expon.cdf(1/6, scale=1/2)
```

where:

- $1/6$ is the value at which we want to evaluate the CDF
- $\text{scale} = 1/\lambda = 1/2$ (scale parameter/mean)

This returns $p = 0.283$, i.e., about 28.3% probability that the flight will be delayed by no more than 10 minutes.

Example 3: Waiting for a flight (cont.)

Solution

(c) In python, we can use the `scipy.stats.expon.cdf` function to find $P(T \leq 1/6)$:

```
from scipy import stats
import numpy as np
p = stats.expon.cdf(1/6, scale=1/2)
```

where:

- $1/6$ is the value at which we want to evaluate the CDF
- $\text{scale} = 1/\lambda = 1/2$ (scale parameter/mean)

This returns $p = 0.283$, i.e., about 28.3% probability that the flight will be delayed by no more than 10 minutes.

Example 3: Waiting for a flight (cont.)

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$P(T > (0.5 + 1) | T > 1) = P(T > 1.5 | T > 1)$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\ &= e^{-2(0.5)} \quad (= P(T > 0.5)) \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\ &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\ &= e^{-1} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\ &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\ &= e^{-1} = \boxed{0.37} \end{aligned}$$

Example 3: Waiting for a flight (cont.)

Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\
 &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\
 &= e^{-1} = \boxed{0.37}
 \end{aligned}$$

In Python: `p = 1 - stats.expon.pdf(.5, scale=1/2)` also returns `p = 0.37`.

Memorylessness of the exponential distribution

Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

Memoryless property

$$P(T > t + s | T > s) =$$

Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

Memoryless property

$$P(T > t + s | T > s) = P(T > t) \quad (15)$$

Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

Memoryless property

$$P(T > t + s | T > s) = P(T > t) \quad (15)$$

That is, it does not matter from which time the waiting begins (i.e. conditioning); the probability of an elapsed time remains the same.

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$
CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$
CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{\left(\mu + \frac{1}{2}\sigma^2\right)} \quad (16)$$

Variance:

$$\mathbb{V}(X) =$$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

PDF: $f_X(x) =$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF: $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

PDF: $f_X(x) = \lambda e^{-\lambda x},$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) =$$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x},$$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Mean:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (20)$$

Recap

- **Lognormal distribution:** $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:** $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Mean:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (20)$$

Variance:

$$\mathbb{V}(X) = \frac{1}{\lambda^2} \quad (21)$$