

# CEE 260/MIE 273: Probability and Statistics in Civil Engineering

## Lecture 3E: The Poisson Distribution

**Jimi Oke**

UMassAmherst  

---

College of Engineering

October 7, 2025

# Outline

## ① Introduction

## ② The Poisson distribution

## ③ Examples

## ④ Outlook

# Recap of Lecture 3b: Binomial distribution

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**
  - Fixed number of statistically independent trials  $n$

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**
  - Fixed number of statistically independent trials  $n$
  - Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**
  - Fixed number of statistically independent trials  $n$
  - Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )
  - Probability of occurrence  $p$  is constant in each trial



# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**
  - Fixed number of statistically independent trials  $n$
  - Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )
  - Probability of occurrence  $p$  is constant in each trial
- Binomial distribution  $X \sim \text{Bin}(n, p)$ :

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**
  - Fixed number of statistically independent trials  $n$
  - Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )
  - Probability of occurrence  $p$  is constant in each trial
- Binomial distribution  $X \sim \text{Bin}(n, p)$ :
  - **Mean:**  $\mathbb{E}(X) = np$

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**
  - Fixed number of statistically independent trials  $n$
  - Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )
  - Probability of occurrence  $p$  is constant in each trial
- Binomial distribution  $X \sim \text{Bin}(n, p)$ :
  - **Mean:**  $\mathbb{E}(X) = np$
  - **Variance:**  $\mathbb{V}(X) = npq = np(1 - p)$

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**

- Fixed number of statistically independent trials  $n$
- Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )
- Probability of occurrence  $p$  is constant in each trial

- Binomial distribution  $X \sim \text{Bin}(n, p)$ :

- **Mean:**  $\mathbb{E}(X) = np$
- **Variance:**  $\mathbb{V}(X) = npq = np(1 - p)$
- **PMF:**

$$P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x} \quad (1)$$

# Recap of Lecture 3b: Binomial distribution

The binomial distribution governs Bernoulli sequences.

- **Bernoulli sequence:**

- Fixed number of statistically independent trials  $n$
- Each trial has only two possible outcomes: occurrence (probability  $p$ ) or nonoccurrence (probability  $q = 1 - p$ )
- Probability of occurrence  $p$  is constant in each trial

- Binomial distribution  $X \sim \text{Bin}(n, p)$ :

- **Mean:**  $\mathbb{E}(X) = np$
- **Variance:**  $\mathbb{V}(X) = npq = np(1 - p)$
- **PMF:**

$$P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x} \quad (1)$$

- **CDF:**

$$P(X \leq x) = \sum_{k=0}^x \binom{n}{k} p^k (1 - p)^{n-k} \quad (2)$$

# Motivation

# Motivation

Consider the following situations:

# Motivation

Consider the following situations:

- Text messages arrive randomly on your cellphone at various times during the day. How might you compute the probability that a given number of texts will arrive in an hour?



# Motivation

Consider the following situations:

- Text messages arrive randomly on your cellphone at various times during the day. How might you compute the probability that a given number of texts will arrive in an hour?
- Accidents happen to occur randomly in time and space along Rte 116. What is the probability that 3 accidents will occur tomorrow?

# Motivation

Consider the following situations:

- Text messages arrive randomly on your cellphone at various times during the day. How might you compute the probability that a given number of texts will arrive in an hour?
- Accidents happen to occur randomly in time and space along Rte 116. What is the probability that 3 accidents will occur tomorrow?
- Students seem to join the Zoom meeting for our class between 12:30 and 1:15 PM in no particular order. How would you determine the probability that between 5 and 10 students will join the Zoom meeting between 1:00 and 1:01 PM?

# Motivation

Consider the following situations:

- Text messages arrive randomly on your cellphone at various times during the day. How might you compute the probability that a given number of texts will arrive in an hour?
- Accidents happen to occur randomly in time and space along Rte 116. What is the probability that 3 accidents will occur tomorrow?
- Students seem to join the Zoom meeting for our class between 12:30 and 1:15 PM in no particular order. How would you determine the probability that between 5 and 10 students will join the Zoom meeting between 1:00 and 1:01 PM?

# Motivation

Consider the following situations:

- Text messages arrive randomly on your cellphone at various times during the day. How might you compute the probability that a given number of texts will arrive in an hour?
- Accidents happen to occur randomly in time and space along Rte 116. What is the probability that 3 accidents will occur tomorrow?
- Students seem to join the Zoom meeting for our class between 12:30 and 1:15 PM in no particular order. How would you determine the probability that between 5 and 10 students will join the Zoom meeting between 1:00 and 1:01 PM?

The above are all **Poisson processes** and the respective probabilities can be modeled by the **Poisson distribution**

# Poisson process

# Poisson process

Key assumptions:

- **Randomness:**

# Poisson process

Key assumptions:

- **Randomness:** An event can occur at random and at any instant of time or any point in space
- **Independence:**

# Poisson process

Key assumptions:

- **Randomness:** An event can occur at random and at any instant of time or any point in space
- **Independence:** The occurrence of an event in a given time (or space) interval is statistically independent of that in any other non-overlapping interval
- **Constant rate:** The number of occurrences of an event in time interval  $t$  can be given by  $\lambda t$ , where  $\lambda$  is the mean rate of occurrence of the event
- **Non-overlapping events:**



# Poisson process

Key assumptions:

- **Randomness:** An event can occur at random and at any instant of time or any point in space
- **Independence:** The occurrence of an event in a given time (or space) interval is statistically independent of that in any other non-overlapping interval
- **Constant rate:** The number of occurrences of an event in time interval  $t$  can be given by  $\lambda t$ , where  $\lambda$  is the mean rate of occurrence of the event
- **Non-overlapping events:** The probability of two or more occurrences in  $t$  is negligible (i.e. only one event occurs each time).

## Further examples of Poisson processes

# Poisson process

Key assumptions:

- **Randomness:** An event can occur at random and at any instant of time or any point in space
- **Independence:** The occurrence of an event in a given time (or space) interval is statistically independent of that in any other non-overlapping interval
- **Constant rate:** The number of occurrences of an event in time interval  $t$  can be given by  $\lambda t$ , where  $\lambda$  is the mean rate of occurrence of the event
- **Non-overlapping events:** The probability of two or more occurrences in  $t$  is negligible (i.e. only one event occurs each time).

## Further examples of Poisson processes

- Breakdowns of a computer system over time

# Poisson process

Key assumptions:

- **Randomness:** An event can occur at random and at any instant of time or any point in space
- **Independence:** The occurrence of an event in a given time (or space) interval is statistically independent of that in any other non-overlapping interval
- **Constant rate:** The number of occurrences of an event in time interval  $t$  can be given by  $\lambda t$ , where  $\lambda$  is the mean rate of occurrence of the event
- **Non-overlapping events:** The probability of two or more occurrences in  $t$  is negligible (i.e. only one event occurs each time).

## Further examples of Poisson processes

- Breakdowns of a computer system over time
- Number of accidents in an industrial facility over time

# Poisson process

Key assumptions:

- **Randomness:** An event can occur at random and at any instant of time or any point in space
- **Independence:** The occurrence of an event in a given time (or space) interval is statistically independent of that in any other non-overlapping interval
- **Constant rate:** The number of occurrences of an event in time interval  $t$  can be given by  $\lambda t$ , where  $\lambda$  is the mean rate of occurrence of the event
- **Non-overlapping events:** The probability of two or more occurrences in  $t$  is negligible (i.e. only one event occurs each time).

## Further examples of Poisson processes

- Breakdowns of a computer system over time
- Number of accidents in an industrial facility over time
- Customer arrivals at a bike store in a given morning

# Poisson distribution: PMF and CDF

# Poisson distribution: PMF and CDF

If  $X$  has a Poisson distribution with rate parameter  $\lambda$ , i.e.  $X \sim \text{Poisson}(\lambda)$ , its PMF is:

# Poisson distribution: PMF and CDF

If  $X$  has a Poisson distribution with rate parameter  $\lambda$ , i.e.  $X \sim \text{Poisson}(\lambda)$ , its PMF is:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x \geq 0 \quad (3)$$

# Poisson distribution: PMF and CDF

If  $X$  has a Poisson distribution with rate parameter  $\lambda$ , i.e.  $X \sim \text{Poisson}(\lambda)$ , its PMF is:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x \geq 0 \quad (3)$$

where:



# Poisson distribution: PMF and CDF

If  $X$  has a Poisson distribution with rate parameter  $\lambda$ , i.e.  $X \sim \text{Poisson}(\lambda)$ , its PMF is:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x \geq 0 \quad (3)$$

where:  $\lambda \equiv$  expected/mean number of occurrences in unit time interval.

# Poisson distribution: PMF and CDF

If  $X$  has a Poisson distribution with rate parameter  $\lambda$ , i.e.  $X \sim \text{Poisson}(\lambda)$ , its PMF is:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x \geq 0 \quad (3)$$

where:  $\lambda \equiv$  expected/mean number of occurrences in unit time interval.

The CDF of  $X$  is given by:

# Poisson distribution: PMF and CDF

If  $X$  has a Poisson distribution with rate parameter  $\lambda$ , i.e.  $X \sim \text{Poisson}(\lambda)$ , its PMF is:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x \geq 0 \quad (3)$$

where:  $\lambda \equiv$  expected/mean number of occurrences in unit time interval.

The CDF of  $X$  is given by:


$$P(X \leq x) = \sum_{k=0}^x \frac{\lambda^k}{k!} e^{-\lambda} \quad (4)$$

# Poisson distribution: PMF

# Poisson distribution: PMF

PMF of  $X \sim \text{Poisson}(\lambda)$


$$P(X = x)$$



# Poisson distribution: PMF

PMF of  $X \sim \text{Poisson}(\lambda)$


$$P(X = x)$$



# Poisson distribution: PMF

PMF of  $X \sim \text{Poisson}(\lambda)$


$$P(X = x)$$



# Poisson distribution: PMF

PMF of  $X \sim \text{Poisson}(\lambda)$

$$P(X = x)$$






# Poisson distribution: PMF


PMF of  $X \sim \text{Poisson}(\lambda)$

$$P(X = x)$$



# Poisson distribution: CDF

# Poisson distribution: CDF



# Mean and variance

# Mean and variance

The mean  $\mathbb{E}(X)$  and variance  $\mathbb{V}(X)$  of a Poisson distributed random variable are equivalent

# Mean and variance

The mean  $\mathbb{E}(X)$  and variance  $\mathbb{V}(X)$  of a Poisson distributed random variable are equivalent

$$\mathbb{E}(X) = \mathbb{V}(X) = \sigma_X^2 =$$

# Mean and variance

The mean  $\mathbb{E}(X)$  and variance  $\mathbb{V}(X)$  of a Poisson distributed random variable are equivalent

$$\mathbb{E}(X) = \mathbb{V}(X) = \sigma_X^2 = \lambda \quad (5)$$

# Mean and variance

The mean  $\mathbb{E}(X)$  and variance  $\mathbb{V}(X)$  of a Poisson distributed random variable are equivalent

$$\mathbb{E}(X) = \mathbb{V}(X) = \sigma_X^2 = \lambda \quad (5)$$

The standard deviation is



# Mean and variance

The mean  $\mathbb{E}(X)$  and variance  $\mathbb{V}(X)$  of a Poisson distributed random variable are equivalent

$$\mathbb{E}(X) = \mathbb{V}(X) = \sigma_X^2 = \lambda \quad (5)$$

The standard deviation is  $\sigma(X) = \sqrt{\mathbb{V}(X)} = \sqrt{\lambda}$

# Computing Poisson probabilities in Python

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`
- Similarly, you can find the CDF  $P(X \leq x)$  as `poisson.cdf(x, lambda)`

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`
- Similarly, you can find the CDF  $P(X \leq x)$  as `poisson.cdf(x, lambda)`
- Note that  $P(X > x)$  (survival function) is obtained by

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`
- Similarly, you can find the CDF  $P(X \leq x)$  as `poisson.cdf(x, lambda)`
- Note that  $P(X > x)$  (survival function) is obtained by

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`
- Similarly, you can find the CDF  $P(X \leq x)$  as `poisson.cdf(x, lambda)`
- Note that  $P(X > x)$  (survival function) is obtained by  
`1 - poisson.cdf(x, lambda)`



# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`
- Similarly, you can find the CDF  $P(X \leq x)$  as `poisson.cdf(x, lambda)`
- Note that  $P(X > x)$  (survival function) is obtained by  
`1 - poisson.cdf(x, lambda)`  
or

# Computing Poisson probabilities in Python

- Load: `from scipy.stats import poisson`
- Compute the PMF  $P(X = x)$  using `poisson.pmf(x, lambda)`
- Similarly, you can find the CDF  $P(X \leq x)$  as `poisson.cdf(x, lambda)`
- Note that  $P(X > x)$  (survival function) is obtained by  
`1 - poisson.cdf(x, lambda)`  
or  
`poisson.sf(x, lambda) #sf = survival function = 1 - cdf`

# Example 1: Amherst Coffee customer arrivals

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM;

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM;

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution



## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30}$$

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30)$$

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30) = \boxed{0.0139}$$

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30) = \boxed{0.0139}$$

(b) The desired probability is given by:

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30) = \boxed{0.0139}$$

(b) The desired probability is given by:

$$P(X \leq 40) =$$



## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30) = \boxed{0.0139}$$

(b) The desired probability is given by:

$$P(X \leq 40) = \sum_{k=0}^{40} \frac{30^k}{k!} e^{-30}$$

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30) = \boxed{0.0139}$$

(b) The desired probability is given by:

$$P(X \leq 40) = \sum_{k=0}^{40} \frac{30^k}{k!} e^{-30} = \text{poisson.cdf}(40, 30)$$

## Example 1: Amherst Coffee customer arrivals

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Find the (a) probability that **exactly 40** customers arrive between 10 and 11 AM; (b) probability that **no more than 40** customers arrive between 11 AM and 12 PM; (c) **uncertainty** in the rate parameter.

### Solution

The number of customers arriving within an hour  $X \sim \text{Poisson}(\lambda = 30)$ .

(a) The desired probability is given by:

$$P(X = 40) = \frac{30^{40}}{40!} e^{-30} = \text{poisson.pmf}(40, 30) = \boxed{0.0139}$$

(b) The desired probability is given by:

$$P(X \leq 40) = \sum_{k=0}^{40} \frac{30^k}{k!} e^{-30} = \text{poisson.cdf}(40, 30) = \boxed{0.9677}$$

# Example 1: Amherst Coffee customer arrivals (cont.)

# Example 1: Amherst Coffee customer arrivals (cont.)

## Solution (cont.)

# Example 1: Amherst Coffee customer arrivals (cont.)

## Solution (cont.)

(c) The uncertainty is given by the standard deviation:

# Example 1: Amherst Coffee customer arrivals (cont.)

## Solution (cont.)

(c) The uncertainty is given by the standard deviation:

$$\sigma_X = \sqrt{\lambda}$$

# Example 1: Amherst Coffee customer arrivals (cont.)

## Solution (cont.)

(c) The uncertainty is given by the standard deviation:

$$\sigma_X = \sqrt{\lambda} = \sqrt{30}$$



# Example 1: Amherst Coffee customer arrivals (cont.)

## Solution (cont.)

(c) The uncertainty is given by the standard deviation:

$$\sigma_X = \sqrt{\lambda} = \sqrt{30} = \boxed{5.477}$$

## Example 2: Aircraft arrivals at airport

## Example 2: Aircraft arrivals at airport

Suppose small aircraft arrive at a certain airport according to a Poisson process with a rate  $\lambda = 8/\text{hr}$ .

## Example 2: Aircraft arrivals at airport

Suppose small aircraft arrive at a certain airport according to a Poisson process with a rate  $\lambda = 8/\text{hr}$ .

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?
- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?
- (c) How many aircraft do you expect to arrive during a 90-minute period?

## Example 2: Aircraft arrivals at airport (cont.)

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} =$$

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} =$$



## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8)$$

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

### Solution

The desired probability is given by:

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

### Solution

The desired probability is given by:

$$P(X_{1hr} > 9) = 1 - P(X_{1hr} \leq 9)$$

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

### Solution

The desired probability is given by:

$$\begin{aligned} P(X_{1hr} > 9) &= 1 - P(X_{1hr} \leq 9) \\ &= 1 - \sum_{x=0}^9 \frac{\lambda^x e^{-\lambda}}{x!} = \end{aligned}$$

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

### Solution

The desired probability is given by:

$$\begin{aligned} P(X_{1hr} > 9) &= 1 - P(X_{1hr} \leq 9) \\ &= 1 - \sum_{x=0}^9 \frac{\lambda^x e^{-\lambda}}{x!} = 1 - \text{poisson.cdf}(9, 8) \end{aligned}$$

## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

### Solution

The desired probability is given by:

$$\begin{aligned} P(X_{1hr} > 9) &= 1 - P(X_{1hr} \leq 9) \\ &= 1 - \sum_{x=0}^9 \frac{\lambda^x e^{-\lambda}}{x!} = 1 - \text{poisson.cdf}(9, 8) \\ &= 1 - 0.7166 = \end{aligned}$$



## Example 2: Aircraft arrivals at airport (cont.)

- (a) What is the probability that exactly 6 small aircraft arrive during a 1-hr period?

### Solution

$$P(X_{1hr} = 6) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{8^6 e^{-8}}{6!} = \text{poisson.pmf}(6, 8) = \boxed{0.1221}$$

- (b) What is the probability that at least 10 small aircraft arrive during a 1-hr period?

### Solution

The desired probability is given by:

$$\begin{aligned} P(X_{1hr} > 9) &= 1 - P(X_{1hr} \leq 9) \\ &= 1 - \sum_{x=0}^9 \frac{\lambda^x e^{-\lambda}}{x!} = 1 - \text{poisson.cdf}(9, 8) \\ &= 1 - 0.7166 = \boxed{0.2834} \end{aligned}$$

# Manipulating the rate parameter

# Manipulating the rate parameter

- The rate parameter  $\lambda$  is given by the mean number of occurrences per unit time.

# Manipulating the rate parameter

- The rate parameter  $\lambda$  is given by the mean number of occurrences per unit time.
- If we are to find the probability of a certain number of events within a different time interval  $T'$ , where

$$T' = T \times t \tag{6}$$

# Manipulating the rate parameter

- The rate parameter  $\lambda$  is given by the mean number of occurrences per unit time.
- If we are to find the probability of a certain number of events within a different time interval  $T'$ , where

$$T' = T \times t \tag{6}$$

# Manipulating the rate parameter

- The rate parameter  $\lambda$  is given by the mean number of occurrences per unit time.
- If we are to find the probability of a certain number of events within a different time interval  $T'$ , where

$$T' = T \times t \quad (6)$$

- Then, we can find the new rate parameter by multiplying  $\lambda$  by the appropriate number of time units  $t$ , such that

$$\lambda' = \lambda t \quad (7)$$

# Manipulating the rate parameter

- The rate parameter  $\lambda$  is given by the mean number of occurrences per unit time.
- If we are to find the probability of a certain number of events within a different time interval  $T'$ , where

$$T' = T \times t \quad (6)$$

- Then, we can find the new rate parameter by multiplying  $\lambda$  by the appropriate number of time units  $t$ , such that

$$\lambda' = \lambda t \quad (7)$$

# Manipulating the rate parameter

- The rate parameter  $\lambda$  is given by the mean number of occurrences per unit time.
- If we are to find the probability of a certain number of events within a different time interval  $T'$ , where

$$T' = T \times t \quad (6)$$

- Then, we can find the new rate parameter by multiplying  $\lambda$  by the appropriate number of time units  $t$ , such that

$$\lambda' = \lambda t \quad (7)$$

- And thus:

$$P(X_t = x) = \frac{(\lambda t)^x}{x!} e^{-(\lambda t)} = \frac{(\lambda')^x}{x!} e^{-(\lambda')} \quad (8)$$



## Example 2: Aircraft arrivals at airport (cont.)

## Example 2: Aircraft arrivals at airport (cont.)

(c) How many aircraft do you expect to arrive during a 90-minute period?

## Example 2: Aircraft arrivals at airport (cont.)

(c) How many aircraft do you expect to arrive during a 90-minute period?

**Solution**

## Example 2: Aircraft arrivals at airport (cont.)

(c) How many aircraft do you expect to arrive during a 90-minute period?

### Solution

Expected arrivals within 90 minutes =  $8/\text{hr} \times 1.5 \text{ hrs} =$

## Example 2: Aircraft arrivals at airport (cont.)

(c) How many aircraft do you expect to arrive during a 90-minute period?

### Solution

Expected arrivals within 90 minutes =  $8/\text{hr} \times 1.5 \text{ hrs} = 12$ .

## Example 3: Amherst Coffee customer arrivals (revisited)

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution



## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long.

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* =$$

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2$$

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

The desired probability is thus given by:

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

The desired probability is thus given by:

$$P(X_{2hr} > 65) =$$

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

The desired probability is thus given by:

$$P(X_{2hr} > 65) = 1 - P(X_{2hr} \leq 65)$$



## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

The desired probability is thus given by:

$$P(X_{2hr} > 65) = 1 - P(X_{2hr} \leq 65) = 1 - \text{poisson.cdf}(65, 60)$$

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

The desired probability is thus given by:

$$\begin{aligned} P(X_{2hr} > 65) &= 1 - P(X_{2hr} \leq 65) = 1 - \text{poisson.cdf}(65, 60) \\ &\equiv \text{poisson.sf}(65, 60) \end{aligned}$$

## Example 3: Amherst Coffee customer arrivals (revisited)

Starting at 7 AM, customers arrive Amherst Coffee according to a Poisson process at the rate of 30 customers per hour. Now find the probability more than 65 customers arrive between 10 and 12 PM.

### Solution

The probability is now desired for an interval twice as long. So we compute a new rate parameter for 2-hr interval:

$$\lambda^* = 30 \times 2 = 60 \text{ (per 2 hrs)}$$

The desired probability is thus given by:

$$\begin{aligned} P(X_{2hr} > 65) &= 1 - P(X_{2hr} \leq 65) = 1 - \text{poisson.cdf}(65, 60) \\ &\equiv \text{poisson.sf}(65, 60) = \boxed{0.2355} \end{aligned}$$

## Example 4: Text messages

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.



## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.

Thus:

$$P(X_{12} = 18 \cap X_5 = 52) =$$

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.

Thus:

$$P(X_{12} = 18 \cap X_5 = 52) = P(X_{12} = 18) \times P(X_5 = 52)$$

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.

Thus:

$$\begin{aligned} P(X_{12} = 18 \cap X_5 = 52) &= P(X_{12} = 18) \times P(X_5 = 52) \\ &= \frac{(10(2))^{18}}{18!} e^{-10(2)} \times \frac{(10(5))^{52}}{52!} e^{-10(5)} \end{aligned}$$

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.

Thus:

$$\begin{aligned} P(X_{12} = 18 \cap X_5 = 52) &= P(X_{12} = 18) \times P(X_5 = 52) \\ &= \frac{(10(2))^{18}}{18!} e^{-10(2)} \times \frac{(10(5))^{52}}{52!} e^{-10(5)} \\ &= \text{poisson.pmf}(18, 20) * \text{poisson.pmf}(52, 50) \end{aligned}$$

## Example 4: Text messages

A student receives text messages starting at 10 AM at the rate of 10 texts per hour according to a Poisson process. Find the probability that they will receive exactly 18 texts by noon and 70 texts by 5 PM.

### Solution

- The student receives 18 texts in the first 2 hrs and then  $70 - 18 = 52$  texts in the next 5 hrs (5:00 PM).
- The events  $X_{12} = 18$  and  $X_5 = 52$  are independent within the specified time intervals.

Thus:

$$\begin{aligned} P(X_{12} = 18 \cap X_5 = 52) &= P(X_{12} = 18) \times P(X_5 = 52) \\ &= \frac{(10(2))^{18}}{18!} e^{-10(2)} \times \frac{(10(5))^{52}}{52!} e^{-10(5)} \\ &= \text{poisson.pmf}(18,20) * \text{poisson.pmf}(52,50) \\ &= \boxed{0.0045} \end{aligned}$$

# Poisson distribution as limit of binomial distribution

# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials,



# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$


where  $\lambda = np$ .

# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .






# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .




# Poisson distribution as limit of binomial distribution

For a binomially distributed r.v.  $X$  with  $n$  trials, as  $n \rightarrow \infty$  and  $p \rightarrow 0$ , then

$$X \sim \text{Bin}(n, p) \rightarrow X \sim \text{Poisson}(\lambda) \quad (9)$$

where  $\lambda = np$ .



# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)

# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)
- Such events are described as **Poisson processes**



# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)
- Such events are described as **Poisson processes**
- The **PMF** of a Poisson random variable with rate parameter  $\lambda$  is given by:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad (10)$$

where  $x \geq 0$

# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)
- Such events are described as **Poisson processes**
- The **PMF** of a Poisson random variable with rate parameter  $\lambda$  is given by:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad (10)$$

where  $x \geq 0$

- The **mean** and **variance** of a Poisson random variable are equal:

$$\mathbb{E}(X) = \mathbb{V}(X) = \lambda \quad (11)$$

# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)
- Such events are described as **Poisson processes**
- The **PMF** of a Poisson random variable with rate parameter  $\lambda$  is given by:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad (10)$$

where  $x \geq 0$

- The **mean** and **variance** of a Poisson random variable are equal:

$$\mathbb{E}(X) = \mathbb{V}(X) = \lambda \quad (11)$$

- Recall that the **standard deviation** is simply the square root of the variance:

$$\sigma(X) = \sqrt{\lambda} \quad (12)$$

# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)
- Such events are described as **Poisson processes**
- The **PMF** of a Poisson random variable with rate parameter  $\lambda$  is given by:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad (10)$$

where  $x \geq 0$

- The **mean** and **variance** of a Poisson random variable are equal:

$$\mathbb{E}(X) = \mathbb{V}(X) = \lambda \quad (11)$$

- Recall that the **standard deviation** is simply the square root of the variance:

$$\sigma(X) = \sqrt{\lambda} \quad (12)$$

# Recap

- The Poisson distribution is used to model the probability that a number of independent events occur within a fixed time interval (or within a finite space)
- Such events are described as **Poisson processes**
- The **PMF** of a Poisson random variable with rate parameter  $\lambda$  is given by:

$$P(X = x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad (10)$$

where  $x \geq 0$

- The **mean** and **variance** of a Poisson random variable are equal:

$$\mathbb{E}(X) = \mathbb{V}(X) = \lambda \quad (11)$$

- Recall that the **standard deviation** is simply the square root of the variance:

$$\sigma(X) = \sqrt{\lambda} \quad (12)$$

## Reading

- Open Intro Statistics Section 4.5