

# CEE 260/MIE 273: Probability and Statistics in Civil Engineering

## Lecture M3c: Lognormal and Exponential Distributions

**Jimi Oke**

UMassAmherst

---

College of Engineering

September 30, 2025

# Outline

- ① Introduction
- ② Lognormal distribution
- ③ Exponential distribution
- ④ Outlook

# Recap of normal distribution

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its **"Z-score"**:

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \tag{1}$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its **“Z-score”**:

$$Z = \frac{X - \mu}{\sigma} \tag{2}$$



# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \quad (1)$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \quad (2)$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \tag{1}$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its **"Z-score"**:

$$Z = \frac{X - \mu}{\sigma} \tag{2}$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol  $\Phi$  ("phi") is used to represent the CDF of the **standard normal distribution**, whose values can be looked up in a table.

# Recap of normal distribution

- The **PDF** of the normal distribution (parameters: mean  $\mu$  and variance  $\sigma^2$ ) is given by

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right] \tag{1}$$

- There is no closed-form solution to the integral of the normal CDF
- Instead, it is customary to standardize a normal variable to its “**Z-score**”:

$$Z = \frac{X - \mu}{\sigma} \tag{2}$$

- The mean and variance of the standard normal distribution are 0 and 1, respectively.
- The symbol  $\Phi$  (“phi”) is used to represent the CDF of the *standard normal distribution*, whose values can be looked up in a table.
- In Python, the `norm.cdf(x, mu, sigma)` and `norm.ppf(p, mu, sigma)` can be used to compute probabilities and inverse CDFs of the normal distribution, respectively.

# Using the standard normal CDF probability table

# Using the standard normal CDF probability table


- First convert the random variable to its  $Z$ -score

# Using the standard normal CDF probability table

- First convert the random variable to its  $Z$ -score
- Find the corresponding value in the table

# Using the standard normal CDF probability table

- First convert the random variable to its  $Z$ -score
- Find the corresponding value in the table



# Objectives of today's lecture



# Objectives of today's lecture

- Lognormal distribution

# Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions

# Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions
- Exponential distribution

# Objectives of today's lecture

- Lognormal distribution
- Relationship between lognormal and normal distributions
- Exponential distribution
- Memoryless property of exponential distribution

# PDF of the lognormal distribution

## Definition

# PDF of the lognormal distribution

## Definition

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma)$ ) has the PDF:

# PDF of the lognormal distribution

## Definition

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma)$ ) has the PDF:


$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$

# PDF of the lognormal distribution

## Definition

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma)$ ) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$






# PDF of the lognormal distribution

## Definition

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma)$ ) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$




# PDF of the lognormal distribution

## Definition

A random variable  $X$  that is lognormally distributed with the parameters  $\mu$  and  $\sigma$  (denoted  $X \sim \mathcal{LN}(\mu, \sigma)$ ) has the PDF:

$$f_X(x) = \frac{1}{(\sigma x)\sqrt{2\pi}} \exp \left[ -\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2 \right] \quad x \geq 0 \quad (3)$$



# Mean, median and variance of a lognormal distribution

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of  $X$  is given by

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of  $X$  is:

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$



# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :

$$X \sim \mathcal{LN}(\mu, \sigma^2)$$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of  $X$  is given by:

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :  
 $X \sim \mathcal{LN}(\mu, \sigma^2)$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \tag{4}$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \tag{5}$$

- The **variance** of  $X$  is given by:

$$\mathbb{V}(X) =$$

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :  
 $X \sim \mathcal{LN}(\mu, \sigma^2)$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of  $X$  is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :  
 $X \sim \mathcal{LN}(\mu, \sigma^2)$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of  $X$  is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :  
 $X \sim \mathcal{LN}(\mu, \sigma^2)$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \quad (4)$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \quad (5)$$

- The **variance** of  $X$  is given by:

$$\mathbb{V}(X) = (\exp[\sigma^2] - 1) \exp[2\mu + \sigma^2] \quad (6)$$

## Notes

- $\mu$  and  $\sigma$  are the mean and standard deviation of the associated normal distribution of  $\ln(X)$ .

# Mean, median and variance of a lognormal distribution

Let  $X$  be a lognormally distributed random variable with parameters  $\mu$  and  $\sigma$ :  
 $X \sim \mathcal{LN}(\mu, \sigma^2)$

- The **mean** of  $X$  is given by

$$\mathbb{E}(X) = \exp \left[ \mu + \frac{1}{2} \sigma^2 \right] \tag{4}$$

- The **median** of  $X$  is:

$$\text{Median}(X) = \exp(\mu) \tag{5}$$

- The **variance** of  $X$  is given by:

$$\mathbb{V}(X) = (\exp [\sigma^2] - 1) \exp [2\mu + \sigma^2] \tag{6}$$

## Notes

- $\mu$  and  $\sigma$  are the mean and standard deviation of the associated normal distribution of  $\ln(X)$ .
- Thus, if  $X \sim \mathcal{LN}(\mu, \sigma)$ , then  $\ln(X) \sim \mathcal{N}(\mu, \sigma)$

# Example 1: Mean and variance of lognormal distribution (1)

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?



# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) =$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$\therefore$  The mean is given by

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2}$$



# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\therefore \text{The mean is given by } \mathbb{E}(X) = e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\begin{aligned} \therefore \text{The mean is given by } \mathbb{E}(X) &= e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21} \\ &= 5(e^{0.21}) \end{aligned}$$

# Example 1: Mean and variance of lognormal distribution (1)

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What are the mean and variance of its distribution?

## Solution

First, we find the parameter  $\mu$ :

$$\text{Median}(X) = e^{\mu}$$

$$5 = e^{\mu}$$

$$\implies \ln(5) = \mu$$

$$\begin{aligned} \therefore \text{The mean is given by } \mathbb{E}(X) &= e^{\mu + \frac{1}{2}\sigma^2} = e^{\ln(5) + 0.21} = e^{\ln(5)} \times e^{0.21} \\ &= 5(e^{0.21}) = \boxed{6.17 \text{ days}} \end{aligned}$$

# Example 1: Mean and variance of lognormal distribution (2)

# Example 1: Mean and variance of lognormal distribution (2)

Solution (cont.)

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

$$\mathbb{V}(X) = (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2])$$



# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

$$\begin{aligned}\mathbb{V}(X) &= (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2]) \\ &= (\exp(0.42) - 1)(\exp(2 \ln(5) + 0.42))\end{aligned}$$

# Example 1: Mean and variance of lognormal distribution (2)

## Solution (cont.)

The variance is given by:

$$\begin{aligned}
 \mathbb{V}(X) &= (\exp(\sigma^2) - 1) (\exp[2\mu + \sigma^2]) \\
 &= (\exp(0.42) - 1)(\exp(2 \ln(5) + 0.42)) \\
 &= \boxed{19.86 \text{ days}^2}
 \end{aligned}$$

# Relationship between normal and lognormal distributions

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma)$$

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies$$

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$



# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma$ , then:
  - $\mu = \mathbb{E}(\ln(X))$  and  $\sigma^2 = \mathbb{V}(\ln(X))$

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma$ , then:
  - $\mu = \mathbb{E}(\ln(X))$  and  $\sigma^2 = \mathbb{V}(\ln(X))$
  - $\ln(X)$  is **normally** distributed with the same parameters.

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:

$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \tag{7}$$

- Conversely, if a random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma$ , then:
  - $\mu = \mathbb{E}(\ln(X))$  and  $\sigma^2 = \mathbb{V}(\ln(X))$
  - $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma)$$

# Relationship between normal and lognormal distributions

- If random variable  $X$  is **normally** distributed with the parameters  $\mu$  and  $\sigma^2$ , then:
  - $\mu = \mathbb{E}(X)$  and  $\sigma^2 = \mathbb{V}(X)$
  - $e^X$  is **lognormally** distributed with the same parameters:


$$X \sim \mathcal{N}(\mu, \sigma) \implies e^X \sim \mathcal{LN}(\mu, \sigma) \quad (7)$$

- Conversely, if a random variable  $X$  is **lognormally** distributed with the **parameters**  $\mu$  and  $\sigma$ , then:
  - $\mu = \mathbb{E}(\ln(X))$  and  $\sigma^2 = \mathbb{V}(\ln(X))$
  - $\ln(X)$  is **normally** distributed with the same parameters.

$$X \sim \mathcal{LN}(\mu, \sigma) \implies \ln(X) \sim \mathcal{N}(\mu, \sigma) \quad (8)$$

# Relationship between normal and lognormal (cont.)

# Relationship between normal and lognormal (cont.)



# Positive skewness of lognormal distribution



# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed

# Positive skewness of lognormal distribution


- The lognormal distribution is positively skewed
- Its mean is always greater than its median

# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median

# Positive skewness of lognormal distribution

- The lognormal distribution is positively skewed
- Its mean is always greater than its median



# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) =$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma_X \sqrt{2\pi}}$$

# Probability of a lognormal random variate


Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma X \sqrt{2\pi}} \int_a^b \exp \left[ -\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2 \right] dx \quad (9)$$

# Probability of a lognormal random variate

Given a r.v.  $X$  that is lognormally distributed with parameters  $\mu$  and  $\sigma^2$ :

$$P(a < X \leq b) = \frac{1}{\sigma x \sqrt{2\pi}} \int_a^b \exp \left[ -\frac{1}{2} \left( \frac{\ln(x) - \mu}{\sigma} \right)^2 \right] dx \quad (9)$$



**Figure:** Lognormal distribution with  $\mu = 0$ ,  $\sigma = 1$ , showing  $P(a < X < b)$  where  $a = 2$  and  $b = 4$



# Probability of a lognormal random variate (cont.)

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma}$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b)$$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma}$$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x)-\mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a)-\mu)/\sigma}^{(\ln(b)-\mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \tag{10}$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:



Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x)-\mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a)-\mu)/\sigma}^{(\ln(b)-\mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \tag{10}$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b)$$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right)$$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x) - \mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a) - \mu)/\sigma}^{(\ln(b) - \mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \quad (10)$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right) -$$

# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x)-\mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a)-\mu)/\sigma}^{(\ln(b)-\mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \tag{10}$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right) - \Phi\left(\frac{\ln a - \mu}{\sigma}\right) \tag{11}$$


# Probability of a lognormal random variate (cont.)

Substituting  $z = \frac{\ln(x)-\mu}{\sigma} \implies dx = \sigma x dz$ , we obtain:

$$P(a < X \leq b) = \frac{1}{\sqrt{2\pi}} \int_{(\ln(a)-\mu)/\sigma}^{(\ln(b)-\mu)/\sigma} \exp\left[-\frac{1}{2}z^2\right] dz \tag{10}$$

Recognizing that the integrand is the PDF of the **standard normal distribution**, we have:

$$P(a < X \leq b) = \Phi\left(\frac{\ln b - \mu}{\sigma}\right) - \Phi\left(\frac{\ln a - \mu}{\sigma}\right) \tag{11}$$



## Example 2: Equipment breakdown

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:



## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) =$$

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:


$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$

## Example 2: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$



**Figure:** Standard normal distribution showing  $P(\ln(2) < Z < \ln(4))$


## Example 2: Probability of incubation period

## Example 2: Probability of incubation period

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What is the probability that a randomly selected person will show symptoms within 7 days of exposure (i.e.,  $P(X \leq 7)$ )?

## Example 2: Probability of incubation period

The incubation period of the COVID-19 infection is assumed to be lognormally distributed with a median of about 5 days and  $\sigma^2 = 0.42$ . What is the probability that a randomly selected person will show symptoms within 7 days of exposure (i.e.,  $P(X \leq 7)$ )?



**Figure:** Lognormal distribution with  $\mu = \ln(5)$ ,  $\sigma^2 = 0.42$ , showing  $P(X \leq 7)$

## Example 2: Probability of incubation period (cont.)

## Example 2: Probability of incubation period (cont.)

### Solution (using Python)



## Example 2: Probability of incubation period (cont.)

### Solution (using Python)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the `scipy.stats.lognorm.cdf` function:

## Example 2: Probability of incubation period (cont.)

### Solution (using Python)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the `scipy.stats.lognorm.cdf` function:

```
from scipy import stats
import numpy as np
p = stats.lognorm.cdf(7, s=0.648, scale=5)
```

## Example 2: Probability of incubation period (cont.)

### Solution (using Python)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the `scipy.stats.lognorm.cdf` function:

```
from scipy import stats
import numpy as np
p = stats.lognorm.cdf(7, s=0.648, scale=5)
```

where:

- $s = \sigma = 0.648 = \sqrt{0.42}$  (shape parameter/standard deviation of underlying normal)
- $\text{scale} = \exp(\mu) = \exp(\ln(5)) = 5$  (scale parameter/median)

The result is:  $p = 0.691$ , i.e., about 69.1% of the people will show symptoms within 7 days of exposure.

## Example 2: Probability of incubation period (cont.)

## Example 2: Probability of incubation period (cont.)

Solution (using tables)

## Example 2: Probability of incubation period (cont.)

### Solution (using tables)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the relationship:

## Example 2: Probability of incubation period (cont.)

### Solution (using tables)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

## Example 2: Probability of incubation period (cont.)

### Solution (using tables)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

where  $\mu = \ln(5) = 1.609$  and  $\sigma = \sqrt{0.42} = 0.648$ . Thus:



## Example 2: Probability of incubation period (cont.)

### Solution (using tables)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

where  $\mu = \ln(5) = 1.609$  and  $\sigma = \sqrt{0.42} = 0.648$ . Thus:

$$\begin{aligned} P(X \leq 7) &= \Phi\left(\frac{\ln 7 - 1.609}{0.648}\right) \\ &= \Phi(0.539) \\ &\approx 0.7054 \text{ (from standard normal table)} \end{aligned}$$

## Example 2: Probability of incubation period (cont.)

### Solution (using tables)

We want to find:  $P(X \leq 7) = F(7)$ ; thus we use the relationship:

$$P(X \leq x) = \Phi\left(\frac{\ln x - \mu}{\sigma}\right)$$

where  $\mu = \ln(5) = 1.609$  and  $\sigma = \sqrt{0.42} = 0.648$ . Thus:

$$\begin{aligned} P(X \leq 7) &= \Phi\left(\frac{\ln 7 - 1.609}{0.648}\right) \\ &= \Phi(0.539) \\ &\approx 0.7054 \text{ (from standard normal table)} \end{aligned}$$

The result is:  $P(X \leq 7) \approx 0.7054$ , i.e., about 70.54% of the people will show symptoms within 7 days of exposure.

## Example 3: Equipment breakdown

## Example 3: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

## Example 3: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

## Example 3: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) =$$

# Example 3: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95$$

## Example 3: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a Median( $X$ ) = 6 months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95 \implies P(X \leq x_0) = 0.05$$




## Example 3: Equipment breakdown

The lifetime  $X$  of a major oil platform equipment is lognormally distributed with a  $\text{Median}(X) = 6$  months and  $\sigma = 0.30$ . To ensure 95% reliability, determine the desired interval  $x_0$  for maintenance.

Given:  $\mu = \ln 6 = 1.792$  and  $\sigma = 0.30$ , we want to find  $x_0$  such that:

$$P(X > x_0) = 1 - P(X \leq x_0) = 0.95 \implies P(X \leq x_0) = 0.05$$



**Figure:** Lognormal CDF showing the 5th percentile at  $x = x_0 = 3.66$

## Example 3: Equipment breakdown (cont.)

## Example 3: Equipment breakdown (cont.)

### Solution (using Python)

## Example 3: Equipment breakdown (cont.)

### Solution (using Python)

$$P(X \leq x_0) = 0.05$$

## Example 3: Equipment breakdown (cont.)

### Solution (using Python)

$P(X \leq x_0) = 0.05$  implies:

$$x_0 = F^{-1}(0.05)$$

## Example 3: Equipment breakdown (cont.)

### Solution (using Python)

$P(X \leq x_0) = 0.05$  implies:

$$x_0 = F^{-1}(0.05)$$

Code:

---

```
from scipy import stats
import numpy as np
x0 = stats.lognorm.ppf(q=0.05, s=0.30, scale=6)
```

---

## Example 3: Equipment breakdown (cont.)

### Solution (using Python)

$P(X \leq x_0) = 0.05$  implies:

$$x_0 = F^{-1}(0.05)$$

Code:

---

```
from scipy import stats
import numpy as np
x0 = stats.lognorm.ppf(q=0.05, s=0.30, scale=6)
```

---

Note the following about the *arguments* `stats.lognorm.ppf` function:

- The first, `q`, is the cumulative probability or quantile (0.05)
- The second, `s`, is the shape parameter  $\sigma$  (0.30)
- The third, `scale`, is  $e^{\mu}$  (median) (6)

This returns  $x_0 = 3.66$  months.

## Example 3: Equipment breakdown (cont.)



## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) =$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 =$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 =$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$



## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Therefore, the required inspection interval is:

$$x_0 =$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Therefore, the required inspection interval is:

$$x_0 = e^{1.297} =$$

## Example 3: Equipment breakdown (cont.)

### Solution (using tables)

Thus:

$$\Phi\left(\frac{\ln(x_0) - 1.792}{0.30}\right) = 0.05$$

$$\frac{\ln x_0 - 1.792}{0.30} = \Phi^{-1}(0.05)$$

$$\ln x_0 - 1.792 = 0.30[-\Phi^{-1}(0.95)]$$

$$\ln x_0 = 1.792 + 0.30(-1.65)$$

$$\ln x_0 = 1.792 - 0.495 = 1.297$$

Therefore, the required inspection interval is:

$$x_0 = e^{1.297} = 3.66 \text{ months}$$

# Modeling probabilities of elapsed times

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

Now consider the variable  $Y$  representing the **elapsed time** between successive arrivals.



# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

Now consider the variable  $Y$  representing the **elapsed time** between successive arrivals.

- What is the probability the time between the third and fourth arrivals is less than  $y$  minutes, for instance?

# Modeling probabilities of elapsed times

Consider the random variable  $X$  which represents the *number of arrivals* at a restaurant within a given time interval.



- The probability of  $X$  in  $t$  time units can be modeled by the Poisson distribution with a rate parameter  $\lambda t$

Now consider the variable  $Y$  representing the **elapsed time** between successive arrivals.

- What is the probability the time between the third and fourth arrivals is less than  $y$  minutes, for instance?
- This is modeled by the **exponential distribution** with parameter  $\lambda$ .

# PDF of exponential distribution

# PDF of exponential distribution

## Definition

# PDF of exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

# PDF of exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) =$$

# PDF of exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) = \lambda e^{-\lambda x}$$

# PDF of exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0 \quad (12)$$




# PDF of exponential distribution

## Definition

A random variable  $X$  that is exponentially distributed with parameter  $\lambda$  has the PDF:

$$f_X(x) = \lambda e^{-\lambda x} \quad x > 0 \quad (12)$$



# CDF of the exponential distribution

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) =$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x)$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$


$$F_X(x) = 1 - e^{-\lambda x}$$

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$






# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$




Note that  $P(X \leq x) = 1 - e^{-\lambda x}$ ,

# CDF of the exponential distribution

The CDF of the exponential distribution is derived as:

$$F_X(x) = P(X \leq x) = \int_0^x \lambda e^{-\lambda t} dt$$

$$F_X(x) = 1 - e^{-\lambda x}$$



Note that  $P(X \leq x) = 1 - e^{-\lambda x}$ , while  $P(X > x) = 1 - (1 - e^{-\lambda x}) = e^{-\lambda x}$

# Mean and variance of the exponential distribution

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X)$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \tag{13}$$

## Variance

The variance of  $X$  is given by:



# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X)$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \tag{13}$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X) =$$

# Mean and variance of the exponential distribution

Let  $X \sim \text{Exponential}(\lambda)$ .

## Mean

The mean of  $X$  is given by:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (13)$$

## Variance

The variance of  $X$  is given by:

$$\mathbb{V}(X) = \frac{1}{\lambda^2} \quad (14)$$

# Example 3: Waiting for a flight

## Example 3: Waiting for a flight

The delay time  $T$  of a flight is exponentially distributed with  $\lambda = 2$  (delays per hour).

# Example 3: Waiting for a flight

- The delay time  $T$  of a flight is exponentially distributed with  $\lambda = 2$  (delays per hour). Answer the following questions:
- (a) What is the mean delay (waiting) time,  $\mathbb{E}(T)$ ?
  - (b) What is the variance of the delay time  $\mathbb{V}(T)$ ?
  - (c) Find the probability that a flight will be delayed by no more than 10 minutes.
  - (d) Assuming you have been waiting for a flight for an hour, what is the probability that the flight will be delayed for an additional 30 minutes? (i.e. Find  $P(T > 1.5 | T > 1)$ ).

## Example 3: Waiting for a flight (cont.)

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by



## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) =$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2}$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2} = \frac{1}{2^2} =$$

## Example 3: Waiting for a flight (cont.)

### Solution

(a) The mean delay is given by

$$\mathbb{E}(T) = \frac{1}{\lambda} = \frac{1}{2} = \boxed{0.5\text{hr}}$$

(b) The variance is:

$$\mathbb{V}(T) = \frac{1}{\lambda^2} = \frac{1}{2^2} = \boxed{0.25\text{hr}^2}$$



## Example 3: Waiting for a flight (cont.)

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right]$$

# Example 3: Waiting for a flight (cont.)

## Solution

(c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] =$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$P\left(T \leq \frac{1}{6}\right) = 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] = 1 - \exp\left[-\frac{1}{3}\right]$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:


$$\begin{aligned} P\left(T \leq \frac{1}{6}\right) &= 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] = 1 - \exp\left[-\frac{1}{3}\right] \\ &= \boxed{0.283} \end{aligned}$$

# Example 3: Waiting for a flight (cont.)

## Solution

- (c) The probability the flight will be delayed by no more than 10 minutes ( $\frac{1}{6}$  hr) is given by:

$$\begin{aligned} P\left(T \leq \frac{1}{6}\right) &= 1 - \exp\left[-\lambda \cdot \frac{1}{6}\right] = 1 - \exp\left[-2\left(\frac{1}{6}\right)\right] = 1 - \exp\left[-\frac{1}{3}\right] \\ &= \boxed{0.283} \end{aligned}$$



## Example 3: Waiting for a flight (cont.)

### Solution

- (c) In python, we can use the `scipy.stats.expon.cdf` function to find  $P(T \leq 1/6)$ :



## Example 3: Waiting for a flight (cont.)

### Solution

(c) In python, we can use the `scipy.stats.expon.cdf` function to find  $P(T \leq 1/6)$ :

---

```
from scipy import stats
import numpy as np
p = stats.expon.cdf(1/6, scale=1/2)
```

---

where:

- $1/6$  is the value at which we want to evaluate the CDF
- $\text{scale} = 1/\lambda = 1/2$  (scale parameter/mean)

This returns  $p = 0.283$ , i.e., about 28.3% probability that the flight will be delayed by no more than 10 minutes.

## Example 3: Waiting for a flight (cont.)

### Solution

(c) In python, we can use the `scipy.stats.expon.cdf` function to find  $P(T \leq 1/6)$ :

---

```
from scipy import stats
import numpy as np
p = stats.expon.cdf(1/6, scale=1/2)
```

---

where:

- $1/6$  is the value at which we want to evaluate the CDF
- $\text{scale} = 1/\lambda = 1/2$  (scale parameter/mean)

This returns  $p = 0.283$ , i.e., about 28.3% probability that the flight will be delayed by no more than 10 minutes.

## Example 3: Waiting for a flight (cont.)

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$P(T > (0.5 + 1) | T > 1) = P(T > 1.5 | T > 1)$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule})
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \end{aligned}$$



## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]}
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\&= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\&= \frac{P(T > 1.5)}{P(T > 1)} \\&= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\&= e^{-2(0.5)} \quad (= P(T > 0.5))\end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\
 &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\
 &= e^{-1}
 \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned} P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\ &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\ &= \frac{P(T > 1.5)}{P(T > 1)} \\ &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\ &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\ &= e^{-1} = \boxed{0.37} \end{aligned}$$

## Example 3: Waiting for a flight (cont.)

### Solution

- (d) The probability that the flight will be delayed by a further 0.5hr after 1hr of waiting is given by:

$$\begin{aligned}
 P(T > (0.5 + 1) | T > 1) &= P(T > 1.5 | T > 1) \\
 &= \frac{P((T > 1.5) \cap (T > 1))}{P(T > 1)} \quad (\text{mult. rule}) \\
 &= \frac{P(T > 1.5)}{P(T > 1)} \\
 &= \frac{e^{-2(1.5)}}{e^{-2(1)}} = e^{-2[1.5-1.0]} \\
 &= e^{-2(0.5)} \quad (= P(T > 0.5)) \\
 &= e^{-1} = \boxed{0.37}
 \end{aligned}$$

In Python: `p = 1 - stats.expon.pdf(.5, scale=1/2)` also returns `p = 0.37`.

# Memorylessness of the exponential distribution

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution



# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

## Memoryless property

$$P(T > t + s | T > s) =$$

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

## Memoryless property

$$P(T > t + s | T > s) = P(T > t) \quad (15)$$

# Memorylessness of the exponential distribution

This leads us to an important property of the exponential distribution

## Memoryless property

$$P(T > t + s | T > s) = P(T > t) \quad (15)$$

That is, it does not matter from which time the waiting begins (i.e. conditioning); the probability of an elapsed time remains the same.

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$   
CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$   
CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) =$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$



# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

PDF:  $f_X(x) =$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

CDF:  $F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

PDF:  $f_X(x) = \lambda e^{-\lambda x},$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) =$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x},$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Mean:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (20)$$

# Recap

- **Lognormal distribution:**  $X \sim \mathcal{LN}(\mu, \sigma^2)$

$$\text{CDF: } F_X(x) = P(X \leq x) = \Phi((\ln(x) - \mu)/\sigma)$$

Mean:

$$\mathbb{E}(X) = e^{(\mu + \frac{1}{2}\sigma^2)} \quad (16)$$

Variance:

$$\mathbb{V}(X) = (e^{\sigma^2} - 1)e^{(2\mu + \sigma^2)} \quad (17)$$

- **Exponential distribution:**  $X \sim \text{Exponential}(\lambda)$

$$\text{PDF: } f_X(x) = \lambda e^{-\lambda x}, \quad x > 0 \quad (18)$$

$$\text{CDF: } F_X(x) = P(X \leq x) = 1 - e^{-\lambda x}, \quad x > 0 \quad (19)$$

Mean:

$$\mathbb{E}(X) = \frac{1}{\lambda} \quad (20)$$

Variance:

$$\mathbb{V}(X) = \frac{1}{\lambda^2} \quad (21)$$